IBUPROFEN SOFTGELS- ibuprofen capsule Breeden Brothers, LLC

Ibuprofen Softgels

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to Pain reliever/ 200 mg ibuprofen (NSAID)* *nonsteroidal anti-inflammatory drug

Purpose

fever reducer (present as the free acid and potassium salt)
Drug Facts (continued)

Uses

- _ temporarily relieves minor aches and pains due to:
- _ muscular aches
- _ the common cold
- _ headache
- _ toothache
- _ backache
- _ menstrual cramps
- _ minor pain of arthritis
- _ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- _blisters
- _ rash
- _ hives
- _ asthma (wheezing)
- _shock
- _ skin reddening
- _ facial swelling

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe

stomach bleeding. The chance is higher if you:

_ take a blood thinning (anticoagulant) or

Drug Facts (continued)

steroid drug

_ are age 60 or older

_ have had stomach ulcers or bleeding problems

_ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- _ have 3 or more alcoholic drinks every day while using this product
- _ take more or for a longer time than directed

Do not use

- _ if you have ever had an allergic reaction to any other pain reliever/fever reducer
- _ right before or after heart surgery

Ask a doctor before use if

- _ stomach bleeding warning applies to you
- _ you have a history of stomach problems, such as heartburn
- _ you are taking a diuretic

Drug Facts (continued)

- _ you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- _ you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- _ taking any other drug
- _ under a doctor's care for any serious condition
- _ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

When using this product

- _ take with food or milk if stomach upset occurs
- _ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

_ you experience any of the following signs of

Drug Facts (continued)

stomach bleeding:

- _ feel faint
- _ have stomach pain that does not get better
- have bloody or black stools
- _ vomit blood
- _ pain gets worse or lasts more than 10 days

- _ fever gets worse or lasts more than 3 days
- _ redness or swelling is present in the painful area
- _ any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (continued)

Directions

_ do not take more than directed

_ the smallest effective dose should be used

- _ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- _ if pain or fever does not respond to 1 capsule, 2 capsules may be used
- _ do not exceed 6 capsules in 24 hours, unless directed by a doctor
- _ children under 12 years: ask a doctor

Other information

- _ each capsule contains: potassium 28 mg
- _ read all warnings and directions before use.
- _ store at 20°-25°C (68°-77°F)
- _ avoid excessive heat above 40°C (104°F)

Drug Facts (continued)

Inactive ingredients

FD&C green #3, gelatin, isopropyl alcohol, medium chain triglycerides, polyethylene glycol, potassium hydroxide, purified water, sorbitol special, and white edible ink

Questions or comments?

1-800-901-2420

Product of China

PRINCIPAL DISPLAY PANEL

b+b better by giving IBUPROFEN SOFTGELS



PRINCIPAL DISPLAY PANEL

b+b better by giving IBUPROFEN SOFTGELS 40 Softgels (200 mg each)



PRINCIPAL DISPLAY PANEL

NDC 70729-509-32 b+b IBUPROFEN SOFTGELS 120 Softgels (200 mg each)



Product Information									
Product T ype		HUMAN OTC DRUG		Item Code (Source)			NDC:70729-000		
Route of Administration		ORAL							
Active Ingredient/Act	ive Moi	atv							
neuve ingreutent/neu		Basis of Strength		Strength					
Ingredient Name IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)					IBUPROFEN		200 mg		
Inactive Ingredients									
		Ingredient Na	ame				Strength		
FD&C GREEN NO. 3 (UNII:		1S)							
GELATIN (UNII: 2G86QN32									
ISOPROPYL ALCOHOL (U									
MEDIUM-CHAIN TRIGLYC									
POLYETHYLENE GLYCOL									
POTASSIUM HYDROXIDE		3C48 M4T)							
WATER (UNII: 059QF0K001 SORBITOL (UNII: 506T60A									
SONDITOL (UNII: SUUTUUA	125K)								
Product Characteristi	ics								
Color	GREEN		Score			no score			
Shape	CAPSUL	E	Size			18 mm			
Flavor			Imprint	Code					
Contains									

# Item Coue		rackage Descript	10 11			Date		Date	
NDC:70729-000-) in 1 BOTTLE, PLASTIC; Type 0: Not a Combination			ion				Dutt	
1 01	roduct				02/01/2018				
/	40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product			ion	0 2/0 1/20 18				
Marketing Info	rmation								
•		n Number er Mener		ion M	[]	ng Ctout Date	Maultad	ing End Data	
Marketing Category	ANDA206568	Application Number or Monograph Citation			лагке и /0 1/20 1	ng Start Date	Market	Marketing End Date	
	1111200300	5		02	10 1/20 1	0			
IBUPROFEN S	OFTGEL	S							
ibuprofen capsule	of follo	5							
Product Informat	ion								
Product Type	e HUMAN OTC DRUG Item C			em Code	e (Sour	ce)	NDC:7072	NDC:70729-509	
Route of Administra									
itoute of Hummistru	lion								
Active Ingredient	/Active Moi	ety							
0		redient Name				Basis of St	Strength Strength		
IBUPROFEN (UNII: WK	-		(YI10 Q M)		IBUPROFEN			200 mg	
Inactive Ingredie	nts								
		Ingredient Na	me				5	Strength	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)									
GELATIN (UNII: 2G860	QN327L)								
ISOPROPYL ALCOHO	DL (UNII: ND2M	416302)							
MEDIUM-CHAIN TRIG	LYCERIDES (U	NII: C9H2L21V7U)							
POLYETHYLENE GLY	COLS (UNII: 3)	WJQ0SDW1A)							
POTASSIUM HYDRO		I3C48M4T)							
WATER (UNII: 059QF0									
SORBITOL (UNII: 506	T60A25R)								
Product Characte	ristics								
Color	GREEN		Score				no score		
Shape	CAPSUI		Size				18 mm		
Flavor			Imprint Co	de					
Contains									
Packaging									
# Item Code		Package Descript	ion		Ma	rketing Start Date	Mar	keting End Date	

1 NDC:70729-509- 32	120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	0 2/0 1/20 18					
Marketing Information							
Marketing Categor	y Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				
ANDA	ANDA206568	02/01/2018					

Labeler - Breeden Brothers, LLC (080131046)

Revised: 1/2018

Breeden Brothers, LLC