UP AND UP CHILDRENS ACETAMINOPHEN- acetaminophen suspension Target Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Target Corporation Children's Acetaminophen Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- · the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- · remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL

36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

^{*}or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions?

Call 1-888-547-7400

Principal Display Panel

Compare to active ingredient in Children's Tylenol®

Oral Suspension

for ages 2 to 11 years

children's acetaminophen 160 mg per 5 mL

oral suspension

fever reducer/pain reliever

ibuprofen free

alcohol free

aspirin free

paraben free

GRAPE FLAVOR

up & up™

DOSAGE CUP INCLUDED

AGES 2-11 YEARS

4 FL OZ (118 mL)



Wamings

warmings: Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes ■ more than 5 doses in 24 hours, which is the maximum

- more than 5 doses in 24 hours, which is the maxim daily amount
 with other drugs containing acetaminophen
 Alleryy alert. Acetaminophen may cause severe skin reactions. Symptoms may include:
 skin reddening
 blisters
 rash
- If a skin reaction occurs, stop use and seek medical help

If a skill reaction of the state of the stat

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor
- or pharmacist.

 if your child has ever had an allergic reaction to this product or any of its ingredients

sk a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
 fever gets worse or lasts more than 3 days
- new symptoms occur redness or swelling is present These could be signs of a serious condition

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical en if you do not notice any signs or symptor

Drug Facts (continued)

- Directions

 This product does not contain directions or complete warnings for adult use warning;

 I do not give more than directed (see overdose warning)

 I shake well before using I'm L = millitier

 Ind right does on chart below. If possible, use weight to does or the price use a contained.

- to dose; otherwise, use age.

 remove the child protective cap and squeeze your child's dose into the dosing cup
 repeat dose every 4 hours while symptoms last

•	than 5 times in 2	4 hours
Weight (lb)	Age (yr)	Dose (mL
under 24	under 2 vears	aek a doct

ı	recigite (ib)	Ago (Ji)	DOSC (IIIL)
Γ	under 24	under 2 years	ask a doctor
	24-35	2-3 years	5 mL
ľ	36-47	4-5 years	7.5 mL
ľ	48-59	6-8 years	10 mL
ľ	60-71	9-10 years	12.5 mL
ľ	72-95	11 years	15 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

- Other information
 each 5 mL contains: sodium 3 mg
 store at 20-25°C (68-77°F)
 do not use if printed neckband is broken or

Inactive ingredients

anhydrus citira acid, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethy cellulose, microcystalline cellulose and carboxymethycellulose sodium, propylene glycel, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions? Call 1-888-547-7400

*This product is not manufactured or distributed Johnson & Johnson Consumer Inc., distributor of Children's Tylenol® Oral Suspension.

Distributed by Target Corporation
Minneapolis, MN 55403

2022 Target Pro-

©2022 Target Brands, Inc. Shop Target.com

children's acetaminophen 160 mg/5 mL oral suspensio

fever reducer/pain reliever



NDC 11673-130-26

Compare to active ingredient in Children's Tylenol® Oral Suspension**

for ages 2 to 11 years

children's acetaminophen 160 mg per 5 mL

oral suspension fever reducer/pain reliever



4 FL 0Z (118 mL)

13026 UW C10



oral suspension fever reducer/pain reliever



DIRECTIONS: STEP 1. Shake well before using




STEP 2. Remove child protective cap



STEP 3. Squeeze your child's dose into the dosing cup



IMPORTANT: Please see dosing chart or ask a doctor for your child's correct dose.





UP AND UP CHILDRENS ACETAMINOPHEN

acetaminophen suspension

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-130	
Route of Administration	ORAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	160 mg in 5 mL		

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WATODDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K6790BS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	

Product Characteristics			
Color	PURPLE (viscous)	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:11673-130- 26	1 in 1 CARTON	07/08/2009		
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product			
2	NDC:11673-130- 28	1 in 1 CARTON	08/30/2012	06/27/2014	
2		148 mL in 1 BOTTLE; Type 0: Not a Combination Product			
3	NDC:11673-130- 34	2 in 1 CARTON	07/08/2009	07/08/2009	
2		118 mL in 1 BOTTLE; Type 0: Not a Combination			

3 P	roduct			
No aleation at the				
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part343	07/08/2009		
Tilliai				

Labeler - Target Corporation (006961700)

Revised: 9/2023 Target Corporation