

## **CHILDRENS PAIN AND FEVER- acetaminophen suspension Best Choice (Valu Merchandisers Company)**

-----

### **Drug Facts**

#### **Active ingredient (in each 5 mL)**

**Acetaminophen 160 mg**

### **Purpose**

**Pain reliever/fever reducer**

### **Uses**

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

### **Warnings**

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this

product

**Ask a doctor before use if your child has**

liver disease.

**Ask a doctor or pharmacist before use if your child is**

taking the blood thinning drug warfarin.

**When using this product,**

**do not exceed recommended dosage (see overdose warning)**

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not take more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (yr)	Dose (tsp or mL)
under 24	under 2	ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 ml
48-59	6-8	10 mL
60-71	9-10	12.5 mL

72-95	11	15 mL
-------	----	-------

### Other information

- **each 5 mL contains:** sodium 2 mg
- store between 20-25°C (68-77°F)
- do not refrigerate
- see bottom panel for lot number and expiration date

### Inactive ingredients

citric acid, butylparaben, carboxymethylcellulose sodium, D&C red #33, FD&C blue #1, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

### Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### Principal Display Panel

Compare to the active ingredient in **Children's Tylenol® Oral Suspension\*\***

#### Children's

#### Pain & Fever

Acetaminophen 160 mg per 5 mL

Oral suspension

- Pain Reliever/Fever reducer
- Alcohol free
- Aspirin free
- Ibuprofen free

Grape flavor

For Ages 2-11

Oral suspension

FL OZ (mL)

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral suspension.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

PROUDLY DISTRIBUTED BY:

ASSOCIATED WHOLESALE GROCERS, INC

KANSAS CITY, KANSAS 66106

Product Label

**Drug Facts (continued)**

- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (year)	Dose (mL)*
under 24	under 2	ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 mL
48-59	6-8	10 mL
60-71	9-10	12.5 mL
72-95	11	15 mL

\*or as directed by a doctor

**Other information**

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F) ■ do not refrigerate
- see bottom panel for lot number and expiration date

**Inactive ingredients** butylparaben, carboxymethylcellulose sodium, citric acid, D&C red #33, FD&C blue #1, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

**Questions or comments?**  
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST.

**Best Choice**

Compare to the active ingredient in Children's Tylenol® Oral Suspension

**Children's Pain Reliever**

**Acetaminophen 160 mg per 5 mL Oral suspension**

**Pain Reliever/Fever Reducer**

- Alcohol free
- Aspirin free
- Ibuprofen free

**Grape Flavor**

For ages 2-11 years 4 FL OZ (118 mL)

**Best Choice**

Compare to the active ingredient in Children's Tylenol® Oral Suspension

**Children's Pain Reliever**

**Acetaminophen 160 mg per 5 mL Oral suspension**

**Pain Reliever/Fever Reducer**

- Alcohol free
- Aspirin free
- Ibuprofen free

**Grape Flavor**

For ages 2-11 years 4 FL OZ (118 mL)

**Drug Facts**

Active ingredient (in each 5 mL)	Purpose
Acetaminophen 160 mg	Pain reliever/fever reducer

**Uses**

- temporarily reduces fever
- temporarily relieves minor aches and pains due to
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dose (see Overdose warning).**

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not take more than directed (see Overdose warning)
- shake well before using
- mL = milliliter

**PROUDLY DISTRIBUTED BY:**  
ASSOCIATED WHOLESALE GROCERS, INC.  
KANSAS CITY, KANSAS 66106

Scan here for more product information.  
Call 1-844-232-1112 for more product information.

**Best Choice,**  
**100% Guaranteed**

www.bestchoicebrand.com

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

BEST CHOICE 58651 0823 PLD

0 70038 58651 7

PLD-H237D FC008405

Lot No.:

Exp. Date:

BEST CHOICE Children's Pain Reliever Grape Flavor

CHILDRENS PAIN AND FEVER

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-288
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	160 mg in 5 mL	
Inactive Ingredients				
Ingredient Name			Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
BUTYLPARABEN (UNII: 3QPI1U3FV8)				
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)				
D&C RED NO. 33 (UNII: 9DBA0SBB0L)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
GLYCERIN (UNII: PDC6A3C0OX)				
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SORBITOL (UNII: 506T60A25R)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
XANTHAN GUM (UNII: TTV12P4NEE)				
Product Characteristics				
Color		Score		
Shape		Size		
Flavor	GRAPE	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-288-04	1 in 1 BOX	02/28/2015	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M013	02/28/2015	

**Labeler** - Best Choice (Valu Merchandisers Company) (868703513)