

ACETAMINOPHEN 500 MG- acetaminophen tablet
Reliable 1 Laboratories LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ACETAMINOPHEN 500 MG TABLET

Drug Facts

Active ingredient (in each tablet)

Acetaminophen USP, 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps

- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

- If a skin reaction occurs, stop use and seek medical help right away.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults & children 12 years and over:

- take 1 tablet every 3-4 hours or 2 tablets every 6 hours while symptoms last
- do not take more than 8 tablets in 24 hours

children under 12 years: do not use

Other information

- store at 15° to 30°C (59° to 86°F)

Inactive ingredients

povidone, pregelatinized corn starch, sodium starch glycolate, stearic acid

Questions or comments?

call **516-341-0666**, 8:30 am - 4:30 pm ET, Monday - Friday

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*Reliable 1 Laboratories LLC is not affiliated with the owner of the trademark **TYLENOL®**.

Distributed by: **Reliable 1 Laboratories LLC, Valley Stream, NY 11580**

www.reliable1labs.com

NDC 69618-011-01

Extra Strength

Acetaminophen USP 500 mg

PAIN RELIEVER

FEVER REDUCER

100 TABLETS

*Compare to Active Ingredient in Extra Strength Tylenol®

NDC 69618-011-01



Extra Strength
Acetaminophen USP
500 mg
PAIN RELIEVER
FEVER REDUCER

100 TABLETS

*Compare to Active Ingredient in Extra Strength TYLENOL®

Drug Facts	Purpose
Active ingredient (in each tablet) Acetaminophen USP, 500 mg Pain reliever/Fever reducer	
Uses	
temporarily relieves minor aches and pains due to: <ul style="list-style-type: none">■ headache■ muscular aches■ backache■ minor pain of arthritis■ the common cold■ toothache■ premenstrual and menstrual cramps temporarily reduces fever	
Warnings	
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take <ul style="list-style-type: none">■ more than 4,000 mg of acetaminophen in 24 hours which is the maximum daily amount■ with other drugs containing acetaminophen■ 3 or more alcoholic drinks every day while using this product Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none">■ skin reddening■ blisters■ rash if a skin reaction occurs, stop use and seek medical help right away.	
	Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
	Ask a doctor before use if you have liver disease

Drug Facts continued on back of label

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*Reliable 1 Laboratories LLC is not affiliated with the owner of the trademark TYLENOL®.

Distributed by: **Reliable 1 Laboratories LLC**
Valley Stream, NY 11580
www.reliable1labs.com



3 69618 01101 4

Rev. 0715

Lot No.:
Exp. Date:

PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
Stop use and ask a doctor if <ul style="list-style-type: none">■ pain gets worse or lasts more than 10 days■ fever gets worse or lasts more than 3 days■ new symptoms occur■ redness or swelling is present These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions ■ do not take more than directed (see overdose warning) adults & children 12 years and over: <ul style="list-style-type: none">■ take 1 tablet every 3-4 hours or 2 tablets every 6 hours while symptoms last■ do not take more than 8 tablets in 24 hours■ children under 12 years: do not use
Other information <ul style="list-style-type: none">■ store at 15° to 30° C (59° to 86° F)
Inactive ingredients novidone, pregelatinized corn starch, sodium starch glycolate, stearic acid
Questions or comments? call 516-341-0666, 8:30 am - 4:30 pm ET, Monday - Friday

ACETAMINOPHEN 500 MG

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69618-011
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	AP;013
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69618-011-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2015	
2	NDC:69618-011-10	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2015	



ACETAMINOPHEN 500mg Tablet

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	11/01/2015	

Labeler - Reliable 1 Laboratories LLC (079718111)

Registrant - Reliable 1 Laboratories LLC (079718111)

Revised: 3/2023

Reliable 1 Laboratories LLC