

HENRY SCHEIN IBUPROFEN- ibuprofen tablet, film coated
Henry Schein Inc.

Henry Schein Ibuprofen

Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

Temporarily relieves minor aches and pains associated with

- headache
- common cold
- toothache
- muscular aches
- backache
- minor arthritis pain
- menstrual cramps

Temporarily reduces fever.

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

■ hives ■ skin reddening ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack or stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint, ■ vomit blood, ■ have bloody or black stools ■ have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new or unexpected symptoms occur

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ **do not use more than directed**

■ **the smallest effective dose should be used**

■ do not take longer than 10 days, unless directed by a doctor (see Warnings)

Adults and children:(12 years and older)

Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used.

Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years:

Ask a doctor

Other information

■ read all product information before using

■ store at 68-77°F (20-25°C)

■ avoid excessive heat 104°F (above40°C)

■ tamper evident sealed packets

■ do not use any opened or torn packets

Inactive ingredients

carnauba wax, corn starch, fumed silica gel, hypromellose, iron oxide red, lactose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? 1-800-472-4346

Henry Schein®

Ibuprofen

NDC #0404-0017-13

For temporary relief of minor aches and pains associated with headache, toothache, backache, menstrual cramps, common cold, muscular aches, minor arthritis pain and for the reduction of fever.

Henry Schein

Seal Of Excellence®

This Package is for Households Without Young Children.

Ibuprofen 200 mg

(NSAID)

250 Packets/2 Tablets

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)
STEARIC ACID (UNII: 4ELV7Z65AP)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;291
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0404-0017-13	250 in 1 BOX	12/15/2015	
1	NDC:0404-0017-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	12/15/2015	

Labeler - Henry Schein Inc. (012430880)

Registrant - Unifirst First Aid Corporation (832947092)