SALICYLIC ACID- liquid corn and callus remover liquid Harris Teeter, LLC

Harris Teeter Liquid Corn & Callus Remover

Active ingredient

Salicylic acid 17%w/w

Purpose

Corn and callus remover

Uses

- for the removal of corns and calluses
- relieves pain by removing corns and calluses

Warnings

For external use only.

Flammable

- keep away from fire or flame
- cap bottle tightly and store at room temperature away heat

Do not use

- on irritated skin
- on any area that is infected or reddened
- if you are a diabetic, or if you have poor blood circulation, except under the advice and supervision of a doctor or podiatrist

When using this product

- avoid contact with eyes
- avoid inhaling vapors
- if product gets into the eyes, flush with water for 15 minutes

Stop and ask a doctor if

discomfort persists, see your doctor or podiatrist

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

• wash affected area and dry thoroughly

- apply one drop at a time with applicator, to sufficiently cover each corn/callus; let dry
- use cushions with hole centered over corn or callus to relieve pressure (additional cushions may be purchased separately)
- repeat this procedure once or twice daily as needed for up to 14 days (until corn/callus is removed)

Other information

store between 20°C to 30°C (68°F to 86°F)

Inactive ingredients

camphor, castro oil, ethanol, ethyl ether, nitrocellulose

Questions?

Call 1-888-423-0139

Principal Display Panel

Harris Teeter Liquid

corn & callus

remover

Salicylic Acid

Safely removes corn and calluses Soft foam cushions and protects 3 Cushions

NET 0.33 FL OZ (9.8 mL)

removal of coms and calluses. Pelieves pain by removing coms and calluses. Warnings. For external use only. Rammable. Keep away from fire or flame. Cap bottle tightly and store at room temperature away from heat. Do not use this product if you are diabetic or have poor blood circulation, except under the advice and supervision of a doctor or podiatrist. Com and callus remover

LIQUID corn & callus remover

Safely removes corns & calluses effectively

CLOSE CAP TIGHTLY AFTER USE

NET 0.31 FL OZ (9 mL)

product avoid contact with eyes, if product gets into the eyes, flush with water for 15 minutes. Avoid inhaling vapors. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Directions: Wash affected area and dry thoroughly. Apply one drop at a time with applicator to sufficiently cover each com/callus. Let not use this product on irritated skin, on any area t is infected or reddened. When using this dry. Use cushions with hole centered over com or callus to relieve pressure. Repeat this procedure once or twice daily as needed for up to 14 days (until corn/callus is removed) **Other Information**: Store at 20° – 25° C (68° – 77° F). Close tightly to prevent solidification. Store in a cool place. **Inactive** Ingredients: camphor, castor oil, ethanol, ethyl ether, nitrocellulose Questions? Call 1-866-964-0939 Keep packaging for complete drug facts Information Salicylic Acid

MATTHEWS, NC 28105

**800-432-6111 or harriste eter.com PROUDLY DISTRIBUTED BY:

MADE IN USA WITH U.S AND IMPORTED PARTS

LIQUID corn & callus remover Salicylic Acid Safely removes corns & calluses Soft foam cushions & protects **FPO Blister and Product** Covers This Area 3 CUSHIONS NET 0.33 FL OZ (9.8 mL)

Drug Facts MPORTANT: Active ingredient Salicylic acid 17% w/w.... Purpose ■ for the removal of corns and calluses ■ relieves pain by removing corns and cal Warnings For external use only Flammable ■ keep away from fire or flame ■ cap bottle tightly and store at room temperature away from heat CLO Do not use on irritated skin on any area that is infected or reddened if you are a diabetic, or if you have poor blood circulation, except under the advice and supervision of a doctor or podiatrist ŠE When using this product avoid contact with eyes avoid inhaling vepors fiproduct gets into the eye, flush with water for 15 min. CAP Stop use and sak a doctor if discomfort persists, see your doctor or podiatrist. Keep out of reach of children. If swallowed, get medical hab or context a Pairse. TIGHTLY wash in a streeted area and dry incroughly apply one drop at a time with applicator to sufficiently cover each com-use oushion with hole centered over com or callus to relieve pressure (additional oushions may be purchased separately) repeat this procedure orice or twice daily as needed for up to 14 days (until com/callus is removed). AFTER Other information store between 20°-25°C (68°-77°F) close tightly to prevent solidification store in a cool place Inactive ingredients camphor, castor oil, ethanol, ethyl ether, nitrocellulose Questions? Call 1-866-964-0939 ■ Wipe top of bottle clean after each use *This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Dr. Scholl's^o. PROUDLY DISTRIBUTED BY: HARRIS TEETER, LLC Matthews, NC 28105 1-800-432-6111 or harristeeter.com MADE IN USA WITH U.S. AND IMPORTED PARTS 41-030HT-01

SALICYLIC ACID

liquid corn and callus remover liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69256-036
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	0.17 mg in 9.8 mL

Inactive Ingredients		
Ingredient Name	Strength	
PYROXYLIN (UNII: KYR8BR2X6O)		
CAMPHOR (NATURAL) (UNII: N20HL7Q941)		
CASTOR OIL (UNII: D5340Y2I9G)		

P	ackaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:69256- 036-09	9.8 mL in 1 PACKAGE; Type 0: Not a Combination Product	12/21/2017		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M030	12/21/2017		

Labeler - Harris Teeter, LLC (048463103)

Revised: 1/2024 Harris Teeter, LLC