

**NYSTATIN- nystatin suspension**  
**PAI Holdings, LLC**


-----  
**Nystatin Oral Suspension USP**  
**[100,000 units per mL]**

**Rx Only**

**DESCRIPTION**

Nystatin is an antimycotic polyene antibiotic obtained from *Streptomyces noursei*.

Structural formula:



C<sub>47</sub>H<sub>75</sub>NO<sub>17</sub>    MW 926.13

Nystatin Oral Suspension USP, for oral administration, contains 100,000 USP Nystatin Units per mL. Inactive ingredients: alcohol ( $\leq 1\%$  v/v), artificial peppermint flavor, cherry flavor, citric acid, D&C Yellow No. 10, FD&C Red No. 40, glycerin, magnesium aluminum silicate, methylparaben, potassium phosphate dibasic, propylene glycol, propylparaben, purified water and sucrose.

**CLINICAL PHARMACOLOGY**

**Pharmacokinetics**

Gastrointestinal absorption of nystatin is insignificant. Most orally administered nystatin is passed unchanged in the stool. In patients with renal insufficiency receiving oral therapy with conventional dosage forms, significant plasma concentrations of nystatin may occasionally occur.

**Microbiology**

Nystatin is both fungistatic and fungicidal *in vitro* against a wide variety of yeasts and yeast-like fungi. *Candida albicans* demonstrates no significant resistance to nystatin *in vitro* on repeated subculture in increasing levels of nystatin; other *Candida* species become quite resistant. Generally, resistance does not develop *in vivo*. Nystatin acts by

binding to sterols in the cell membrane of susceptible *Candida* species with a resultant change in membrane permeability allowing leakage of intracellular components. Nystatin exhibits no appreciable activity against bacteria, protozoa, or viruses.

## **INDICATIONS AND USAGE**

Nystatin Oral Suspension is indicated for the treatment of candidiasis in the oral cavity.

## **CONTRAINDICATIONS**

The preparation is contraindicated in patients with a history of hypersensitivity to any of its components.

## **PRECAUTIONS**

### **General**

This medication is not to be used for the treatment of systemic mycoses. Discontinue treatment if sensitization or irritation is reported during use.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

No long-term animal studies have been performed to evaluate carcinogenic potential. There also have been no studies to determine mutagenicity or whether this medication affects fertility in males or females.

### **Pregnancy**

#### **Teratogenic Effects**

Animal reproduction studies have not been conducted with nystatin oral suspension. It is also not known whether nystatin oral suspension can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Nystatin oral suspension should be given to a pregnant woman only if clearly needed.

### **Nursing Mothers**

It is not known whether nystatin is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when nystatin is administered to a nursing woman.

### **Pediatric Use**

See **DOSAGE AND ADMINISTRATION**.

## **ADVERSE REACTIONS**

Nystatin is well tolerated even with prolonged therapy. Oral irritation and sensitization have been reported. (See **PRECAUTIONS: General**).

*Gastrointestinal:* Diarrhea (including one case of bloody diarrhea), nausea, vomiting,

gastrointestinal upset/disturbances.

*Dermatologic:* Rash, including urticaria has been reported rarely. Stevens-Johnson syndrome has been reported very rarely.

*Other:* Tachycardia, bronchospasm, facial swelling, and non-specific myalgia have also been rarely reported.

**To report SUSPECTED ADVERSE REACTIONS, contact Pharmaceutical Associates, Inc. at 1-800-845-8210 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## **OVERDOSAGE**

Oral doses of nystatin in excess of five million units daily have caused nausea and gastrointestinal upset. There have been no reports of serious toxic effects of superinfections (See **CLINICAL PHARMACOLOGY, Pharmacokinetics**).

## **DOSAGE AND ADMINISTRATION**

**INFANTS:** 2 mL (200,000 units) four times daily (in infants and young children, use dropper to place one-half of dose in each side of mouth and avoid feeding for 5 to 10 minutes).

**NOTE:** Limited clinical studies in premature and low birth weight infants indicate that 1 mL four times daily is effective.

**CHILDREN AND ADULTS:** 4 to 6 mL (400,000 to 600,000 units) four times daily (one-half of dose in each side of mouth). The preparation should be retained in the mouth as long as possible before swallowing.

Continue treatment for at least 48 hours after perioral symptoms have disappeared and cultures demonstrate eradication of *Candida albicans*.

## **HOW SUPPLIED**

Nystatin Oral Suspension USP, 100,000 USP Nystatin Units per mL, is available in a cherry, peppermint flavored, light creamy yellow, ready-to-use suspension, supplied in the following oral dosage forms:

NDC 0121-0868-02: 2 fl oz (60mL) bottle with calibrated dropper

NDC 0121-0868-16: 16 fl oz (473mL) bottle

NDC 0121-0868-05: 5mL unit dose cup. Case contains 40 unit dose cups of 5 mL (NDC 0121-0868-40) packaged in 4 trays of 10 unit dose cups each, 50 unit dose cups of 5 mL (NDC 0121-0868-50) packaged in 5 trays of 10 unit dose cups each and 100 unit dose cups of 5 mL (NDC 0121-0868-00) packaged in 10 trays of 10 unit dose cups each.

## **Storage**

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Avoid freezing.

**R02/22**

***Pharmaceutical Associates, Inc.***

Greenville, SC 29605

www.paipharma.com

**PRINCIPAL DISPLAY PANEL - 60 mL Bottle Label**

**60 mL**

**NDC 0121-0868-02**

**Nystatin Oral Suspension, USP**

**100,000 units per mL**

**SHAKE WELL BEFORE USING**

**Packaged with Calibrated Dropper**

**DO NOT USE IF TAMPER EVIDENT  
SEAL IS BROKEN OR MISSING.**

**Rx ONLY**

Pharmaceutical Associates, Inc.

Greenville, SC 29605

Each mL of cherry, peppermint flavored oral suspension contains 100,000 units Nystatin, USP and alcohol ( $\leq 1\%$  v/v).

**USUAL DOSAGE:** For Infants: 2 mL(200,000 units) four times daily (in infants and young children, use dropper to place one half of dose in each side of the mouth and avoid feeding for 5 to 10 minutes). See insert.

**WARNINGS:** Keep this and all drugs out of reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.


**Keep tightly closed. Protect from light.**

**Store at 20° to 25°C (68° to 77°F)**

**[See USP Controlled Room Temperature].**

**AVOID FREEZING**

X0868020120      R01/20




60 mL    NDC 0121-0868-02

## Nystatin Oral Suspension, USP

**100,000 units per mL**

SHAKE WELL BEFORE USING  
Packaged with Calibrated Dropper  
DO NOT USE IF TAMPER EVIDENT  
SEAL IS BROKEN OR MISSING.

**Rx ONLY**



**Pharmaceutical  
Associates, Inc.**  
Greenville, SC 29605

Each mL of cherry, peppermint flavored oral suspension contains 100,000 units Nystatin, USP and alcohol ( $\leq 1\%$  v/v).

**Usual Dosage:** For infants: 2 mL (200,000 units) four times daily (in infants and young children, use dropper to place one-half of dose in each side of mouth, and avoid feeding for 5 to 10 minutes). See insert.

**Warnings:** Keep this and all drugs out of reach of children. In case of accidental overdose, seek professional assistance or contact Poison Control Center immediately.

**Keep tightly closed. Protect from light.**

**Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature].**

**AVOID FREEZING.**

X0868020120

R01/20

### PRINCIPAL DISPLAY PANEL - 5 mL Unit Dose Cup Label

Delivers **5 mL**

**NDC 0121-0868-05**

**NYSTATIN ORAL   SUSPENSION, USP**

**500,000 units/5 mL**

Alcohol  $\leq 1\%$  v/v    **SHAKE WELL**

**Package Not Child-Resistant**

**Rx ONLY**

PHARMACEUTICAL ASSOCIATES, INC.,  
GREENVILLE, SC 29605

**SEE INSERT**

A0868051021



NYSTATIN

nystatin suspension

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0121-0868
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
NYSTATIN (UNII: BDF1O1C72E) (NYSTATIN - UNII:BDF1O1C72E)	NYSTATIN	100000 [USP'U] in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SUCROSE (UNII: C151H8M554)	
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)	

Product Characteristics			
Color	yellow (Light - Creamy)	Score	
Shape		Size	

Flavor		CHERRY (w/Peppermint)	Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0121-0868-02	60 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product	07/17/2020	
2	NDC:0121-0868-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/17/2020	
3	NDC:0121-0868-40	4 in 1 CASE	07/17/2020	
3		10 in 1 TRAY		
3	NDC:0121-0868-05	5 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		
4	NDC:0121-0868-50	5 in 1 CASE	07/17/2020	
4		10 in 1 TRAY		
4	NDC:0121-0868-05	5 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		
5	NDC:0121-0868-00	10 in 1 CASE	07/17/2020	
5		10 in 1 TRAY		
5	NDC:0121-0868-05	5 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA203621	07/17/2020	

**Labeler -** PAI Holdings, LLC (044940096)

## Establishment

Name	Address	ID/FEI	Business Operations
PAI Holdings, LLC dba Pharmaceutical Associates, Inc.		097630693	analysis(0121-0868) , label(0121-0868) , manufacture(0121-0868) , pack(0121-0868)

Revised: 3/2023

PAI Holdings, LLC