

**MEDIQUE IPRIN- ibuprofen tablet, film coated**  
**DOVER ADDAPRIN- ibuprofen tablet, film coated**  
**MEDI-FIRST IBUPROFEN- ibuprofen tablet, film coated**  
**MEDI-FIRST PLUS IBUPROFEN- ibuprofen tablet, film coated**  
**OTIS CLAPP ULTRAPRIN- ibuprofen tablet, film coated**  
**Unifirst First Aid Corporation**

-----

## **UniFirst First Aid Ibuprofen**

### ***Drug Facts***

#### ***Active ingredient***

Ibuprofen 200 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

#### ***Purpose***

Pain reliever/fever reducer

#### ***Uses***

Temporarily relieves minor aches and pains associated with

■ headache ■ toothache ■ backache ■ menstrual cramps

■ common cold ■ muscular aches ■ minor arthritis pain

Temporarily reduces fever.

#### ***Warnings***

**Allergy Alert :** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives ■ skin reddening ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

■ are age 60 or older

■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

**Heart attack or stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new or unexpected symptoms occur

**If pregnant or breast feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of Children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## ***Directions***

- **do not use more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

### **Adults and children:(12 years and older)**

Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used.

Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

### **Children under 12 years:**

Ask a doctor

## ***Other information***

- read all product information before using
- store at 68-77°F (20-25°C)
- avoid excessive heat 104°F (above 40°C)
- tamper evident sealed packets
- do not use any opened or torn packets

## ***Inactive ingredients***

carnauba wax\*, corn starch, hypromellose\*, iron oxide red, lactose\*, magnesium stearate\*, microcrystalline cellulose\*, polydextrose\*, polyethylene glycol, polyvinyl alcohol\*, povidone K30\*, silicon dioxide, sodium starch glycolate, stearic acid, talc\*, titanium dioxide

***\*may contain***

***Questions or comments?*** 1-800-634-7680

## **Medique Iprin Label**

Medique® I-Prin

Ibuprofen 200 mg

Anti-inflammatory (NSAID)

This Package is for Households without Young Children.

Pain Reliever/Fever Reducer • Ibuprofen 200 mg

24 Tablets (12 x 2)

Tamper Evident Unit Dose Packets



## Medi-First Ibuprofen Label

Medi-First®

Ibuprofen 200 mg

100 tablets (50 x 2)

Pain Reliever/Fever Reducer

Aches, Fever • Ibuprofen (NSAID) 200mg


Pull to Open

Compare active ingredient to: Advil®

Registered Trademark of Pfizer Consumer Healthcare

This Package is for Households without Young Children.

Tamper Evident Unit Dose Packets



## Medi-First Plus Ibuprofen Label

Medi-First® Plus

Ibuprofen

250 tablets (125 x 2's)

This Package is for Households without Young Children

Pull to Open

Ibuprofen 200 mg (NSAID)

Pain Reliever/Fever Reducer

Compare active ingredient to:

Advil®

Registered Trademark of Pfizer Consumer Healthcare

Tamper Evident Unit Dose Packets

**Drug Facts**

**Active ingredient (in each tablet)** Purpose  
Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses**  
Temporarily relieves minor aches and pains associated with:  
■ headache ■ backache ■ muscle aches ■ menstrual cramps  
■ common cold ■ minor arthritis pain  
Temporarily reduces fever.

**Warnings**  
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives ■ skin redness ■ asthma (wheezing)  
■ facial swelling ■ rash ■ shock ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.  
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
NSAIDs (aspirin, ibuprofen, naproxen, or others)

**Drug Facts (continued)**  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed  
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
Do not use:  
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ right before or after heart surgery  
Ask a doctor before use if:  
■ you have problems or serious side effects from taking pain relievers or fever reducers  
■ stomach bleeding warning applies to you  
■ you have a history of stomach problems such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke  
■ you are taking a diuretic  
Ask a doctor or pharmacist before use if you are:  
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin  
■ under a doctor's care for any serious condition  
■ taking any other drug (continued on opposite panel)

**When using this product**  
■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**  
■ you experience any of the following signs of stomach bleeding:  
■ red, dark, or bloody stools  
■ have bloody or black stools  
■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke  
■ chest pain ■ trouble breathing  
■ weakness in one part or side of body  
■ slurred speech ■ leg swelling  
■ skin gets worse or lasts for more than 10 days  
■ fever gets worse or lasts for more than 3 days  
■ redness or swelling is present in the painful area  
■ any new or unexpected symptoms occur

**Drug Facts (continued)**  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.  
**Directions**  
■ do not use more than directed  
■ the smallest effective dose should be used  
■ do not take longer than 10 days, unless directed by a doctor (see Warnings)  
**Adults and children (12 years and older)**  
Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.  
**Children under 12 years:**  
Ask a doctor  
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Other information**  
■ read all product information before using  
■ store at 68-77°F (20-25°C)  
■ avoid excessive heat 104°F (above 40°C)  
■ tamper evident sealed packets  
■ do not use any opened or torn packets (continued on opposite panel)

**Drug Facts (continued)**  
**Inactive ingredients**  
camphor wax\*, corn starch, hypromellose\*, iron oxide red, lactose\*, magnesium stearate\*, microcrystalline cellulose\*, polydextrose\*, polyethylene glycol, povidone (K-30)\*, silicon dioxide, sodium starch glycolate, stearic acid, talc\*, titanium dioxide  
\*may contain  
**Questions or comments?** 1-800-434-7680

**Compare active ingredient to:**  
Advil®  
Registered Trademark of Pfizer Consumer Healthcare  
Tamper Evident Unit Dose Packets

**RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION**

**Barcode:** 3 47682 90833

**Batch #0833**

**Manufactured by Medi-First Plus**  
Fort Myers, FL 33947 USA  
1-800-434-7680  
www.medi-qproducts.com

Dover Addaprin Label

Dover Addaprin™

Pain Reliever-Fever Reducer

Ibuprofen 200 mg Tablets (NSAID)

This Package is for Households without Young Children.

Dover Pharmaceutical

Products of the highest quality and effectiveness

Tamper Evident

Sealed Packets

Unit Dose Packs

500 Tablets

(250 Packets of 2)

All Dover Pharmaceutical formulas conform to federal regulations

Antihistamine free

No danger of drowsiness

Sugar Free

for safer use by diabetics

Salt free

Minimizes high blood pressure

Caffeine Free

Avoides over stimulation

**Otis Clapp Ultraprin Label**

OC Otis Clapp

Quality & Integrity Since 1840

Ultraprin<sup>™</sup>

Pain Reliever-Fever Reducer (NSAID)

Ibuprofen Tablets USP 200 mg

For Deep Seated Pain

See Warnings and Directions on Side Panel

Tear Out Along Perforation To Dispense

Professional Healthcare

500 Tablets (250 Packets of 2)

**MEDIQUE IPRIN**  
ibuprofen tablet, film coated

Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	NDC:47682-600
Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)			IBUPROFEN	200 mg
Inactive Ingredients				
Ingredient Name				Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)				
MAGNESIUM STEARATE (UNII: 70097M6I3O)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
STARCH, CORN (UNII: O8232NY3SJ)				
HYPROMELLOSES (UNII: 3NXW29V3WO)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
FERRIC OXIDE RED (UNII: 1K09F3G675)				
LACTOSE (UNII: J2B2A4N98G)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
POLYDEXTROSE (UNII: VH2XOU12IE)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
Product Characteristics				
Color	brown	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;291	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-600-13	250 in 1 BOX	01/26/2017	
1	NDC:47682-600-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-600-47	100 in 1 BOX	01/26/2017	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-600-64	12 in 1 BOX	01/26/2017	
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-600-	2 in 1 BOX	01/26/2017	







Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-614-13	250 in 1 BOX	01/26/2017	
1	NDC:47682-614-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	02/01/2016	

MEDI-FIRST IBUPROFEN			
ibuprofen tablet, film coated			

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-608
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients	
Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	brown	Score	no score

Shape		ROUND	Size	10mm
Flavor			Imprint Code	44;291
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-608-30	4 in 1 BOX	01/26/2017	
1	NDC:47682-608-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-608-33	50 in 1 BOX	01/26/2017	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-608-48	125 in 1 BOX	01/26/2017	
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-608-13	250 in 1 BOX	01/26/2017	
4		2 in 1 PACKET; Type 0: Not a Combination Product		
5	NDC:47682-608-50	25 in 1 BOX	04/01/2019	
5		2 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA075010	02/01/2016	

<b>MEDI-FIRST PLUS IBUPROFEN</b>			
ibuprofen tablet, film coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-609
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg
<b>Inactive Ingredients</b>			

Ingredient Name	Strength
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>LACTOSE</b> (UNII: J2B2A4N98G)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I3O)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;291
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-609-13	250 in 1 BOX	01/26/2017	01/27/2017
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-609-33	50 in 1 BOX	01/26/2017	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-609-48	125 in 1 BOX	01/26/2017	
3		2 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	02/01/2016	

## OTIS CLAPP ULTRAPRIN

ibuprofen tablet, film coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-602	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg	
Inactive Ingredients				
Ingredient Name			Strength	
HYPROMELLOSES (UNII: 3NXW29V3WO)				
LACTOSE (UNII: J2B2A4N98G)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
POLYDEXTROSE (UNII: VH2XOU12IE)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
CARNAUBA WAX (UNII: R12CBM0EIZ)				
STARCH, CORN (UNII: O8232NY3SJ)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
FERRIC OXIDE RED (UNII: 1K09F3G675)				
Product Characteristics				
Color	brown	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;291	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-602-13	250 in 1 BOX	02/01/2016	04/03/2017
1	NDC:47682-602-99	2 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA075010		02/01/2016	04/03/2017

Revised: 3/2024

Unifirst First Aid Corporation