FAMOTIDINE- famotidine tablet, film coated Select Brand

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Famotidine USP, 10 mg

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- with other acid reducers
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness**, **sweating**, or **dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

OTHER INFORMATION

- store at 20° to 25°C (68° to 77°F)
- protect from moisture
- read the directions and warnings before use
- keep the carton. It contains important information.
- TAMPER EVIDENT: DO NOT USE IF THE CARTON OR INDIVIDUAL BLISTER UNIT IS OPEN OR TORN.

INACTIVE INGREDIENTS

Colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide

QUESTIONS?

Call 1-800-406-7984

PATIENT INFORMATION

- 1 tablet relieves heartburn due to acid indigestion
- Famotidine prevents heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

PRINCIPAL DISPLAY PANEL

select brand_® NDC 15127-370-12 Original Strength Acid Controller Famotidine Tablets USP, 10 mg

Acid Reducer

 $^{\dagger}\text{Compare}$ to the active ingredient of Original Strength Pepcid AC $^{\circledast}$

Just One Tablet

Prevents & Relieves Heartburn Due to Acid Indigestion

12 Tablets

Distributed by: SELECT BRAND®DISTRIBUTORS

5102118/R0413



FAMOTIDINE								
famotidine tablet, film coated								
Product Information								
Product T ype	HUMAN OTC DRUG	Item Code (Source)		NDC:15127-370				
Route of Administration	ORAL							
A - T - 1 / A - T	· .							
Active Ingredient/Active Moiety								
Ingredient Name			Basis of Strength		Strength			
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)			FAMOTIDINE		10 mg			

	Strength					
COLLOIDAL SILICON						
HYDRO XYPRO PYL CEI	L ULOSE (UNII: RFW2E	T671P)				
HYPROMELLOSES (UN	II: 3NXW29V3WO)					
FERRIC OXIDE RED (UN	NII: 1K09F3G675)					
MAGNESIUM STEARAT	E (UNII: 70097M6I30)					
CELLULOSE, MICROCI	RYSTALLINE (UNII: OPI	IR32D61U)				
POLYETHYLENE GLYC	COL 400 (UNII: B697894	4SGQ)				
STARCH, PREGELATIN	IZED CORN (UNII: 0823	32NY3SJ)				
TALC (UNII: 7SEV7J4R10	J)					
TITANIUM DIO XIDE (UI	NII: 15FIX9V2JP)					
Product Character	istics					
Color	pink	pink Score			no score	
Shape	ROUND	Size	Size		8 mm	
Flavor		Imprint Code	Imprint Code		035	
Contains						
Packaging						
# Item Code	Package Desc	ription Marketi	Marketing Start Date		Marketing End Date	
1 NDC:15127-370-12	1 in 1 CARTON					
1	12 in 1 BLISTER PAC	CK				
2 NDC:15127-370-30	1 in 1 CARTON					
2	30 in 1 BLISTER PAG	CK				
Markating Info	mation					
Marketing Info					Marketing End Date	
Marketing Category		plication Number or Monograph Citation		Marketing Start Date		
ANDA	ANDA090283		02/13/2012			

Labeler - Select Brand (043562370)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	manufacture(15127-370)

Revised: 7/2013