# MILK OF MAGNESIA- magnesium hydroxide liquid H E B

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

## **HEB Milk of Magnesia Drug Facts**

## Active ingredient (in each 15 mL TBSP)

Magnesium hydroxide 1200 mg

## Purpose

Saline laxative

#### Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

## Warnings

## Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

## Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

## Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

### If pregnant or breast-feeding,

ask a health professional before use.

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### Directions

shake well before use

- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- TBSP = tablespoonful

adults and children 12 years and older	2 to 4 TBSP.
children 6 to 11 years	1 to 2 TBSP.
children under 6 years	ask a doctor

### Other information

- each tablespoon (15 mL) contains: calcium 20 mg and magnesium 595 mg
- does not meet USP requirements for preservative effectiveness
- store at 20°-25°C (68°-77°F)
- keep tightly closed
- avoid freezing

## **Inactive ingredients**

purified water

## Questions or comments?

1-800-719-9260

## **Principal Display Panel**

Compare to Phillips'® Milk of Magnesia Original active ingredient

MILK OF MAGNESIA

Magnesium Hydroxide

Saline Laxative

Comfortable, Cramp Free Relief

Stimulant Free

Sugar Free

Original Flavor

**GLUTEN FREE** 

# Compare to Phillips'® Milk of Magnesia Original active ingredient\*

NDC 37808-396-40



# MILK OF MAGNESIA

# Magnesium Hydroxide

# Saline Laxative

- Comfortable, Cramp Free Relief
- Stimulant Free
- Sugar Free

**Original Flavor** 

GLUTEN FREE

12 FL OZ (355 ml)

: 39640 LJ F4

# **Drug Facts**

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.

# Active ingredient (in each 15 mL TBSP)

Purpose

Magnesium hydroxide 1200 mg......Saline laxative

## Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

# Warnings

# Ask a doctor before use if you have

- kidney disease
   a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

## Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## **Directions**

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
   TBSP = tablespoonful

adults and children 12 years and older	2 to 4 TBSP.	
children 6 to 11 years	1 to 2 TBSP.	
children under 6 years	ask a doctor	

## Other information

- each tablespoon (15 mL) contains: calcium 20 mg and magnesium 595 mg
- does not meet USP requirements for preservative effectiveness
- store at 20°-25°C (68°-77°F)
   keep tightly closed
   avoid freezing

# Inactive ingredient purified water

# (Questions or comments? 1-800-719-9260

\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips'® Milk of Magnesia Original.

# E8 rt 0h96E

## SAN ANTONIO, TX 78204 GUARANTEE

We believe the high quality of this H-E-B product makes it an outstanding value. We hope you'll agree. If not, we'll cheerfully refund your money. Thanks for shopping with us.



## **MILK OF MAGNESIA**

magnesium hydroxide liquid

P	ro	duct	Info	rmatio	n

Product Type HUMAN OTC DRUG LABEL Item Cod

Item Code (Source)

NDC:37808-396

**Route of Administration** 

ORAL

**DEA Schedule** 

## **Active Ingredient/Active Moiety**

Ingredient Name

MAGNESIUM HYDRO XIDE (MAGNESIUM CATION)

M

Basis of Strength Strength

MAGNESIUM HYDROXIDE 1200 mg in 15 mL

## **Inactive Ingredients**

**Ingredient Name** 

Strength

WATER

## **Product Characteristics**

ColorWHITE (opaque)ScoreShapeSizeFlavorImprin

ize

Imprint Code

Dackaging

Contains

1	1 ackaging			
#	Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date
1	NDC:37808-396-40	355 mL in 1 BOTTLE		

## **Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	06/09/1992	

# **Labeler** - HEB (007924756)

Revised: 10/2012 HE B