LAXATIVE ENTERIC COATED- bisacodyl tablet, coated P & L Development, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Bisacodyl Enteric Coated Tablet

Active ingredient (in each tablet)

Bisacodyl, USP 5 mg

Purpose

Stimulant laxative

Use

- for temporary relief of occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts for more than 2 weeks

When using this product

- do not chew or crush tablet(s)
- it may cause stomach discomfort, faintness and cramps
- do not use within 1 hour after taking an antacid or milk

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

take with a glass of water

adults and children 12 years of age and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	take 1 tablet in a single dose
children under 6 years of age	ask a doctor

Other information

• store between 20-25°C (68-77°F)

Inactive ingredients

acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C Yellow #10 Aluminum Lake, FD&C Yellow #6 Aluminum Lake, gelatin, iron oxide, iron oxide Black, iron oxide Yellow (iron oxide ochre), magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, and titanium dioxide

Questions or Comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient in DULCOLAX®

LAXATIVE

Bisacodyl, 5 mg stimulant laxative

Gentle, Predictable, Overnight relief

*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of Dulcolax®Lasative.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Distributed by:

PL Developments

200 Hicks St Westbury, NY 11590 USA

Package Label



Drug Facts Active ingredient Purpose (in each tablet) Bsacody (USF 5 mg	product generally produces a bowel movement in 6 to 12 hours movement in 6 to 12 hours bo not use if you cannot swallow without chewing. Ask a doctor before use if you have a smorab pain, nauses or vomitting a smorab pain, nauses or vomitting a sudden change in bowel habits that lasts more than 2 weeks.	When using this product	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overclose, get medical help or contact a Poison Control Center right away.
Drug Facts Active ingredient (in each tablet) Bisacody USP 5 mgSI USE = tor temporary relie	product generally produces a bowel movement in 6 to 12 hours Warnings Do not use if you cannot swallow with chewing. Ask a doctor before use if you have a stoden change in bowel habits the more than 2 weeks.	When using this product or crush tablet(s) or crush tablet(s) or crush tablet(s) or or an antacid or may cause stomach discomfor and cramps Stop use and ask a doctor if rectal bleeding or no bowel mo using this product. These could serious condition. Stop used serious condition.	If pregnant or br professional befor of children. In ca help or contact a away.

Readyincase Laxative Bisacodyl

LAXATIVE ENTERIC COATED

bisacodyl tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59726-190
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg	

Inactive Ingredients			
Ingredient Name	Strength		
ACACIA (UNII: 5C5403N260)			
CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H)			
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)			
CARNAUBA WAX (UNII: R12CBM0EIZ)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
STARCH, CORN (UNII: O8232NY3SJ)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			

FD&C YELLOW NO. 6 (UNII: H77VEI93A8) **ALUMINUM OXIDE (UNII: LMI2606933)** GELATIN (UNII: 2G86QN327L) FERRIC OXIDE RED (UNII: 1K09F3G675) FERROSOFERRIC OXIDE (UNII: XM0M87F357) FERRIC OXIDE YELLOW (UNII: EX43802MRT) MAGNESIUM STEARATE (UNII: 70097M6I30) CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ) **Polyvinyl Acetate Phthalate** (UNII: 58QVG85GW3) POVIDONE (UNII: FZ 989GH94E) **SHELLAC** (UNII: 46N107B710) **SODIUM STARCH GLYCOLATE TYPE A CORN** (UNII: AG9B65PV6B) STEARIC ACID (UNII: 4ELV7Z65AP) TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics			
Color	orange	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	TCL;003
Contains			

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:59726- 190-03	1 in 1 CARTON	09/01/2010	12/27/2024		
1		25 in 1 BLISTER PACK; Type 0: Not a Combination Product				

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	09/01/2010	12/27/2024

Labeler - P & L Development, LLC (800014821)

Revised: 3/2023 P & L Development, LLC