

**FAMILY CARE COLD AND FLU MULTI SYMPTOM- acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride liquid  
United Exchange Corp.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

Active ingredients (in each 15 mL dose cup)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 (30 mL each) doses in 24 hours, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Sore throat warning:** if sore throat is severe, lasts more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

☐ **Ask a doctor or pharmacist before use if you are** ☐ taking the blood thinning drug warfarin

☐ **When using this product**

- do not use more than directed

☐ **Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

☐ **If pregnant or breast-feeding,** ☐ ask a health professional before use.

☐ **Keep out of reach of children. Overdose warning:** ☐ Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

☐ **Directions**

- take only as directed - see Overdose warning
- use dose cup or tablespoon (TBSP)
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs over: 30 mL (2TBSP) every 4 hours

children 6 to under 12 yrs: 15 mL (1 TBSP) every 4 hours

children 4 to under 6 yrs: ask a doctor

children under 4 yrs: do not use ☐

☐ **Other information**

- each 15 mL dose cup contains: sodium 25 mg
- store at room temperature

Inactive ingredients

citric acid hydrate, D-sorbitol, FD&C Yellow No. 6, glycerin, L-menthol, methylparaben, polyethylene glycol, propylene glycol, propylparaben, purified water, sodium chloride, sodium citrate hydrate, sodium saccharin, xanthan gum

Distributed by:

United Exchange Corp.

17211 Valley View Ave.

Cerritos, CA 90703

Made in Korea

TWIST TO OPEN

**Drug Facts (continued)**

• a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis or emphysema • trouble urinating due to enlarged prostate gland • a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**When using this product • do not use more than directed**

**Stop use and ask a doctor if •** you get nervous, dizzy or sleepless • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** **Overdose warning:** Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

**Directions •** take only as directed - see Overdose warning • use dose cup or tablespoon (TBS)P) • do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL (2 TBS)P) every 4 hrs
children 6 to under 12 yrs	15 mL (1 TBS)P) every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

**Other information •** each 15 mL dose cup contains: sodium 25 mg • store at room temperature

**Inactive ingredients** citric acid hydrate, D-sorbitol, FD&C Yellow No.6, glycerin, L-menthol, methylparaben, polyethylene glycol, propylene glycol, propylparaben, purified water, sodium chloride, sodium citrate hydrate, sodium saccharin, xanthan gum

**TAMPER EVIDENT:** Do not use if printed shrinkband is missing or broken. Failure to follow these warnings could result in serious consequences.

\*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Cold & Flu.

\*Compare to the active ingredients in VICKS® DAYQUIL® Cold & Flu

**Drug Facts**

**Active ingredients (in each 15 mL dose cup) Purpose**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Phenylephrine HCl 5 mg.....Nasal decongestant

**Uses** temporarily relieves common cold/flu symptoms: • nasal congestion • cough due to minor throat and bronchial irritation • sore throat • headache • minor aches and pains • fever

**Warnings Liver warning:** This product contains acetaminophen. Severe liver damage may occur if • adult takes more than 4 (30 mL each) doses in 24 hours, which is the maximum daily amount for this product • child takes more than 4 doses (15 mL each) in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks daily while using this product **Sore throat warning:** If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use •** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have •** liver disease • heart disease • high blood pressure • thyroid disease • diabetes • cough that occurs with too much phlegm (mucus)

Dist. by: United Exchange Corp., Cerritos, CA 90703 USA. Made in Korea. 7 80707 00627 6

## FAMILY CARE COLD AND FLU MULTI SYMPTOM

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:65923-627
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	

GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL (UNII: L7T10EIP3A)	
METHYLPARABEN (UNII: A2I8C7H9T)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0K00R)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
XANTHAN GUM (UNII: TTV12P4NEE)	

<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65923-627-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/23/2015	

**Labeler** - United Exchange Corp. (840130579)