

OXYGEN - oxygen gas
Ameri-Quipt of North Carolina, Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

OXYGEN COMPRESSED LABEL

OXYGEN COMPRESSED USP UN1072 C.A.S. NO. 7782-44-7 OXYGEN-2 WARNING: HIGH PRESSURE OXIDIZING GAS. VIGOROUSLY ACCELERATES COMBUSTION. NO SMOKING IN THE PRESENCE OF OXYGEN OR A FIRE MAY RESULT. KEEP OIL AND GREASE AWAY. OPEN VALVE SLOWLY. STORE AND USE WITH ADEQUATE VENTILATION. USE ONLY WITH EQUIPMENT CLEANED FOR OXYGEN SERVICE AND RATED FOR CYLINDER PRESSURE. USE A BACK FLOW PREVENTATIVE DEVICE IN THE PIPING. CLOSE VALVE AFTER EACH USE AND WHEN EMPTY. CYLINDER TEMPERATURE SHOULD NOT EXCEED 52 C (125 F) USE IN ACCORDANCE WITH THE MATERIAL SAFETY DATA SHEET (MSDS) DO NOT REMOVE THIS PRODUCT LABEL.

WARNING: FOR EMERGENCY USE ONLY WHEN ADMINISTERED BY PROPERLY TRAINED PERSONNEL FOR OXYGEN DEFICIENCY AND RESUSCITATION. FOR ALL OTHER MEDICAL APPLICATIONS Rx ONLY. UNINTERRUPTED USE OF HIGH CONCENTRATIONS OF OXYGEN OVER A LONG DURATION WITHOUT MONITORING ITS EFFECT ON OXYGEN CONTENT OF ARTERIAL BLOOD MAY BE HARMFUL. USE ONLY WITH PRESSURE REDUCING EQUIPMENT AND APPARATUS DESIGNED FOR OXYGEN. DO NOT ATTEMPT TO USE ON PATIENTS WHO HAVE STOPPED BREATHING UNLESS USED IN CONJUNCTION WITH RESUSCITATIVE EQUIPMENT. PRODUCED BY AIR LIQUEFACTION.

**OXYGEN,
COMPRESSED
USP**
UN 1072

OXYGEN
2

C.A.S. NO. 7782-44-7

Ameri-Quipt of N.C.
708 A Fleming Street
Hendersonville, NC 28791
828-692-5577
828-884-2808

PRODUCED BY AIR LIQUEFACTION

Δ WARNING: HIGH PRESSURE OXIDIZING GAS, VIGOROUSLY ACCELERATES COMBUSTION, NO SMOKING IN THE PRESENCE OF OXYGEN OR A FIRE MAY RESULT.

Δ WARNING: For emergency use only when administered by properly trained personnel for oxygen deficiency and resuscitation. For all other medical applications, Rx only. Uninterrupted use of high concentrations of oxygen over a long duration, without monitoring its effect on oxygen content of arterial blood, may be harmful. Use only with pressure reducing equipment and apparatus designed for oxygen. Do not attempt to use on patients who have stopped breathing, unless used in conjunction with resuscitative equipment.

Δ CAUTION: Keep oil and grease away. Store and use with adequate ventilation. Use only with equipment cleaned for oxygen service and rated for cylinder pressure. Open valve slowly. Close valve after each use and when empty.

Cylinder temperature should not exceed 52°C (125°F).

USE IN ACCORDANCE WITH MATERIAL SAFETY DATA SHEET.
DO NOT REMOVE THIS PRODUCT LABEL.

Size	Contents	Size	Contents
<input type="checkbox"/> "E"	682 Liters	<input type="checkbox"/> "M-6"	165 Liters
<input type="checkbox"/> "D"	415 Liters	<input type="checkbox"/> "M-60"	1725 Liters
<input type="checkbox"/> "C"	248 Liters	<input type="checkbox"/> Other	____ Liters

OXYGEN REFRIGERATED LIQUID LABEL

OXYGEN REFRIGERATED LIQUID USP UN1073 CONTENTS _____ LITERS ALWAYS KEEP CONTAINER IN UPRIGHT POSITION. WARNING: DO NOT CHANGE OR FORCE FIT CONNECTIONS. EXTREMELY COLD LIQUID AND GAS UNDER PRESSURE. VIGOROUSLY ACCELERATES COMBUSTION. NO SMOKING IN THE PRESENCE OF OXYGEN OR A FIRE MAY RESULT. COMBUSTIBLES IN CONTACT WITH LIQUID OXYGEN MAY EXPLODE ON IGNITION OR IMPACT. CAN CAUSE SEVERE FROSTBITE. KEEP OIL, GREASE AND COMBUSTIBLES AWAY. USE ONLY WITH EQUIPMENT CLEANED FOR OXYGEN SERVICE. STORE AND USE WITH ADEQUATE VENTILATION. DO NOT GET LIQUID IN EYES, ON SKIN OR CLOTHING. FOR LIQUID WITHDRAWAL, WEAR FACE SHIELD AND GLOVES. DO NOT DROP. USE SUITABLE HAND TRUCK FOR CONTAINER MOVEMENT. AVOID SPILLS. DO NOT WALK OR ROLL EQUIPMENT OVER SPILLS. CONTAINER TEMPERATURE SHOULD NOT EXCEED 52C (125 F) CLOSE VALVE AFTER EACH USE AND WHEN EMPTY. USE A BACK FLOW PREVENTATIVE DEVICE IN THE PIPING. USE IN ACCORDANCE WITH THE MATERIAL SAFETY DATA SHEET (MSDS)

FIRST AID: IN CASE OF FROSTBITE OBTAIN MEDICAL TREATMENT IMMEDIATELY.

WARNING: FOR EMERGENCY USE ONLY WHEN ADMINISTERED BY PROPERLY TRAINED PERSONNEL FOR OXYGEN DEFICIENCY AND RESUSCITATION. FOR ALL OTHER MEDICAL APPLICATIONS R_x ONLY. UNINTERRUPTED USE OF HIGH CONCENTRATIONS OF OXYGEN OVER A LONG DURATION WITHOUT MONITORING ITS EFFECT ON OXYGEN CONTENT OF ARTERIAL BLOOD MAY BE HARMFUL. USE ONLY WITH PRESSURE REDUCING EQUIPMENT AND APPARATUS DESIGNED FOR OXYGEN. DO NOT ATTEMPT TO USE ON PATIENTS WHO HAVE STOPPED BREATHING, UNLESS USED IN CONJUNCTION WITH RESUSCITATIVE EQUIPMENT. PRODUCED BY AIR LIQUEFACTION. DO NOT REMOVE THIS PRODUCT LABEL.

**OXYGEN,
REFRIGERATED
LIQUID
USP**
UN 1073



CONTENTS: _____ LITERS

↑ ALWAYS KEEP CONTAINER IN UPRIGHT POSITION. ↓

- ⚠ WARNING:** DO NOT CHANGE OR FORCE FIT CONNECTIONS.
EXTREMELY COLD OXIDIZING LIQUID AND GAS UNDER PRESSURE
VIGOROUSLY ACCELERATES COMBUSTION.
NO SMOKING IN THE PRESENCE OF OXYGEN OR A FIRE MAY RESULT
COMBUSTIBLES IN CONTACT WITH LIQUID OXYGEN MAY EXPLODE ON IGNITION OR IMPACT.
CAN CAUSE SEVERE FROSTBITE.
Keep oil, grease, and combustibles away.
Use only with equipment cleaned for oxygen service.
Store and use with adequate ventilation.
Do not get liquid in eyes, on skin or clothing.
For liquid withdrawal, wear face shield and gloves.
Do not drop. Use suitable hand truck for container movement.
Avoid spills. Do not walk on or roll equipment over spills.
Container temperature should not exceed 52°C (125°F).
Close valve after each use and when empty.
Use a back flow preventive device in the piping.
Use in accordance with the Material Safety Data Sheet (MSDS).
- ⚠ FIRST AID:** IN CASE OF FROSTBITE, obtain medical treatment immediately.
- ⚠ WARNING:** For emergency use only when administered by properly trained personnel for oxygen deficiency and resuscitation. For all other medical applications, Rx Only.
Uninterrupted use of high concentrations of oxygen over a long duration, without monitoring its effect on oxygen content of arterial blood, may be harmful. Use only with pressure reducing equipment and apparatus designed for oxygen. Do not attempt to use on patients who have stopped breathing, unless used in conjunction with resuscitative equipment.
Produced by Air Liquefaction
DO NOT REMOVE THIS PRODUCT LABEL.

CAS NO: 7782-44-7

OXYGEN

oxygen gas

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:60555-622
Route of Administration	RESPIRATORY (INHALATION)		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYGEN (UNII: S88TT14065) (OXYGEN - UNII:S88TT14065)	OXYGEN	99 L in 100 L

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60555-622-01	40 L in 1 CYLINDER		
2	NDC:60555-622-02	113 L in 1 CYLINDER		
3	NDC:60555-622-03	170 L in 1 CYLINDER		
4	NDC:60555-622-04	164 L in 1 CYLINDER		
5	NDC:60555-622-05	240 L in 1 CYLINDER		
6	NDC:60555-622-06	430 L in 1 CYLINDER		
7	NDC:60555-622-07	675 L in 1 CYLINDER		
8	NDC:60555-622-08	1725 L in 1 CYLINDER		
9	NDC:60555-622-09	46 L in 1 DEWAR		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved medical gas		01/01/1982	

Labeler - Ameri-Quipt of North Carolina, Inc (062549233)

Registrant - Ameri-Quipt of North Carolina, Inc (062549233)

Establishment

Name	Address	ID/FEI	Business Operations
Ameri-Quipt of North Carolina, Inc		062549233	manufacture

Revised: 1/2010

Ameri-Quipt of North Carolina, Inc