

OXYGEN- oxygen gas
Dade Medical, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

Oxygen, USP

Principal for Oxygen Product

**OXYGEN,
COMPRESSED
USP**

UN 1072



C.A.S. NO: 7782-44-7

Manufactured / Filled by:
Dade Medical, Inc.
3700 Commerce Parkway
Miramar, FL 33025
305-485-0804

PRODUCED BY AIR LIQUEFACTION

⚠ WARNING: CONTENTS UNDER HIGH PRESSURE AND CAN VIGOROUSLY ACCELERATE COMBUSTION. NO SMOKING IN THE PRESENCE OF OXYGEN OR A FIRE MAY RESULT. DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME AND USE ONLY WITH EQUIPMENT CONDITIONED FOR OXYGEN SERVICE.

⚠ WARNING: For emergency use only when administered by properly trained personnel for oxygen deficiency and resuscitation. For all other medical applications, **Rx** only. **Caution: Federal Law prohibits dispensing without prescription.**

WARNING: Uninterrupted use of high concentrations of oxygen for more than five (5) hours may be harmful. Do NOT attempt to use on patients who have stopped breathing, unless used in conjunction with resuscitative equipment.

KEEP OUT OF REACH OF CHILDREN.

⚠ Caution: Keep free from oil and grease. Open valve slowly and close valve after each use and when empty. Federal Law Requires that this Container be Filled with Oxygen U.S.P. Only by Establishments Registered as a Drug Producer in accordance with the Federal Food, Drug and Cosmetic Act.

DO NOT REMOVE THIS PRODUCT LABEL

Size	Contents	Size	Contents
<input type="checkbox"/> "E"	682 Liters	<input type="checkbox"/> "M-6"	165 Liters
<input type="checkbox"/> "D"	415 Liters	<input type="checkbox"/> "H"	6900 Liters
<input type="checkbox"/> "C"	248 Liters	<input type="checkbox"/> "Other"	_____ Liters

OXYGEN

COMPRESSED

USP

UN 1072

PRODUCED BY AIR LIQUIFICATION

WARNING: HIGH PRESSURE OXIDIZING GAS, VIGOROUSLY ACCELERATES COMBUSTION. NO SMOKING IN THE PRESENCE OF OXYGEN OR A FIRE MAY RESULT.


WARNING: For emergency use only when administered by properly trained personnel for oxygen deficiency and resuscitation. For all other medical applications, Rx only.

WARNING: Federal law prohibits dispensing without prescription. Uninterrupted use of high concentrations of oxygen over a long duration, without monitoring its effect on oxygen content of arterial blood, may be harmful. Use only with pressure reducing equipment and apparatus designed for oxygen. Do not attempt to use on patients who have stopped breathing, unless used in conjunction with resuscitative equipment.

WARNING: Keep oil and grease away. Use only with equipment cleaned for oxygen service and rated for cylinder pressure. Open valve slowly. Close valve after each use and when empty.

USE IN ACCORDANCE WITH MATERIAL SAFETY DATA SHEET.

DO NOT REMOVE THIS PRODUCT LABEL.



**USP
UN1073**



CONTENTS: _____ LITERS

↑ ALWAYS KEEP CONTAINER IN UPRIGHT POSITION ↑

■ ALWAYS KEEP CONTAINER IN UPRIGHT POSITION. ■

⚠ WARNING: EXTREMELY COLD OXIDIZING LIQUID AND GAS UNDER PRESSURE.

VIGOROUSLY ACCELERATES COMBUSTION.

COMBUSTIBLES IN CONTACT WITH LIQUID OXYGEN MAY EXPLODE ON IGNITION OR IMPACT.

CAN CAUSE SEVERE FROSTBITE.

Keep oil, grease, and combustibles away.

Use only with equipment cleaned for oxygen service.

Store and use with adequate ventilation.

Do not get liquid in eyes, on skin or clothing.

For liquid withdrawal, wear face shield and gloves.

Do not drop. Use hand truck for container movement.

Avoid spills. Do not walk on or roll equipment over spills.

Container temperature should not exceed 52°C (125° F).

Close valve after each use and when empty.

Use in accordance with the Material Safety Data Sheet (MSDS).

⚠ FIRST AID: IN CASE OF FROSTBITE, obtain medical treatment immediately.

⚠ WARNING: For emergency use only when administered by properly trained personnel for oxygen deficiency and resuscitation. For all other medical applications, **R_x only**.

Uninterrupted use of high concentrations of oxygen over a long duration, without monitoring its effect on oxygen content of arterial blood, may be harmful. Use only with pressure reducing equipment and apparatus designed for oxygen. Do not attempt to use on patients who have stopped breathing, unless used in conjunction with resuscitative equipment.

Keep Out of Reach of Children

Produced by Air Liquefaction

DO NOT REMOVE THIS PRODUCT LABEL

Manufactured / Filled By:

Dade Medical, Inc.

3700 Commerce Parkway

Miramar, FL 33025

305-485-0804

CAS NO: 7782-44-7

OXYGEN,
REFRIGERATED

LIQUID USP

UN 1073

CONTENTS _____ LITERS

CAS NO: 7782-44-7

DANGER:

REFRIGERATED LIQUEFIED OXIDIZING GAS. ALWAYS
KEEP CONTAINER IN UPRIGHT POSITION. DO NOT
CHANGE OR FORCE FIT CONNECTIONS. MAY CAUSE OR INTENSIFY
FIRE; OXIDIZER, CONTAINS REFRIGERATED GAS: MAY CAUSE
CRYOGENIC BURNS OR INJURY. COMBUSTIBLES IN CONTACT WITH
LIQUID OXYGEN MAY EXPLODE ON IGNITION OR IMPACT.

Do not get in eyes, on skin, or on clothing. Keep reduction valves free
from grease and oil. In case of fire, stop leak if safe to do so. Avoid
spills. Do not walk or roll equipment over spills. Store and use in a well
ventilated place. Use only with equipment of compatible materials
of construction and rated for cylinder pressure. Wear cold insulating
gloves/face shield/eye protection. Protect from sunlight when ambient
temperature exceeds 52 F. Container temperature should
not exceed 52 F. Close valve after each use and when
empty. Use a back flow preventative device in the piping. Do not
handle until all safety precautions have been read and understood.
Use only with equipment cleaned for oxygen service. Keep/store away
from combustible materials. Do not drop. Use suitable hand truck for
container movement.

FIRST AID:

IN CASE OF FROSTBITE: Thaw frosted parts with lukewarm water.

Do not rub affected area. Get immediate medical advice/attention.

DO NOT REMOVE THIS PRODUCT LABEL.

For emergency use only when administered by properly trained
personnel for oxygen deficiency and resuscitation. For all other medical
applications, Rx only. Uninterrupted use of high concentrations of
oxygen over a long duration, without monitoring its effect on oxygen
content of arterial blood, may be harmful. Use only with pressure
reducing equipment and apparatus designed for oxygen. Do not
attempt to use on patients who have stopped breathing unless used in
conjunction with resuscitative equipment. Produced by Air Liquefaction.
ALWAYS KEEP CONTAINER IN UPRIGHT POSITION

OXYGEN

oxygen gas

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70672-123
Route of Administration	RESPIRATORY (INHALATION)		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYGEN (UNII: S88TT14065) (OXYGEN - UNII:S88TT14065)	OXYGEN	99 L in 100 L

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70672-123-01	165 L in 1 CYLINDER; Type 0: Not a Combination Product	03/01/2016	
2	NDC:70672-123-02	248 L in 1 CYLINDER; Type 0: Not a Combination Product	03/01/2016	
3	NDC:70672-123-03	415 L in 1 CYLINDER; Type 0: Not a Combination Product	03/01/2016	
4	NDC:70672-123-04	682 L in 1 CYLINDER; Type 0: Not a Combination Product	03/01/2016	
5	NDC:70672-123-06	43 L in 1 DEWAR; Type 0: Not a Combination Product	03/01/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved medical gas		03/01/2016	

Labeler - Dade Medical, Inc. (808840953)

Registrant - Dade Medical, Inc. (808840953)

Establishment

Name	Address	ID/FEI	Business Operations
Dade Medical, Inc.		808840953	manufacture(70672-123)

Revised: 12/2017

Dade Medical, Inc.