

ASPIRIN- aspirin tablet, chewable
Meijer Distribution Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Meijer Distribution, Inc. Aspirin 81mg Drug Facts

Active ingredient (in each chewable tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

- children under 12 years: consult a doctor

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium

Questions or comments?

1-800-719-9260

Principal Display Panel

VALUE 3 PACK 108 TOTAL TABLETS

meijer®

LOW DOSE

Compare to Bayer® Chewable Aspirin active ingredient

aspirin Pain Reliever (NSAID)

SEE NEW WARNING

81 mg | Aspirin Regimen | Talk to your doctor or other healthcare provider before using this product for your heart.

Chewable Orange Flavor

36 Tablets

actual size

3 - 36 Count Bottles Inside
= 108 Total Tablets



meijer
LOW DOSE
aspirin Pain Reliever (NSAID)
NDC 41250-467-08
81 mg | Aspirin Regimen | Talk to your doctor or other healthcare provider before using this product for your heart.
SEE NEW WARNING
Chewable Orange Flavor
36 Tablets
actual size

VALUE 3 PACK NDC 41250-467-08
108 TOTAL TABLETS

meijer **LOW DOSE** **aspirin** Pain Reliever (NSAID)
Compare to Bayer® Chewable Aspirin active ingredient**

SEE NEW WARNING 81 mg | Aspirin Regimen | Talk to your doctor or other healthcare provider before using this product for your heart. 

 Chewable Orange Flavor **36 Tablets**  actual size

meijer **LOW DOSE** **aspirin** Pain Reliever (NSAID)
Compare to Bayer® Chewable Aspirin active ingredient**

SEE NEW WARNING 81 mg | Aspirin Regimen | Talk to your doctor or other healthcare provider before using this product for your heart. 


 Chewable Orange Flavor **36 Tablets**  actual size

meijer **LOW DOSE** **aspirin** Pain Reliever (NSAID)
Compare to Bayer® Chewable Aspirin active ingredient**

SEE NEW WARNING 81 mg | Aspirin Regimen | Talk to your doctor or other healthcare provider before using this product for your heart. 

 Chewable Orange Flavor **36 Tablets**  actual size

A2082 6E C6



Drug Facts

Active ingredient (in each chewable tablet) Purpose
 Aspirin 81 mg (NSAID*).....Pain reliever
 *nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 ■ are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
 - you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood ■ have bloody or black stools
 ■ have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present ■ new symptoms occur
 - ringing in the ears or a loss of hearing occurs
- These could be signs of a serious condition

Drug Facts (continued)

Other information

■ store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium


Questions or comments?

1-800-719-9260

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

**This product is not manufactured or distributed by Bayer Aktiengesellschaft, owner of the registered trademark Bayer®.

DIST. BY MEJER
 DISTRIBUTION, INC.
 GRAND RAPIDS, MI 49544
 www.mejer.com



Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

ASPIRIN

aspirin tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-467
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	L467
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-467-08	3 in 1 CARTON	08/27/1992	
1	NDC:41250-467-68	36 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	08/27/1992	

Labeler - Meijer Distribution Inc (006959555)

Revised: 2/2023

Meijer Distribution Inc