

## **MAPAP EXTRA STRENGTH- acetaminophen capsule**

### **Major Pharmaceuticals**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **0617 - Major**

#### ***Drug Facts***

##### **Active Ingredient** (in each banded capsule)

Acetaminophen 500 mg

#### ***Purpose***

##### ***Purpose***

Pain reliever/ Fever reducer

#### ***Uses***

Temporarily reduces fever and relieves minor aches and pains due to:

- headache
- muscular aches
- common cold
- toothache
- backache
- minor pain of arthritis
- premenstrual and menstrual cramps

#### ***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

#### **Do not use**

- with any other drugs containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

#### **Ask a doctor before use if you have**

liver disease

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:**

In the case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

**Do not take more than directed (see overdose warning).**

**adults and children 12 years and over:**

- take 2 capsules every 6 hours while symptoms last
- Do not take more than 6 capsules in 24 hours, unless directed by a doctor
- Do not take for more than 10 days unless directed by a doctor

**children under 12 years:** ask a doctor

***Other information***

- store at room temperature

***Inactive Ingredients***

FD&C Blue #1, FD&C Red #40, Gelatin, Polyvinylpyrrolidone, Pregelatinized Starch, Sodium Starch Glycolate, Stearic Acid and Titanium Dioxide

**Questions?**

**Call: (800) 616-2471**

Distributed by: **Major® Pharmaceuticals**

17177 N Laurel Park Drive, Suite 233, Livonia, MI 48152 USA

Re-order No. 700691 Rev.05/19 M-29 4050-05-19

**Major**

**NDC 0904-1987-60**

**EXTRA STRENGTH**

**Mapap**

Acetaminophen 500 mg Each

CAPSULES SAFETY SEALED FOR YOUR PROTECTION

SEE NEW WARNINGS INFORMATION AND DIRECTIONS

ASPIRIN FREE

Strong Pain & Fever Relief

100 CAPSULES

MAJOR<sup>®</sup>

EXTRA STRENGTH

Māpap<sup>®</sup>

Acetaminophen 500 mg Each

SEE NEW WARNINGS INFORMATION AND DIRECTIONS

ASPIRIN FREE  
STRONG PAIN & FEVER RELIEF

100 CAPSULES

NDC 0904-1987-60

TAMPER EVIDENT: DO NOT USE IF CAPSULE BAND OR IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each banded capsule) Purpose

Acetaminophen 500 mg ..... Pain reliever/ Fever reducer

Uses

Temporarily reduces fever and relieves minor aches and pains due to: • headache • muscular aches • common cold • toothache • backache • minor pain of arthritis • premenstrual and menstrual cramps

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take • more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Allergy alert:

acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are allergic to acetaminophen or any of the inactive ingredients in this product

Distributed By: MAJOR<sup>®</sup> PHARMACEUTICALS

17177 N Laurel Park Drive, Suite 233, Livonia, MI 48152 USA

Re-order No. 700691 Rev.05/19 MF-29 4050-06-19

3

0904198760

9

Let # & Exp. Date:

FEEL BETTER

Drug Facts (continued)

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In the case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

• do not take more than directed (see overdose warning)

adults and children 12 years and over: • take 2 capsules every 6 hours while symptoms last

• do not take more than 6 capsules in 24 hours, unless directed by a doctor

• do not take for more than 10 days unless directed by a doctor

children under 12 years: ask a doctor

Other information store at room temperature

Inactive ingredients FD&C Blue #1, FD&C Red #40, Gelatin, Polyvinylpyrrolidone, Pregelatinized Starch, Sodium Starch Glycolate, Stearic Acid and Titanium Dioxide

Questions or comments? Call (800) 616-2471

MAPAP EXTRA STRENGTH

acetaminophen capsule

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-1987
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	500 mg

Inactive Ingredients				
Ingredient Name				Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
GELATIN (UNII: 2G86QN327L)				
POVIDONE K30 (UNII: U725QWY32X)				
STARCH, CORN (UNII: O8232NY3SJ)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	white, red	Score	no score	
Shape	CAPSULE	Size	22mm	
Flavor		Imprint Code	CPC;617	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-1987-60	100 in 1 BOTTLE; Type 0: Not a Combination Product	12/22/1997	
2	NDC:0904-1987-80	1000 in 1 BOTTLE; Type 0: Not a Combination Product	01/29/2001	04/30/2017
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final		part343	12/22/1997	

**Labeler** - Major Pharmaceuticals (191427277)