CAREONE ASPIRIN- as pirin tablet, chewable American Sales Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

American Sales Company Aspirin Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

• store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, D&C red #27 aluminum lake, dextrose excipient, FD&C red #40 aluminum lake, flavors, saccharin sodium

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Active Ingredient in Bayer® Chewable Aspirin

Adult Low Strength

ASPIRIN 81 mg

Actual Size

Pain Reliever (NSAID)

Chewable Cherry Flavor

OUR PHARMACISTS RECOMMEND

Low Dose Aspirin Regimen

Talk to your doctor or other healthcare provider before using this product for your heart.

36 TABLETS



NDC 41520-274-68

Compare to Active Ingredient in Bayer® Chewable Aspirin†

81 mg

Pain Reliever (NSAID) Chewable Cherry Flavor



Low Dose Aspirin Regimen Talk to your doctor or other healthcare provider before using this product for your heart.

36 TABLETS

51468 OF C2

CODE VILEV



DO NOT USE IF PRINTED SEAL UNDER

Drug Facts

Active ingredient Purpose *(in each tablet)* Aspirin 81 mg (NSAID)*. .Pain relieve

*nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious ill ness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives facial swelling asthma (wheezing) shock
- Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

Drug Facts (continued)

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ for at least 7 days after ton sillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- ■you experience any of the following signs of stomach bleeding:
- feel faint vomit blood
- have bloody or black stools
- have stomach pain that does not get better ▶

Drug Facts (continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days ■ redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 1 2 years and over: take 4 to 8 table ts every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

store at 20-25°C (68-77°F)

o la

Questions or comments? 1-800-719-9260 IBUTEDBY FOODHOLD U.S.A., LLC OV ER, MD 20785: 1-877-846-9949 Ahold Licensing Sarl not manufactured or distributed b distributor of Bayer® Chewable A

Drug Facts (contin ued)
Inactive Ingredients corn starch
D&C red #27 alumin um lake, dextross
excipient, FD &C red #40 alumin um la saccharin sodium

orn starch, e, dextrose ıminum lake,

CAREONE ASPIRIN

aspirin tablet, chewable

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-274
--------------	----------------	--------------------	---------------

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color	PINK	Score	no score
Shape	ROUND	Size	8 mm
Flavor	CHERRY	Imprint Code	L274
Contains			

Packaging

П	i uchuşing			
l	# Item Code	Package Description	Marketing Start Date	Marketing End Date
l	1 NDC:41520-274-68	1 in 1 CARTON	10/09/2017	
	1	36 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	10/09/2017	

Labeler - American Sales Company (809183973)

Revised: 8/2018 American Sales Company