

RANITIDINE- ranitidine tablet, coated
Sam's West Inc

Drug Facts

Active ingredient (in each tablet)

Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg)

Purpose

Acid reducer

Use(s)

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

Other information

- do not use if printed foil under bottle cap is open or torn
- store at 20°-25°C (68°-77°F)
- avoid excessive heat or humidity
- protect from light
- this product is sodium and sugar free

Inactive ingredients

FD&C red #40 aluminum lake, hypromellose, iron oxide black, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide.

Questions?

Call **1-888-375-3784**

Revised: 0419

Consumer Information**What you should know about****MAXIMUM STRENGTH****Ranitidine Tablets USP, 150 mg**

(Please read all of this information before taking MAXIMUM STRENGTH Ranitidine Tablets USP, 150 mg. Save this leaflet for future reference.)

What are MAXIMUM STRENGTH Ranitidine Tablets USP, 150 mg?

- MAXIMUM STRENGTH Ranitidine Tablets USP, 150 mg contains 150 mg of ranitidine


(as ranitidine hydrochloride USP, 168 mg), a medicine that doctors have prescribed more than 200 million times worldwide.

Excellent Safety Record

- The ingredient in MAXIMUM STRENGTH Ranitidine Tablets USP, 150 mg, ranitidine, has been prescribed by doctors for years to treat millions of patients safely and effectively. The active ingredient in MAXIMUM STRENGTH Ranitidine Tablets USP, 150 mg has been taken safely with many frequently prescribed medications.
- MAXIMUM STRENGTH Ranitidine Tablets USP, 150 mg are sodium and sugar free.

What symptoms does MAXIMUM STRENGTH Ranitidine Tablets USP, 150 mg relieve and prevent?

- MAXIMUM STRENGTH Ranitidine Tablets USP, 150 mg relieve and prevent heartburn associated with acid indigestion and sour stomach.
- Certain foods or beverages, and even lying down to sleep, can cause heartburn associated with acid indigestion and sour stomach. It is normal to the stomach to produce acid, especially after consuming food or beverages. However, acid in the wrong place, such as the esophagus, or too much acid, can cause burning pain and discomfort.



How should I take MAXIMUM STRENGTH Ranitidine Tablets USP, 150 mg?

- To **relieve** symptoms, swallow 1 tablet with a glass of water.
- To **prevent** symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn.
- This medicine can be used up to twice daily (up to 2 tablets in 24 hours).
- MAXIMUM STRENGTH Ranitidine Tablets USP, 150 mg should not be given to children under 12 years old unless directed by a doctor.
- **Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers.

How does MAXIMUM STRENGTH Ranitidine Tablets USP, 150 mg work?

- MAXIMUM STRENGTH Ranitidine Tablets USP, 150 mg reduce the production of stomach acid. This is what makes MAXIMUM STRENGTH Ranitidine Tablets USP, 150 mg different from antacids, which neutralize the acid already in your stomach.

Antacids do not reduce the production of acid.

Tips for managing heartburn

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Do not Use

- If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

When should I see a doctor?

- **Ask a doctor before use if you have**
 - had heartburn over 3 months. This may be a sign of a more serious condition.
 - heartburn with **lightheadedness, sweating or dizziness**
 - chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 - frequent **chest pain**
 - frequent wheezing, particularly with heartburn
 - unexplained weight loss
 - nausea or vomiting
 - stomach pain
 - kidney disease
- **Ask a doctor or pharmacist before use if you are**
 - taking a prescription drug. Acid reducers may interact with certain prescription drugs.
- **Stop use and ask a doctor if**
 - your heartburn continues or worsens
 - you need to take this product for more than 14 days
- **If pregnant or breast-feeding**, ask a health professional before use.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions? Call 1-888-375-3784

BOTTLES: Bottle is sealed with printed foil under cap. Do not use if printed foil is open or torn.

BLISTERS: Do not use if the individual blister unit is open or torn.

Revised: 0419

SEALED WITH PRINTED FOIL UNDER CAP FOR YOUR PROTECTION. IMPORTANT:

Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

PACKAGE LABEL PRINCIPAL DISPLAY PANEL

Container Label:

Carton Label

Maximum Strength

Ranitidine Tablets USP, 150 mg
ACID REDUCER

PREVENTS & RELIEVES

heartburn associated with acid indigestion
and sour stomach

190 Tablets(190 doses)

Drug Facts
Active ingredient (in each tablet) Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg) **Purpose** Acid reducer

Uses
■ relieves heartburn associated with acid indigestion and sour stomach
■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers
Do not use
■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
■ with other acid reducers
Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck, or shoulder; or lightheadedness
■ frequent chest pain
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain
■ kidney disease
Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)
Directions
■ adults and children 12 years and over:
■ to relieve symptoms, swallow 1 tablet with a glass of water
■ to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
■ can be used up to twice daily (do not take more than 2 tablets in 24 hours)
■ children under 12 years: ask a doctor
Other information
■ do not use if printed foil under bottle cap is open or torn
■ store at 20°-25°C (68°-77°F)
■ avoid excessive heat or humidity
■ protect from light
■ this product is sodium and sugar free
Inactive ingredients
FD&C red #40 aluminum lake, hypromellose, iron oxide black, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide.
Questions? Call 1-888-375-3784
TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING
IMPORTANT: Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.
*This product is not manufactured or distributed by Chatter, Inc., distributor of Zantac 150® Tablets. Zantac 150® is a registered trademark of Chatter, Inc.

100% MONEY BACK GUARANTEE
SUPERIOR QUALITY AND PERFORMANCE
We would like to hear from you with any comments or suggestions. In the continental U.S. or Canada, you can call us at 1-888-375-3784 from 9:00 a.m. to 7:00 p.m. EST Monday - Friday.

Compare to Zantac 150[®] active ingredient*

Member's Mark. QUALITY GUARANTEED

maximum strength ranitidine tablets USP 150 mg acid reducer

PREVENTS & RELIEVES HEARTBURN associated with acid indigestion and sour stomach

190 Tablets (190 Doses)
2X95 count bottles

LOT
EXP

DISTRIBUTED BY:
SAM'S WEST, INC.,
BENTONVILLE, AR 72716

Revised: 07/19

Leave this area blank (no color or no markings) for "Made in" In-line Printers ONLY

Bottle

Maximum Strength

Ranitidine Tablets USP, 150 mg

ACID REDUCER

PREVENTS & RELIEVES

heartburn associated with acid indigestion
and sour stomach

95 Tablets(95 doses)

Compare to Zantac 150[®]
active ingredient*



maximum strength
ranitidine
tablets USP
150 mg
acid reducer

actual
size

95
Tablets
(95 Doses)

• PREVENTS & RELIEVES
HEARTBURN associated
with acid indigestion
and sour stomach

TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING
IMPORTANT: This label does not contain full product information. See carton for complete information. Read the directions, consumer information leaflet and warnings before use. Retain carton and leaflet for reference.

Active ingredient (in each tablet) Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg) **Purpose** Acid reducer

Uses ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers
Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers
Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent chest pain ■ frequent wheezing, particularly with heartburn
■ unexplained weight loss ■ nausea or vomiting
■ stomach pain ■ kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if ■ your heartburn worsens ■ you need to take this product for more than 14 days ■ **pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ adults and children 12 years and over: ■ **to relieve symptoms**, swallow 1 tablet with a glass of water ■ **to prevent symptoms**, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn ■ can be used up to twice daily (do not take more than 2 tablets in 24 hours) ■ children under 12 years: ask a doctor

Questions? Call **1-888-375-3784**

Revised: 07/19

 LOT
EXP

DISTRIBUTED BY:
SAM'S WEST, INC.
BENTONVILLE, AR 72716

*This product is not manufactured or distributed by Chatterm, Inc., distributor of Zantac 150[®] Tablets. Zantac 150[®] is a registered trademark of Chatterm, Inc.

150077012

RANITIDINE

ranitidine tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68196-985(NDC:55111-404)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Ranitidine Hydrochloride (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	150 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
magnesium stearate (UNII: 70097M6I30)	
cellulose, microcrystalline (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
titanium dioxide (UNII: 15FIX9V2JP)	

Product Characteristics

Color	PINK	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	R150
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68196-985-19	2 in 1 CARTON	01/05/2011	09/30/2019
1		95 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078192	01/05/2011	

Labeler - Sam's West Inc (051957769)

Revised: 12/2019

Sam's West Inc