

NEONATAL VITAMIN- neonatal vitamin tablet
NEOMED PHARMACEUTICAL

NEONATAL VITAMIN TABLETS

Prenatal Vitamin With Other Vitamins & Minerals For Nutritional Supplementation

SUPPLEMENT FACTS:

Serving Size: One Tablet

Each Tablet contains:

Vitamin A (as acetate, beta carotene)	1200 mcg
Vitamin D3 (as cholecalciferol)	10 mcg
Vitamin E (as dl-alpha tocopheryl acetate)	4.6 mg
Vitamin C (Ascorbic acid)	100 mg
Vitamin B1 (as thiamine HCl)	1.5 mg
Vitamin B2 (as riboflavin)	1.7 mg
Vitamin B3 (as niacinamide)	1.8 mg
Vitamin B6 (as pyridoxine HCl)	2.6 mg
Folic acid	800 mcg
Vitamin B12 (as cyanocobalamin)	4 mcg
Iron (as ferrous fumarate)	27 mg
Citrus bioflavonoids complex 50% bio flavonoids	2 mg
Calcium (as carbonate sulfate)	263 mg
Zinc (as zinc oxide)	25 mg
Boron (as boron citrate)	0.2 mg
Inositol	5 mg

Other Ingredients: Dibasic calcium phosphate, microcrystalline croscarmellose sodium, stearic acid, silicon dioxide, titanium dioxide (as color), polyethylene glycol, sucrose, magnesium stearate, talc, triacetin, HPMC E15, HPMC E5/E6, FD&C RED #40 (LAKES), FD&C YELLOW # 6 (LAKE)

WARNINGS

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

DOSAGE & ADMINISTRATION

As with any supplement, if you are pregnant, nursing, or taking medication, consult your doctor before use.

The safe upper limit for daily intake of folic acid is 1000 mcg (1 mg).

Do not take Prenatal with other multivitamin formulas unless recommended by your doctor.

IMPORTANT INFORMATION

Long-term intake of high levels of vitamin A (excluding that sourced from beta-carotene) may increase the risk of osteoporosis in adults. Do not take this product if taking other vitamin A supplements.

Information for Patients

Patients should be counseled to disclose all medical conditions, including use of all medications,

vitamins and supplements, pregnancy, and breastfeeding.

Pediatric Use

Not recommended for pediatric use.

HOW SUPPLIED

NEONATAL VITAMIN ; available in bottles of 100 Tablets. 70898-105-01

SAFE HANDLING WARNING

Store at room temperature.

Avoid excessive heat above 40°C (104°F.)

QUESTIONS? COMMENTS?

Manufactured for:

NEOMED Pharmaceutical,

Hauppauge, NY-11788.

REV.:08/20

Distributed by:

AUM Pharmaceuticals,

320 Oser Ave.

Hauppauge, NY-11788.

DISCLAIMER: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. [For further information about unapproved drugs, click here.](#)



Each prenatal tablet contains the following ingredients for nutritional supplement:

Vitamin A (As Acetate and Beta Carotene)	1200 MCG
Vitamin C (Ascorbic Acid)	100 MG
Vitamin D3 (As Cholecalciferol)	10 MCG
Vitamin E (As Di-Alpha Tocopheryl Acetate)	4.6 MG
Vitamin B1 (As Thiamine Mononitrate)	1.5 MG
Vitamin B2 (Riboflavin)	1.7 MG
Vitamin B3 (As Nicotinamide)	18 MG
Vitamin B6 (As Pyridoxine Hcl)	2.6 MG
Folic Acid	800 MCG
Vitamin B12 (As Cyanocobalamin) On DCP	4 MCG
Calcium (As Calcium Carbonate and Sulfate)	263 MG
Iron (As Ferrous Fumarate)	27 MG
Zinc (As Zinc Oxide)	25 MG
Inositol	5 MG
Citrus Bioflavonoid Complex 50%	2 MG
Boron (As Boron Citrate)	0.2 MG

OTHER INGREDIENTS

MICROCRYSTALLINE CELLULOSE, DICALCIUM PHOSPHATE, STEARIC ACID, CROSCARMELLOSE SODIUM, SILICON DIOXIDE, MAGNESIUM STEARATE, HPMC K15, HPVC K5/76, PEG - 8000, TALC, FD&C RED # 40 (LAKES), FD&C YELLOW # 6 (LAKES), TRIACETIN.

Description: Neonatal Tablet is an orally administered prenatal dietary supplement and should be administered under the supervision of a licensed medical practitioner.

See package insert for full prescribing information.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

CAUTION: High levels of folic acid may, especially in older adults, hide signs of Vitamin B-12 deficiency (such as pernicious anemia), a condition that can cause nerve damage.

Store at 15-30°C (59-86°F). [See USP Controlled Room Temperature.] Protect from light and moisture. Dispense in a tight, light-resistant container. Notice: Contact with moisture may produce surface discoloration or erosion.

Call your licensed medical practitioner about side effect. You may report side effects by calling 1-866-760-6565.

Manufactured for:
NEOMED Pharmaceutical,
Hauppauge, NY-11788.

Distributed by:
AUM Pharmaceuticals,
320 Oser Ave.
Hauppauge, NY-11788.

REV:08/20

NEONATAL VITAMIN

neonatal vitamin tablet

Product Information

Product Type	DIETARY SUPPLEMENT	Item Code (Source)	NHRC:70898-105
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
VITAMIN A (UNII: 8 1G40H8B0T) (VITAMIN A - UNII:8 1G40H8B0T)	VITAMIN A	1200 ug
ASCORBIC ACID (UNII: PQ6CK8PD0R) (ASCORBIC ACID - UNII:PQ6CK8PD0R)	ASCORBIC ACID	100 mg
CHOLECALCIFEROL (UNII: 1C6V77QF41) (CHOLECALCIFEROL - UNII:1C6V77QF41)	CHOLECALCIFEROL	10 ug
.ALPHA.-TOCOPHEROL ACETATE (UNII: 9 E8X80D2L0) (.ALPHA.-TOCOPHEROL - UNII:H4N855PNZ1)	.ALPHA.-TOCOPHEROL ACETATE	4.6 mg
THIAMINE (UNII: X66NSO3N35) (THIAMINE ION - UNII:4ABT0J945J)	THIAMINE	1.5 mg
RIBOFLAVIN (UNII: TLM2976OFR) (RIBOFLAVIN - UNII:TLM2976OFR)	RIBOFLAVIN	1.7 mg
NIACINAMIDE (UNII: 25X51I8RD4) (NIACINAMIDE - UNII:25X51I8RD4)	NIACINAMIDE	1.8 mg
PYRIDOXINE HYDROCHLORIDE (UNII: 68Y4CF58BV) (PYRIDOXINE - UNII:KV2JZ1B16Z)	PYRIDOXINE	2.6 mg
FOLIC ACID (UNII: 935E97BOY8) (FOLIC ACID - UNII:935E97BOY8)	FOLIC ACID	800 ug
CYANOCOBALAMIN (UNII: P6YC3EG204) (CYANOCOBALAMIN - UNII:P6YC3EG204)	CYANOCOBALAMIN	4 ug
FERROUS FUMARATE (UNII: R5L488RY0Q) (FERROUS CATION - UNII:GW89581OWR)	FERROUS CATION	27 mg
INOSITOL (UNII: 4L6452S749) (INOSITOL - UNII:4L6452S749)	INOSITOL	5 mg
CITRUS BIOFLAVONOIDS (UNII: BD70459I50) (HESPERIDIN - UNII:E750O06Y6O)	CITRUS BIOFLAVONOIDS	2 mg
CALCIUM (UNII: SY7Q814VUP) (CALCIUM - UNII:SY7Q814VUP)	CALCIUM	263 mg
BORON (UNII: N9E3X5056Q) (BORON - UNII:N9E3X5056Q)	BORON	0.2 mg
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC CATION - UNII:13S1S8SF37)	ZINC CATION	25 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CALCIUM PHOSPHATE, DIBASIC, ANHYDROUS (UNII: L11K75P92J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TALC (UNII: 7SEV7J4R1U)	
POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
TRIACETIN (UNII: XHX3C3X673)	
SUCROSE (UNII: C151H8M554)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NHRIC:70898-105-01	1 in 1 PACKAGE		
1		100 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
DIETARY SUPPLEMENT		07/31/2018	

Supplement Facts		
Serving Size :		Serving per Container :
	Amount Per Serving	% Daily Value
color		
scoring	1	
shape		
size (solid drugs)	5 mm	

Labeler - NEOMED PHARMACEUTICAL (048388130)

Revised: 9/2020

NEOMED PHARMACEUTICAL