

**SILACE - docusate sodium liquid**  
**Silarx Pharmaceuticals, Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Silace Liquid**

Active Ingredient: Docusate sodium 10 mg (in each 1 mL)

Purpose: Stool Softener

**Uses**

- for gentle, reliable relief from occasional constipation (irregularity)
- generally produces bowel movement in 12 to 72 hours

**Warning**

- **Do not use** laxative products for longer than 1 week unless told to do so by a doctor
- if you are presently taking mineral oil unless told to do by a doctor

**Ask doctor before use if you have**

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that last over two weeks

**Stop use and ask a doctor** if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition

**If pregnant or breast feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- take as indicated by doctor
- this product must be given in a 6 to 8 oz. glass of milk or fruit juice or infant's formula to prevent throat irritation. Dose may be taken as a single daily dose or in divided doses
- dosage should be adjusted to individual response
- higher doses are recommended for initial therapy
- the effect on stools is usually apparent 1 to 3 days after the first dose

Adults and children over 12 years	50 to 200 mg (1 to 4 teaspoonful)
Children 6 to under 12 years	50 to 100 mg (1 to 2 teaspoonful)
Children 3 to under 6 years	25 to 50 mg (1/2 to 1 teaspoonful)
Children under 3 years of age	Ask a doctor

**Other information**

- store at room temperature 20°-25°C (68°-77°F)

- protect from freezing and excessive heat
- do not use if tamperevident safety seal around cap is broken or missing
- dispense in tight, light-resistant container with a child-resistant closure

**Inactive ingredients**

citric acid, D&C red no. 33, flavor, methylparaben, Poloxamer, propylene glycol, propylparaben, purified water, sodium citrate.

**Questions**

888-974-5279

This product is not manufactured or distributed by Purdue Products LP, distributor of Colace® Liquid.

Manufactured by:

Silarx Pharmaceuticals, Inc.  
1033 Stoneleigh Ave  
Carmel, NY 10512  
USA

## Drug Facts

**Active ingredient (in each 1 mL)** Purpose  
Docusate sodium 10 mg Stool softener

**Uses** ■ for gentle, reliable relief from occasional constipation (irregularity) ■ generally produces bowel movement in 12 to 72 hours

### Warnings

**Do not use** ■ laxative products for longer than 1 week unless told to do so by a doctor ■ if you are presently taking mineral oil unless told to do so by a doctor

**Ask a doctor before use if you have** ■ stomach pain ■ nausea ■ vomiting ■ noticed a sudden change in bowel habits that last over two weeks

**Stop use and ask a doctor if you have** rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.

**If pregnant or breast feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ take as directed by a doctor ■ this product must be given in a 6 to 8 oz. glass of milk or fruit juice to prevent throat irritation. Dose may be taken as a single daily dose or in divided doses ■ dosage should be adjusted to individual response ■ higher doses are recommended for initial therapy ■ the effect on stools is usually apparent 1 to 3 days after the first dose

adults and children 12 years of age and older	50 to 200 mg (1 to 4 teaspoonful)
children 6 to under 12 years of age	50 to 100 mg (1 to 2 teaspoonful)
children 3 to under 6 years of age	25 to 50 mg (1/2 to 1 teaspoonful)
children under 3 years of age	ask a doctor

**Other information** ■ store at room temperature 20°-25°C (68°-77°F) ■ protect from freezing and excessive heat

**Inactive ingredients** citric acid, D&C red no. 33, lemon-vanilla flavor, methylparaben, poloxamer, propylene glycol, propylparaben, purified water, sodium citrate

**Questions?** ☎ 888-974-5279



N  
3

54838-116-80

9

† This product is not manufactured or distributed by Purdue Products LP, distributor of Colace® Liquid.  
Rev. 08/13

Control # Exp. Date

# SILARX

NDC 54838-116-80

# SILACE LIQUID

**Docusate Sodium  
(1% Solution)**

**Stool Softener  
Stimulant Free**

**BULK CONTAINER –  
NOT FOR HOUSEHOLD USE**  
Pharmacist - Dispense in a tight, light-resistant container with child-resistant closure.

**TAMPER EVIDENT:**  
Do not use if tamper-evident safety seal around cap is broken or missing

† Compare to the active ingredient of Colace® Liquid

**1 Pint (473 mL)**

Manufactured by:  
Silarx Pharmaceuticals, Inc.  
1033 Stoneleigh Ave., Carmel, NY 10512

## SILACE

docusate sodium liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG LABEL	<b>Item Code (Source)</b>	NDC:54838-116
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>Docusate sodium</b> (Docusate)	Docusate sodium	10 mg in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
anhydrous citric acid	
D&C red no. 33	
methylparaben	
propylene glycol	
propylparaben	
sodium citrate	
water	
Poloxamer 407	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	LEMON (Lemon Vanilla Flavor)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54838-116-80	473 mL in 1 BOTTLE, PLASTIC		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	10/05/1990	

**Labeler** - Silarx Pharmaceuticals, Inc (161630033)