



## **CONTRAINDICATIONS**

Hypersensitivity to probenecid.

Children under 2 years of age.

Not recommended in persons with known blood dyscrasias or uric acid kidney stones.

Therapy with probenecid should not be started until an acute gouty attack has subsided.

## **WARNINGS**

Exacerbation of gout following therapy with probenecid may occur; in such cases colchicine or other appropriate therapy is advisable.

Probenecid increases plasma concentrations of methotrexate in both animals and humans. In animal studies, increased methotrexate toxicity has been reported. If probenecid is given with methotrexate, the dosage of methotrexate should be reduced and serum levels may need to be monitored.

In patients on probenecid the use of salicylates in either small or large doses is contraindicated because it antagonizes the uricosuric action of probenecid. The biphasic action of salicylates in the renal tubules accounts for the so-called "paradoxical effect" of uricosuric agents. In patients on probenecid who require a mild analgesic agent the use of acetaminophen rather than small doses of salicylates would be preferred.

Rarely, severe allergic reactions and anaphylaxis have been reported with the use of probenecid. Most of these have been reported to occur within several hours after readministration following prior usage of the drug.

The appearance of hypersensitivity reactions requires cessation of therapy with probenecid.

### **Use in Pregnancy**

Probenecid crosses the placental barrier and appears in cord blood. The use of any drug in women of childbearing potential requires that the anticipated benefit be weighed against possible hazards.

## **PRECAUTIONS**

### **General**

Hematuria, renal colic, costovertebral pain, and formation of uric acid stones associated with the use of probenecid in gouty patients may be prevented by alkalization of the urine and a liberal fluid intake (see DOSAGE AND ADMINISTRATION). In these cases when alkali is administered, the acid-base balance of the patient should be watched.

Use with caution in patients with a history of peptic ulcer.

Probenecid has been used in patients with some renal impairment but dosage requirements may be increased. Probenecid may not be effective in chronic renal insufficiency particularly when the glomerular filtration rate is 30 mL/minute or less. Because of its mechanism of action, probenecid is not recommended in conjunction with a penicillin in the presence of *known* renal impairment.

A reducing substance may appear in the urine of patients receiving probenecid. This disappears with discontinuance of therapy. Suspected glycosuria should be confirmed by using a test specific for glucose.

### **Drug Interactions**

When probenecid is used to elevate plasma concentrations of penicillin or other betalactams, or when

such drugs are given to patients taking probenecid therapeutically, high plasma concentrations of the other drug may increase the incidence of adverse reactions associated with that drug. In the case of penicillin or other beta-lactams, psychic disturbances have been reported.

The use of salicylates antagonizes the uricosuric action of probenecid (see WARNINGS). The uricosuric action of probenecid is also antagonized by pyrazinamide.

Probenecid produces an insignificant increase in free sulfonamide plasma concentrations but a significant increase in total sulfonamide plasma levels. Since probenecid decreases the renal excretion of conjugated sulfonamides, plasma concentrations of the latter should be determined from time to time when a sulfonamide and probenecid are coadministered for prolonged periods. Probenecid may prolong or enhance the action of oral sulfonylureas and thereby increase the risk of hypoglycemia.

It has been reported that patients receiving probenecid require significantly less thiopental for induction of anesthesia. In addition, ketamine and thiopental anesthesia were significantly prolonged in rats receiving probenecid.

The concomitant administration of probenecid increases the mean plasma elimination half-life of a number of drugs which can lead to increased plasma concentrations. These include agents such as indomethacin, acetaminophen, naproxen, ketoprofen, meclofenamate, lorazepam, and rifampin. Although the clinical significance of this observation has not been established, a lower dosage of the drug may be required to produce a therapeutic effect, and increases in dosage of the drug in question should be made cautiously and in small increments when probenecid is being co-administered. Although specific instances of toxicity due to this potential interaction have not been observed to date, physicians should be alert to this possibility.

Probenecid given concomitantly with sulindac had only a slight effect on plasma sulfide levels, while plasma levels of sulindac and sulfone were increased. Sulindac was shown to produce a modest reduction in the uricosuric action of probenecid, which probably is not significant under most circumstances.

In animals and in humans, probenecid has been reported to increase plasma concentrations of methotrexate (see WARNINGS).

Falsely high readings for theophylline have been reported in an *in vitro* study, using the Schack and Waxler technic, when therapeutic concentrations of theophylline and probenecid were added to human plasma.

## **ADVERSE REACTIONS**

The following adverse reactions have been observed and within each category are listed in order of decreasing severity.

**Central Nervous System:** headache, dizziness.

**Metabolic:** precipitation of acute gouty arthritis.

**Gastrointestinal:** hepatic necrosis, vomiting, nausea, anorexia, sore gums.

**Genitourinary:** nephrotic syndrome, uric acid stones with or without hematuria, renal colic, costovertebral pain, urinary frequency.

**Hypersensitivity:** anaphylaxis, fever, urticaria, pruritus.

**Hematologic:** aplastic anemia, leukopenia, hemolytic anemia which in some patients could be related to genetic deficiency of glucose-6-phosphate dehydrogenase in red blood cells, anemia.

**Integumentary:** dermatitis, alopecia, flushing.

## **DOSAGE AND ADMINISTRATION**

## Gout

Therapy with probenecid should not be *started* until an acute gouty attack has subsided. However, if an acute attack is precipitated *during* therapy, probenecid may be continued without changing the dosage, and full therapeutic dosage of colchicine or other appropriate therapy should be given to control the acute attack.

The recommended adult dosage is 250 mg (1/2 tablet of probenecid) twice a day for one week, followed by 500 mg (1 tablet) twice a day thereafter.

Some degree of renal impairment may be present in patients with gout. A daily dosage of 1000 mg may be adequate. However, if necessary, the daily dosage may be increased by 500 mg increments every 4 weeks within tolerance (and usually not above 2000 mg per day) if symptoms of gouty arthritis are not controlled or the 24 hour uric acid excretion is not above 700 mg. As noted, probenecid may not be effective in chronic renal insufficiency, particularly when the glomerular filtration rate is 30 mL/minute or less.

Gastric intolerance may be indicative of overdosage, and may be corrected by decreasing the dosage.

As uric acid tends to crystallize out of an acid urine, a liberal fluid intake is recommended, as well as sufficient sodium bicarbonate (3 to 7.5 g daily), or potassium citrate (7.5 g daily) to maintain an alkaline urine (see PRECAUTIONS).

Alkalinization of the urine is recommended until the serum urate level returns to normal limits and tophaceous deposits disappear, i.e., during the period when urinary excretion of uric acid is at a high level. Thereafter, alkalinization of the urine and the usual restriction of purine-producing foods may be somewhat relaxed.

Probenecid should be continued at the dosage that will maintain normal serum urate levels. When acute attacks have been absent for 6 months or more and serum urate levels remain within normal limits, the daily dosage may be decreased by 500 mg every 6 months. The maintenance dosage should not be reduced to the point where serum urate levels tend to rise.

### Probenecid and Penicillin Therapy (General)

#### Adults

The recommended dosage is 2000 mg (4 tablets of probenecid) daily in divided doses. This dosage should be reduced in older patients in whom renal impairment may be present.

#### Children 2-14 years of age

**Initial dose:** 25 mg/kg body weight ( or 0.7 g/square meter body surface).

**Maintenance dose:** 40 mg/kg body weight ( or 1.2 g/square meter body surface) per day, divided into 4 doses.

For children weighing more than 50 kg (110 lb) the adult dosage is recommended.

Probenecid is contraindicated in children under 2 years of age.

The PSP excretion test may be used to determine the effectiveness of probenecid in retarding penicillin excretion and maintaining therapeutic levels. The renal clearance of PSP is reduced to about one-fifth the normal rate when dosage of probenecid is adequate.

#### Penicillin Therapy (Gonorrhea)\*

In uncomplicated gonococcal infections in men and women (urethral, cervical, rectal), 1 g of probenecid should be given orally with 4.8 million units of aqueous procaine penicillin G\*\* (given IM), or 3 g of amoxicillin\*\* (given orally), or 3.5 g of ampicillin\*\* (given orally).

For further guidance, see CDC recommendations for definition of regimens of choice, alternative regimens, treatment of hypersensitive patients, and other aspects of therapy.

\*Recommended by the Centers for Disease Control, U.S. Department of Health and Human Services, Public Health Service (Morbidity and Mortality Weekly Report Supplement, Volume 34, Number 4S, October 18, 1985).

\*\*See package circulars of manufacturers for detailed information about CONTRAINDICATIONS, WARNINGS, PRECAUTIONS, and ADVERSE REACTIONS.

## **HOW SUPPLIED**

Probenecid Tablets USP 500 mg are bisected, capsule-shaped, yellow, film-coated tablets imprinted **DAN DAN** and **5347** supplied in bottles of 8.

Dispense in well-closed container with child-resistant closure.

**Store** at 20°-25° (68°-77°F). [See USP controlled room temperature.]

### **Protect From Light**

Manufactured By:

**Watson Pharma Private Ltd.**

Verna, Salcette Goa 403 722 INDIA

Distributed By:

**Westminster Pharmaceuticals, LLC**

154 Downing Street, Unit 1 & 2

Olive Branch, MS 38654

Revised: 08/16

1540816

0609B

## **PRINCIPAL DISPLAY PANEL**

**NDC 69367-154-00**

Probenecid Tablets USP

500 mg

8 Tablets Rx only

**Each tablet contains:**

Probenecid USP, 500 mg

**Dosage:** See package insert for dosage and full prescribing information.

**Dispense** in a well-closed container with child-resistant closure.

**Store** at 20°C-25°C (68°-77°F).[See USP controlled room temperature.]

Manufactured By:

**Watson Pharma Private Limited**

Verna, Salcette Goa 403 722 INDIA

Code No. GO/DRUGS/741 190810

Distributed By: **Westminster Pharmaceuticals, LLC**

Each Tablet Contains:  
Probenecid USP, 500 mg

Dosage: See package insert for dosage and full prescribing information.

Dispense in a well-closed container with child-resistant closure.

Store at 20°-25°C (68° -77°F). [See USP controlled room temperature.]

Protect From Light

Code No. GO/DRUGS/741 190810

**NDC 69367-154-00**  
**R<sub>x</sub> Only**

LOT: No Varnish  
EXP: .75" x .5"

**Probenecid Tablets USP** **500 mg**

**8 Tablets**

Manufactured by: Rev: 08/16  
Watson Pharma Private Limited  
Verna, Salcette Goa 403 722 India

Repackaged By:  
Westminster Pharmaceuticals, LLC  
154 Downing Street, Unit 1 & 2  
Olive Branch, MS 38654

**WP Westminster Pharmaceuticals**



## PROBENECID

probenecid tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:69367-154(NDC:0591-5347)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PROBENECID (UNII: PO572Z7917) (PROBENECID - UNII:PO572Z7917)	PROBENECID	500 mg

### Inactive Ingredients

Ingredient Name	Strength
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	2 pieces
--------------	--------	--------------	----------

<b>Shape</b>	OVAL (capsule-shaped)	<b>Size</b>	18 mm
<b>Flavor</b>		<b>Imprint Code</b>	DAN;DAN;5347
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69367-154-00	8 in 1 BOTTLE; Type 0: Not a Combination Product	07/28/2016	

**Each Tablet Contains:**  
Probenecid USP, 500 mg

**Dosage:** See package insert for dosage and full prescribing information.

**Dispense** in a well-closed container with child-resistant closure.

**Store** at 20°-25°C (68° -77°F). [See USP controlled room temperature.]

**Protect From Light**

Code No. GO/DRUGS/741 190810

**Manufactured by:** Rev: 08/16  
Watson Pharma Private Limited  
Verna, Salcette Goa 403 722 India

**Repackaged By:**  
Westminster Pharmaceuticals, LLC  
154 Downing Street, Unit 1 & 2  
Olive Branch, MS 38654

**NDC 69367-154-00**  
**Rx Only**

**Probenecid Tablets USP**

**500 mg**

**8 Tablets**

LOT:  
EXP:

No Varnish  
.75" x .5"

**WP Westminster Pharmaceuticals**

3 69367 15400 0

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA084442	07/28/2016	

**Labeler** - Westminster Pharmaceuticals, LLC (079516651)

**Registrant** - Watson Pharma Private Limited (677605709)

### Establishment

Name	Address	ID/FEI	Business Operations
Nutra-Med Packaging, Inc.		022004902	relabel(69367-154) , repack(69367-154)

### Establishment

Name	Address	ID/FEI	Business Operations
Watson Pharma Private Limited		677605709	manufacture(69367-154)