BABIES R US PAIN AND FEVER- acetaminophen suspension Toys "R"Us, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Toys "R" Us, Inc. Infants' Pain & Fever Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended does (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- push air out of syringe. Firmly push syringe into bottle opening.
- turn bottle upside down. Pull syringe to the first dose line and then push product back into bottle.
- pull syringe until it reaches and stays at the correct dose
- dispense liquid slowly into child's mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

Dosing Chart			
Weight (lb)	Age (yr)	Dose (mL)*	
under 24	under 2 years	ask a doctor	
24-35	2-3 years	5 mL	

^{*}or as directed by a doctor

Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

Other information

- store at 20-25°C (68-77°F)
- do not use if printed bottle wrap is broken or missing

Inactive ingredients

anhydrous citric acid, butylparaben, calcium sulfate, carrageenan, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions?

1-800-719-9260

Package/Label Principal Display Panel

Infants' Pain & Fever

Acetaminophen Suspension

Fever Reducer – Pain Reliever

Aspirin Free

Ibuprofen Free

Alcohol Free

Cherry Flavor

Use only with enclosed syringe

See side panel for more information

See New Dosage & Directions

Pediatrician Preferred Dosing System

Compare to the active ingredient of Infants' Tylenol® Oral Suspension

1 fl oz (30 mL) 160 mg per 5 mL





Infants' Pain & Fever

Drug Facts

Active ingredient (in each 5 mL) cetaminophen 160 mg.....

Purpose

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat

Warnings

Liver warning: This product contains acetaminophen. Severe liver

damage may occur if your child takes

■ more than 5 doses in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child has ever had an allergic reaction to this product or any of its ingredients Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin When using this product do not exceed recommended dose

(see overdose warning)

Stop use and ask a doctor if redness or swelling is present

■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days new symptoms occur These could be signs of a serious condition

Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions this product does not contain directions or

- complete warnings for adult use

 do not give more than directed (see overdose warning)

 shake well before using
 m.l.=milliliter

 find right dose on chart. If possible, use weight to dose; otherwise,
- push air out of syringe. Firmly push syringe into bottle opening. ■ turn bottle upside down. Pull syringe to the first dose line and then push product back into bottle

Drug Facts (continued)

- pull syringe until it reaches and stays at the correct dose
- dispense liquid slowly into child's mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in
- 24 hours ■ replace cap tightly to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*	
under 24	under 2 years	ask a doctor	
24-35	2-3 years	5 mL	
According to the state of the s			

or as directed by a doctor Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

Other information

 store at 20-25°C (68-77°F) ■ do not use if printed bottle wrap is broken or missing

Inactive ingredients anhydrous citric acid, butylparaben, calcium sulfate carrageenan, FD&C red #40, flavor, glycerin, high fructose com syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions? 1-800-719-9260

www.babiesrus.com

BABIES "R" US is a registered trademark of Geoffrey, LLC. Part of the "R US" Family of Brands. © 2010 Geoffrey, LLC. All rights reserved. Distributed by Toys "R" Us, Inc., Wayne, NJ 07470

NDC 51334-161-10

BABIES

Fever Reducer - Pain Reliever

Aspirin Free Ibuprofen Free Alcohol Free



Cherry Flavor


See side panel for more See New Dosage &

Directions

Pediatrician Preferred Dosing System

Compare to the active ingredient of Infants' Tylenol® Oral Suspension*

1 fl oz (30 mL) 160 mg per 5 mL



STEP 2. Shake well before using.

STEP 3. Push air out of syringe. Firmly push syringe into bottle opening.



STEP 4. Turn the bottle upside down. Pull syringe to the first dose line and then push product back into bottle.

STEP 5. Pull syringe until it reach stays at the correct dose



IMPORTANT:

New Dosage and Directions. Please see dosing chart or ask a doctor for your child's correct dose

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenok® Oral Suspension.





LOT NO.

EXP.

: 19110 MC CS



BABIES R US PAIN AND FEVER

acetaminophen suspension

Product Information

Product Type HUMAN OTC DRUG LABEL Item Code (Source) NDC:51334-161 Route of Administration ORAL DEA Schedule

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength **ACETAMINO PHEN** (ACETAMINO PHEN) ACETAMINOPHEN $160 \ mg \ in \ 5 \ mL$

Inactive Ingredients

Ingredient Name	Strength
ANHYDRO US CITRIC ACID	
BUTYLPARABEN	
CALCIUM SULFATE	
CARRAGEENAN	
FD&C RED NO. 40	
GLYCERIN	
HIGH FRUCTO SE CORN SYRUP	
CELLULOSE, MICRO CRYSTALLINE	
CARBO XYMETHYLCELLULO SE SO DIUM	
PROPYLENE GLYCOL	
WATER	
SO DIUM BENZO ATE	
SORBITOL	
SO DIUM PHO SPHATE, TRIBASIC	

Product Characteristics	roduct Characteristics			
Color	RED	Score		
Shape		Size		
Flavor		Imprint Code		
Contains				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:51334-161-10	1 in 1 CARTON			
1		30 mL in 1 BOTTLE			
2	NDC:51334-161-16	1 in 1 CARTON			
2		59 mL in 1 BOTTLE			

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph not final	part343	11/29/2011			

Labeler - Toys"R"Us, Inc. (006985808)

Revised: 1/2014 Toys"R"Us, Inc.