

BENZONATATE- benzonatate capsule
Northwind Pharmaceuticals

Benzonatate

DESCRIPTION

Benzonatate, a non-narcotic oral antitussive agent, is 2, 5, 8, 11, 14, 17, 20, 23, 26-nonaoxaoctacosan-28-yl p-(butylamino) benzoate. The molecular formula is C₃₀H₅₃NO₁₁ with a molecular weight of 603.7

Each capsule, for oral administration, contains 200 mg of benzonatate, USP.

Benzonatate Capsules also contain: Alcohol, ammonium hydroxide, D&C Yellow 10, gelatin, glycerin, propylene glycol, purified water, shellac glaze, simethicone and titanium dioxide. In addition, the capsule may contain trace amounts of fractionated coconut oil.

CLINICAL PHARMACOLOGY

Benzonatate acts peripherally by anesthetizing the stretch receptors located in the respiratory passages, lungs, and pleura by dampening their activity and thereby reducing the cough reflex at its source. It begins to act within 15 to 20 minutes and its effect lasts for 3 to 8 hours. Benzonatate capsules have no inhibitory effect on the respiratory center in recommended dosage.

INDICATIONS & USAGE

Benzonatate capsules are indicated for the symptomatic relief of cough.

CONTRAINDICATIONS

Hypersensitivity to benzonatate or related compounds.

WARNINGS

Hypersensitivity

Severe hypersensitivity reactions (including bronchospasm, laryngospasm and cardiovascular collapse) have been reported which are possibly related to local anesthesia from sucking or chewing the capsule instead of swallowing it. Severe reactions have required intervention with vasopressor agents and supportive measures.

Psychiatric Effects

Isolated instances of bizarre behavior, including mental confusion and visual hallucinations, have also been reported in patients taking benzonatate capsules in combination with other prescribed drugs.

Accidental Ingestion and Death in Children

Keep benzonatate capsules out of reach of children. Accidental ingestion of benzonatate capsules resulting in death has been reported in children below age 10. Signs and symptoms of overdose have been reported within 15-20 minutes and death has been reported within one hour of ingestion. If accidental ingestion occurs, seek medical attention immediately (see OVERDOSAGE).

PRECAUTIONS

Benzonatate is chemically related to anesthetic agents of the para-amino-benzoic acid class (e.g. procaine; tetracaine) and has been associated with adverse CNS effects possibly related to a prior sensitivity to related agents or interaction with concomitant medication.

Information for Patients:

Swallow benzonatate capsules whole. Do not break, chew, dissolve, cut or crush benzonatate capsules. Release of benzonatate from the capsule in the mouth can produce a temporary local anesthesia of the oral mucosa and choking could occur. If numbness or tingling of the tongue, mouth, throat, or face occurs, refrain from oral ingestion of food or liquids until the numbness has resolved. If the symptoms worsen or persist, seek medical attention.

Keep benzonatate capsules out of reach of children. Accidental ingestion resulting in death has been reported in children. Signs and symptoms of overdose have been reported within 15-20 minutes and death has been reported within one hour of ingestion. Signs and symptoms may include restlessness, tremors, convulsions, coma and cardiac arrest. If accidental ingestion occurs, seek medical attention immediately.

Overdosage resulting in death may occur in adults.

Do not exceed a single dose of 200 mg and a total daily dosage of 600 mg. If you miss a dose of benzonatate capsules, skip that dose and take the next dose at the next scheduled time. Do not take 2 doses of benzonatate capsules at one time.

Usage in Pregnancy:

Pregnancy Category C. Animal reproduction studies have not been conducted with benzonatate. It is also not known whether benzonatate can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Benzonatate capsules should be given to a pregnant woman only if clearly needed.

Nursing Mothers:

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk caution should be exercised when benzonatate capsules are administered to a nursing woman.

Carcinogenesis, Mutagenesis, Impairment of Fertility:

Carcinogenicity, mutagenicity, and reproduction studies have not been conducted with benzonatate.

Pediatric Use:

Safety and effectiveness in children below the age of 10 have not been established. Accidental ingestion resulting in death has been reported in children below age 10. Keep out of reach of children.

ADVERSE REACTIONS

Potential Adverse Reactions to benzonatate capsules may include:

Hypersensitivity reactions including bronchospasm, laryngospasm, cardiovascular collapse possibly related to local anesthesia from chewing or sucking the capsule.

CNS: sedation; headache; dizziness; mental confusion; visual hallucinations.

GI: constipation; nausea; GI upset.

Dermatologic: pruritus; skin eruptions.

Other: nasal congestion; sensation of burning in the eyes; vague "chilly" sensation; numbness of the chest; hypersensitivity.

Deliberate or accidental overdose has resulted in death, particularly in children.

OVERDOSAGE

Intentional and unintentional overdose may result in death, particularly in children.

The drug is chemically related to tetracaine and other topical anesthetics and shares various aspects of their pharmacology and toxicology. Drugs of this type are generally well absorbed after ingestion.

Signs and Symptoms:

The signs and symptoms of overdose of benzonatate have been reported within 15-20 minutes. If capsules are chewed or dissolved in the mouth, oropharyngeal anesthesia will develop rapidly, which may cause choking and airway compromise.

CNS stimulation may cause restlessness and tremors which may proceed to clonic convulsions followed by profound CNS depression. Convulsions, coma, cerebral edema and cardiac arrest leading to death have been reported within 1 hour of ingestion.

Treatment:

In case of overdose, seek medical attention immediately. Evacuate gastric contents and administer copious amounts of activated charcoal slurry. Even in the conscious patient, cough and gag reflexes may be so depressed as to necessitate special attention to protection against aspiration of gastric contents and orally administered materials. Convulsions should be treated with a short-acting barbiturate given intravenously and carefully titrated for the smallest effective dosage. Intensive support of respiration and cardiovascular-renal function is an essential feature of the treatment of severe intoxication from overdose.

Do not use CNS stimulants.

DOSAGE & ADMINISTRATION

Adults and Children over 10 years of age: Usual dose is one 100 mg or 200 mg capsule three times a day as needed for cough. If necessary to control cough, up to 600 mg daily in three divided doses may be given. Benzonatate Capsules should be swallowed whole. Benzonatate Capsules are not to be broken, chewed, dissolved, cut or crushed.

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC: 51655-771-20

MFG: 57664-134-88

Benzonatate 200 MG

20 Capsules

Rx Only

Lot#:

Exp. Date:

Each soft gelatin capsule contains Benzonatate, USP....200 MG

Dosage: See accompanying prescribing information

Store at 68 to 77 degrees F.

Keep tightly closed.

Protect from light. Keep out of the reach of children.

Medication guide is found at www.fda.gov/drugs/drugsafety/ucm/085729

Distributed by: Caraco Pharmaceutical Laboratories, Detroit, MI 48202 Lot#

Repackaged by Northwind Pharmaceuticals, Indianapolis, IN 46256

NDC: 51655-771-20 LCN# 01
MFG: 57664-134-88 Rev. 2

Benzonatate 200MG

20 Capsules
Rx Only

Lot #: NW73010005
Exp. Date: 11/2015

Each soft gelatin capsule contains:
Benzonatate, USP....200MG
Dosage: See accompanying
prescribing information

Store at 68 to 77 degrees F.
Keep tightly closed. Protect from
light. Keep out of the reach of
children.

Medication guide is found at
www.fda.gov/drugs/drugsafety/ucm085729

Distributed by: Caraco Pharmaceutical
Laboratories, Detroit, MI 48202 Lot# 3308 026A

Repackaged By: Northwind Pharmaceuticals,
Indianapolis, IN 46256



NDC: 51655-771-52

MFG: 57664-134-88

Benzonatate 200 MG

30 Capsules

Rx Only

Lot#:

Exp. Date:

Each soft gelatin capsule contains Benzonatate, USP....200 MG

Dosage: See accompanying prescribing information

Store at 68 to 77 degrees F.


Keep tightly closed.

Protect from light. Keep out of the reach of children.

Medication guide is found at www.fda.gov/drugs/drugsafety/ucm/085729

Distributed by: Caraco Pharmaceutical Laboratories, Detroit, MI 48202 Lot#

Repackaged by Northwind Pharmaceuticals, Indianapolis, IN 46256



BENZONATATE

benzonatate capsule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:51655-771(NDC:57664-134)
Route of Administration	Oral		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
benzonatate (UNII: 5P4DHS6ENR) (BENZONATATE - UNII:5P4DHS6ENR)	benzonatate	200 mg

Product Characteristics

Color	yellow	Score	no score
Shape	capsule	Size	5mm
Flavor		Imprint Code	134
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51655-771-20	20 in 1 BOTTLE, DISPENSING		
2	NDC:51655-771-52	30 in 1 BOTTLE, DISPENSING		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040587	01/15/2015	

Labeler - Northwind Pharmaceuticals (036986393)

Registrant - Northwind Pharmaceuticals (036986393)

Establishment

Name	Address	ID/FEI	Business Operations
Northwind Pharmaceuticals		036986393	repack(51655-771)

Revised: 4/2015

Northwind Pharmaceuticals