PERFECT CHOICE- sodium fluoride and hydrofluoric acid gel Young Dental Manufacturing Co 1, LLC.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

APF - Peppermint Stick

INDICATIONS AND USAGE:

Topical application of APF gel is indicated in the prevention of dental caries.

Warning:

Keep out of reach of children. Do not swallow.

If swallowed during treatment, administer milk, limewater, or antacid. If larger dose is ingested (more than 1/4 ounce), contact poison control and immediately seek medical help. Do not use on patients with an allergy to fluoride.

Precautions:

For professional use only. This product is not intended for home or unsupervised consumer use. Repeated use may cause dulling of porcelain and ceramic restorations.

Adverse reactions:

Developing teeth of children under age 6 may become permanently discolored if excessive amounts are repeatedly swallowed. Individuals hypersensitive to fluoride may have an adverse reaction.

DOSAGE AND ADMINISTRATION:

After thorough prophylaxis, place gel in applicator tray (not more than 1/3 full). Throughly air-dry teeth and insert tray(s) in mouth. Gel may also be swab applied. Instruct patient to bite down slightly but firmly for one to four minutes. A slight chewing motion enhances coverage interproximally. Remove tray(s) and have patient expectorate excess gel. Instruct patient not to eat, drink or rinse for at least 30 minutes.

HOW SUPPLIED:

A thixotropic gel, one bottle contains up to 68 individual applications. Store at controlled room temperature 20°- 25°C (68°-77°F).

Do not allow to freeze.

ACTIVE INGREDIENTS:

1.23% w/w Fluoride Ion (0.79% from Sodium Fluoride and 0.44% from Hydrogen Fluoride) in 0.1 Molar Phosphoric Acid.

Contains: FD&C Red #40

Principal Display Panel

NDC 0273-7006-16

 $ZOOBY \\ \mathbb{R}$

PERFECT CHOICE®

1.23% APF TOPICAL GEL

STRIPED

PEPPERMINT

STICK

60-Second

Treatment Acidulated

Phosphate Fluoride

1.23% Fluoride Ion

Net Weight: 17.5 oz. (497g)



NDC 0273-7006-16

control and immediately seek medical help. Do not use on patients If swallowed during treatment, administer milk, limewater, or antacid If larger dose is ingested (more than 1/4 ounce), contact poison Warning: Keep out of reach of children. Do not swallow.

intended for home or unsupervised consumer use. Repeated use may **Precautions:** For professional use only. This product is not cause dulling of porcelain and ceramic restorations.

6 may become permanently discolored if excessive amounts are have an adverse reaction. repeatedly swallowed. Individuals hypersensitive to fluoride may Adverse reactions: Developing teeth of children under age

DOSAGE AND ADMINISTRATION:

Affer thorough prophylaxis, place gel in applicator tray (not more than 1/3 full). Thoroughly air-dry teeth and insert tray(s) in mouth

1.23% APF TOPICAL GEL

PERFECT CHOICE®

HOW SUPPLIED:

Store at controlled room temperature 20°-25°C (68°-77°F). A thixotropic gel, one bottle contains up to 68 individual applications.

Do not allow to freeze.

ACTIVE INGREDIENTS:

from Hydrogen Fluoride) in 0,1 Molar Phosphoric Acid .23% w/w Fluoride Ion (0.79% from Sodium Fluoride and 0.44%

Contains: FD&C Red #40.

INDICATIONS AND USAGE:

dental caries. Topical application of APF gel is indicated in the prevention of

with an allergy to fluoride.

interproximal coverage. Remove tray(s) and have patient expectorate Gel may also be swab applied. Instruct patient to bite down slightly 30 minutes excess gel. Instruct patient not to eat, drink or rinse for at least but firmly for one to four minutes. A slight chewing motion enhances



Phosphate Fluoride

.23% Fluoride Ion

Ireatment Acidulated

60-Second

STICK

PEPPERMINT

STRIPED

TIME TO REORDER! CALL YOUR DENTAL DEALER.



www.ZoobyDental.com Earth City, MO 63045 (800)227-3321 manufactured for: **BIOTROL**[™]



X only

12000235C - 05/15

REF APF 205PS





NET WEIGHT: 17.5 oz. (497g)



sodium fluoride and hydrofluoric acid gel

Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0273-7006	
Route of Administration	DENTAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
SODIUM FLUORIDE (UNII: 8ZYQ1474W7) (FLUORIDE ION - UNII:Q80 VPU408O)	FLUORIDE ION	7.9 mg in 1 g	
HYDROFLUORIC ACID (UNII: RGL5YE86CZ) (FLUORIDE ION - UNII:Q80VPU408O)	FLUORIDE ION	4.4 mg in 1 g	

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	PEPPERMINT (Peppermint Stick)	Imprint Code	
Contains			

ı	Packaging			
ı	# Item Code	Package Description	Marketing Start Date	Marketing End Date
ı	1 NDC:0273-7006-16	497 g in 1 BOTTLE; Type 0: Not a Combination Product	12/22/2018	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
unapproved drug other		12/22/2018		

Labeler - Young Dental Manufacturing Co 1, LLC. (006309355)

Registrant - Medical Products Laboratories, Inc (002290302)

Establishment					
Name	Address	ID/FEI	Business Operations		
Medical Products Laboratories, Inc		002290302	manufacture(0273-7006)		

Revised: 12/2018 Young Dental Manufacturing Co 1, LLC.