

**IBUPROFEN - ibuprofen tablet**  
**MARKSANS PHARMA LIMITED**

-----  
**Ibuprofen Tablets 200mg (OTC; Brown Tablets & Caplets)**

**Active ingredient(s)**

Ibuprofen 200 mg (NSAID)\*

\* nonstreoidal anti-inflammatory drug

**Purpose**

Pain reliever / fever reducer

**Use(s)**

temporarily relieves minor aches and pains due to :

- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps

temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergy reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAID's except aspirin increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask doctor if**

you experience any of the following signs of stomach bleeding:

- feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days  
 fever gets worse or lasts more than 3 days  
 redness or swelling is present in the painful area  
 any new symptoms appear

### **If pregnant or breast-feeding**

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause a problems in the unborn child or complications during delivery.

### **Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and older	<ul style="list-style-type: none"> <li>▪ take 1 tablet every 4 to 6 hours while symptoms persist</li> <li>▪ if pain or fever does not respond to 1 tablet, 2 tablets may be used</li> <li>▪ do not exceed 6 tablets in 24 hours, unless directed by a doctor.</li> </ul>
children under 12 years	<ul style="list-style-type: none"> <li>▪ ask a doctor</li> </ul>

### **Other information**

- store between 20-25 °C (68-77 °F).
- **do not use if carton is opened. do not use if seal under bottle cap imprinted with” SEALED for YOUR PROTECTION” is broken or missing.**
- see end panel for lot number and expiration date

### **Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

### ***Questions or Comments?***

Call 1-877-290-4008

Manufactured for:

**Time-Cap Labs, Inc.**

7 Michael Avenue,  
Farmingdale,  
NY 11735, USA

Manufactured by:

**Marksans Pharma Ltd.**

Plot No. L-82, L-83  
Verna Industrial Estate  
Verna, Goa-403722, India

### **Storage**

### **Principal Display Panel**

NDC 25000-114-03, Ibuprofen Tablets USP, 200mg

30 counts Bottle Label



NDC 25000-114-03

# Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer  
(NSAID)

30 Coated Tablets



Marksans

**Important: Read all product information before using.  
Keep the box for important information.**

## Active ingredient (in each tablet)

## Purpose

Ibuprofen 200 mg (NSAID) \* ..... Pain reliever/fever reducer  
\* nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to:  
■ headache ■ muscular aches ■ minor pain of arthritis ➔

LOT :

EXP :

R912/21

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. ➔

Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This ➔

product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you : ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ➔

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than ➔

directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ ➔

if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel ➔

faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part ➔

■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. ➔

a health professional before use. **Keep out of reach of children. Directions :** ■ do not take more than directed ■ the smallest effective dose should be used. adults and children 12 years and ➔

or side of body ■ slurred speech ➔	If pregnant or breast-feeding, ask ➔	older : ■ take 1 tablet every 4 to 6 ➔
------------------------------------	--------------------------------------	--

hours while symptoms persist ■  
 if pain or fever does not respond to  
 1 tablet, 2 tablets may be used ■  
 do not exceed 6 tablets in 24  
 hours, unless directed by a doctor.  
 children under 12 years: ■ ask a  
 doctor. **Other information** ■ store ➔

between 20-25°C (68-77°F)

***Do not use if seal under bottle  
 cap imprinted with "SEALED  
 for YOUR PROTECTION" is  
 broken or missing.***


Mfg. Lic. No. : GO/DRUGS/515 ➔

Manufactured for:  
**Time-Cap Labs, Inc.**  
 7 Michael Avenue, Farmingdale,  
 NY 11735, USA

Manufactured by:  
**Marksans Pharma Ltd.**  
 Plot No. L-82, L-83, Verna Indl. Estate,  
 Verna, GOA - 403722, India.

NDC 25000-114-03, Ibuprofen Tablets USP, 200mg

30 counts Carton Label



NDC 25000-117-03, Ibuprofen Tablets USP, 200mg (Caplets)

30 counts Bottle Label



NDC 25000-117-03

# Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer  
(NSAID)

**30 Coated Caplets\***

\* Capsule-Shaped Tablets



Marksans

**Important: Read all product information before using.  
Keep the box for important information.**

## **Active ingredient (in each caplet)**

Ibuprofen 200 mg (NSAID) \* ..... Pain reliever/fever reducer  
\* nonsteroidal anti-inflammatory drug

## **Purpose**

**Uses** ■ temporarily relieves minor aches and pains due to:  
■ headache ■ muscular aches ■ minor pain of arthritis ➔

LOT :

EXP :

R912/21

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. ➔

Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This ➔

product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you : ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ➔

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than ➔

directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ ➔

if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel ➔

faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part ➔

■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. ➔

a health professional before use. **Keep out of reach of children. Directions :** ■ do not take more than directed ■ the smallest effective dose should be used. adults and children 12 years and ➔

or side of body ■ slurred speech ➔

If pregnant or breast-feeding, ask ➔

older : ■ take 1 caplet every 4 to 6 ➔

hours while symptoms persist ■  
if pain or fever does not respond to  
1 caplet, 2 caplets may be used ■  
do not exceed 6 caplets in 24  
hours, unless directed by a doctor.  
children under 12 years: ■ ask a  
doctor. **Other information** ■ store ➔

between 20-25°C (68-77°F)

***Do not use if seal under bottle  
cap imprinted with "SEALED  
for YOUR PROTECTION" is  
broken or missing.***


Mfg. Lic. No. : GO/DRUGS/515 ➔

Manufactured for:  
**Time-Cap Labs, Inc.**  
7 Michael Avenue, Farmingdale,  
NY 11735, USA

Manufactured by:  
**Marksans Pharma Ltd.**  
Plot No. L-82, L-83, Verna Indl. Estate,  
Verna, GOA - 403722, India.

NDC 25000-117-03, Ibuprofen Tablets USP, 200mg (Caplets)

30 counts Carton Label



NDC 25000-114-70, Ibuprofen Tablets USP, 200mg

500 counts Bottle Label



NDC 25000-114-70

# Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer (NSAID)

**THIS PACKAGE FOR HOUSEHOLDS  
WITHOUT YOUNG CHILDREN**



**500 Coated Tablets**  
**Marksans**

## Drug Facts

**Active ingredient (in each tablet)** **Purpose**  
Ibuprofen 200 mg (NSAID)\* Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

## Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

## Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include :

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

LOT :

EXP :

R11/21



## Drug Facts (Continued)

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you :

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAID's except aspirin, increases the risk of heart attack, heart failure, and stroke.

## Drug Facts (Continued)

These can be fatal. The risk is higher if you use more than directed or for longer than directed.

## Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

## Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.
- you are taking a diuretic

## Drug Facts (Continued)

### Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

## When using this product

- take with food or milk if stomach upset occurs

## Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part

## Drug Facts (Continued)

- or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Drug Facts (Continued)

## Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children  
12 years and older

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years

- ask a doctor

## Other information

- store between 20-25°C (68-77°F).
- do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

## Drug Facts (Continued)

## Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

## Questions or comments?

Call 1-877-290-4008

Manufactured for: **Time-Cap Labs, Inc.**  
7 Michael Avenue, Farmingdale, NY 11735, USA

Manufactured by: **Marksans Pharma Ltd.**  
Plot No. L-82, L-83, Verna Indl. Estate,  
Verna, GOA - 403722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-114-14, Ibuprofen Tablets USP, 200mg

1000 counts Bottle Label





NDC 25000-114-14

# Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer  
(NSAID)

**THIS PACKAGE FOR HOUSEHOLDS  
WITHOUT YOUNG CHILDREN**

**1000 Coated Tablets**



Marksans

## Drug Facts

**Active ingredient (in each tablet)**      **Purpose**  
Ibuprofen 200 mg (NSAID)\* Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

### Uses

- temporarily relieves minor aches and pains due to:
  - headache ■ muscular aches ■ minor pain of arthritis ■ toothache ■ backache ■ the common cold ■ menstrual cramps
- temporarily reduces fever

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include :

- hives ■ facial swelling ■ asthma (wheezing)
- shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.


The chance is higher if you :

- are age 60 or older

LOT :

EXP :

R912/21



## Drug Facts (Continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

### Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** →

## Drug Facts (Continued)

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

### When using this product

- take with food or milk if stomach upset occurs

### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint ■ vomit blood ■ have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. →

## Drug Facts (Continued)

## Drug Facts (Continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- do not take more than directed
- the smallest effective dose should be used

adults and children  
12 years and older

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor.

children under 12 years

- ask a doctor

**Other information**

- store between 20-25°C (68-77°F).
- do not use if carton is opened. Do not use if seal under bottle cap imprinted with “SEALED for YOUR PROTECTION” is broken or missing

**Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol,

polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

**Questions or comments?**

Call 1-877-290-4008

Manufactured for:  
**Time-Cap Labs, Inc.**  
7 Michael Avenue,  
Farmingdale,  
NY 11735, USA

Manufactured by:  
**Marksans Pharma Ltd.**  
Plot No. L-82, L-83,  
Verna Indl. Estate,  
Verna, GOA - 403722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-114-71, Ibuprofen Tablets USP, 200mg

1000 counts (PET) Bottle Label

NDC 25000-114-71

# Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer (NSAID)

THIS PACKAGE FOR HOUSEHOLDS  
WITHOUT YOUNG CHILDREN

1000 Coated Tablets



Marksans

## Drug Facts

**Active ingredient (in each tablet)** **Purpose**  
Ibuprofen 200 mg (NSAID)\* Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

### Uses

■ temporarily relieves minor aches and pains due to:  
■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache ■ backache ■ the common cold ■ menstrual cramps  
■ temporarily reduces fever

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

■ hives ■ facial swelling ■ asthma (wheezing)  
■ shock ■ skin reddening ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

## Drug Facts (Continued)

or others)  
■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

**Heart attack and stroke warning:** NSAID's except aspirin, increases the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### Do not use

■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer  
■ right before or after heart surgery

### Ask a doctor before use if

■ you have problems or serious side effects from taking pain relievers or fever reducers  
■ the stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.  
■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

LOT :

EXP :

R11221



## Drug Facts (Continued)

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
■ under a doctor's care for any serious condition  
■ taking any other drug

### When using this product

■ take with food or milk if stomach upset occurs

### Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools  
■ have stomach pain that does not get better  
■ you have symptoms of heart problem or stroke:  
■ chest pain ■ trouble breathing ■ weakness in one part or side of body  
■ slurred speech ■ leg swelling  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Drug Facts (Continued)

### Directions

■ do not take more than directed  
■ the smallest effective dose should be used

adults and children  
12 years and older

■ take 1 tablet every 4 to 6 hours while symptoms persist  
■ if pain or fever does not respond to 1 tablet, 2 tablets may be used  
■ do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years

■ ask a doctor

### Other information

■ store between 20-25°C (68-77°F).  
■ do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing

### Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

### Questions or comments?

Call 1-877-290-4008

## Drug Facts (Continued)

Manufactured for:  
**Time-Cap Labs, Inc.**  
7 Michael Avenue,  
Farmingdale,  
NY 11735, USA

Manufactured by:  
**Marksans Pharma Ltd.**  
Plot No. L-82, L-83,  
Verna Indl. Estate,  
Verna, GOA - 403722, India.

Mfg. Lic. No.: GO/DRUGS/515

NDC 25000-117-14, Ibuprofen Tablets USP, 200mg

1000 counts Bottle Label (Caplets)



NDC 25000-117-14

# Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer  
(NSAID)

**THIS PACKAGE FOR HOUSEHOLDS  
WITHOUT YOUNG CHILDREN**

**1000 Coated Caplets\***  
\*Capsule-Shaped Tablets



Marksans

## Drug Facts

**Active ingredient (in each caplet)** **Purpose**  
Ibuprofen 200 mg (NSAID)\* Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.


The chance is higher if you:

- are age 60 or older

LOT:

EXP:

R912/21



## Drug Facts (Continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

### Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

## Drug Facts (Continued)

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

### When using this product

- take with food or milk if stomach upset occurs

### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## Drug Facts (Continued)

## Drug Facts (Continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- do not take more than directed
- the smallest effective dose should be used

adults and children  
12 years and older

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor.

children under 12 years

- ask a doctor

**Other information**

- store between 20-25°C (68-77°F).
- do not use if carton is opened. Do not use if seal under bottle cap imprinted with “SEALED for YOUR PROTECTION” is broken or missing

**Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol,

polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

**Questions or comments?**

Call 1-877-290-4008

Manufactured for:  
**Time-Cap Labs, Inc.**  
7 Michael Avenue,  
Farmingdale,  
NY 11735, USA

Manufactured by:  
**Marksans Pharma Ltd.**  
Plot No. L-82, L-83,  
Verna Indl. Estate,  
Verna, GOA - 403722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-117-69, Ibuprofen Tablets USP, 200mg (Caplets)

300 counts Bottle Label

NDC 25000-117-69

# Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer (NSAID)

**THIS PACKAGE FOR HOUSEHOLDS  
WITHOUT YOUNG CHILDREN**

**300 Coated Caplets\***

\* Capsule-Shaped Tablets

**Marksans**



## Drug Facts

**Active ingredient (in each caplet)** **Purpose**  
Ibuprofen 200 mg (NSAID)\* Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

## Uses

- temporarily relieves minor aches and pains due to:
  - headache ■ muscular aches ■ minor pain of arthritis ■ toothache ■ backache ■ the common cold ■ menstrual cramps
- temporarily reduces fever

## Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.


Symptoms may include :

- hives ■ facial swelling ■ asthma (wheezing) ■ shock
- skin reddening ■ rash ■ blisters

LOT :

EXP :

R11/21



## Drug Facts (Continued)

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you :

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAID's except aspirin, increases the risk of heart attack, heart failure, and stroke.

## Drug Facts (Continued)

These can be fatal. The risk is higher if you use more than directed or for longer than directed.

## Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

## Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.
- you are taking a diuretic

## Drug Facts (Continued)

### Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

## When using this product

- take with food or milk if stomach upset occurs

## Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint ■ vomit blood ■ have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
  - chest pain ■ trouble breathing ■ weakness in one part

## Drug Facts (Continued)

- or side of body ■ slurred speech ■ leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Drug Facts (Continued)

## Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children  
12 years and older

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor.

Children under 12 years

- ask a doctor

## Other information

- store between 20-25°C (68-77°F).
- do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

## Drug Facts (Continued)

## Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

## Questions or comments?

Call 1-877-290-4008

Manufactured for: **Time-Cap Labs, Inc.**  
7 Michael Avenue, Farmingdale, NY 11735, USA

Manufactured by: **Marksans Pharma Ltd.**  
Plot No. L-82, L-83, Verna Indl. Estate,  
Verna, GOA - 403722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-114-02, Ibuprofen Tablets USP, 200mg

24 counts Bottle Label



NDC 25000-114-02

## Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer  
(NSAID)

24 Coated Tablets



Marksans

**Important: Read all product information before using.  
Keep the box for important information.**

### **Active ingredient (in each tablet)**

Ibuprofen 200 mg (NSAID)\* ..... Pain reliever/fever reducer

\*nonsteroidal anti-inflammatory drug

### **Purpose**

**Uses** ■ temporarily relieves minor aches and pains due to:

■ headache ■ muscular aches ■ minor pain of arthritis ➔

LOT :

EXP :

R912/21

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. ➔

Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This ➔

product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you : ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ➔

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than ➔

directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ ➔

if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel ➔

faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part ➔

■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear ➔

a health professional before use. **Keep out of reach of children. Directions :** ■ do not take more than directed ■ the smallest effective dose should be used. adults and children 12 years and ➔

breathing ■ weakness in one part  
or side of body ■ slurred speech ➔

new symptoms appear.

**If pregnant or breast-feeding, ask** ➔

adults and children 12 years and  
older : ■ take 1 tablet every 4 to 6 ➔

hours while symptoms persist ■  
if pain or fever does not respond to  
1 tablet, 2 tablets may be used ■  
do not exceed 6 tablets in 24  
hours, unless directed by a doctor.  
children under 12 years: ■ ask a  
doctor. **Other information** ■ store ➔

between 20-25°C (68-77°F)

***Do not use if seal under bottle  
cap imprinted with "SEALED  
for YOUR PROTECTION" is  
broken or missing.***

Mfg. Lic. No. : GO/DRUGS/515 ➔

Manufactured for:

**Time-Cap Labs, Inc.**

7 Michael Avenue, Farmingdale,  
NY 11735, USA

Manufactured by:


**Marksans Pharma Ltd.**

Plot No. L-82, L-83, Verna Indl. Estate,  
Verna, GOA - 403722, India.

NDC 25000-114-02, Ibuprofen Tablets USP, 200mg


24 counts Carton Label





NDC 25000-117-02, Ibuprofen Tablets USP, 200mg (Caplets)

24 counts Bottle Label



■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. ➔

Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This ➔

product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you : ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ➔

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than ➔

directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ ➔

if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel ➔

faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ➔

■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding,** ask ➔

a health professional before use. **Keep out of reach of children. Directions :** ■ do not take more than directed ■ the smallest effective dose should be used. adults and children 12 years and older : ■ take 1 caplet every 4 to 6 ➔

hours while symptoms persist ■ if pain or fever does not respond to ➔

between 20-25°C (68-77°F)

© 2014 Time-Cap Labs, Inc. All rights reserved.

Manufactured for:  
**Time-Cap Labs, Inc.**  
7 Michael Avenue, Farmingdale



# IBUPROFEN

ibuprofen tablet

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:25000-114
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	

## Product Characteristics

Color	BROWN	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	114
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-114-03	30 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	
2	NDC:25000-114-14	1000 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	
3	NDC:25000-114-20	4 in 1 BOX	07/25/2011	
3		6500 in 1 BAG; Type 0: Not a Combination Product		
4	NDC:25000-114-	6 in 1 BOX	07/25/2011	

4	30	6500 in 1 BAG; Type 0: Not a Combination Product	07/25/2011	
4		6500 in 1 BAG; Type 0: Not a Combination Product		
5	NDC:25000-114-98	1 in 1 BOX	07/25/2011	
5		34065 in 1 BAG; Type 0: Not a Combination Product		
6	NDC:25000-114-93	1 in 1 BOX	07/25/2011	
6		37470 in 1 BAG; Type 0: Not a Combination Product		
7	NDC:25000-114-70	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	
8	NDC:25000-114-71	1000 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	
9	NDC:25000-114-02	1 in 1 CARTON	12/20/2019	
9		24 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091239	07/25/2011	

## IBUPROFEN

ibuprofen tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:25000-117
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	



<b>POLYVINYL ALCOHOL</b> (UNII: 532B59J990)	
<b>POLYETHYLENE GLYCOL</b> (UNII: 3WJQ0SDW1A)	

### Product Characteristics

<b>Color</b>	BROWN	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	15mm
<b>Flavor</b>		<b>Imprint Code</b>	117
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-117-03	30 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	
2	NDC:25000-117-14	1000 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	
3	NDC:25000-117-20	4 in 1 BOX	07/25/2011	
3		6500 in 1 BAG; Type 0: Not a Combination Product		
4	NDC:25000-117-30	6 in 1 BOX	07/25/2011	
4		6500 in 1 BAG; Type 0: Not a Combination Product		
5	NDC:25000-117-98	1 in 1 BOX	07/25/2011	
5		34065 in 1 BAG; Type 0: Not a Combination Product		
6	NDC:25000-117-93	1 in 1 BOX	07/25/2011	
6		37470 in 1 BAG; Type 0: Not a Combination Product		
7	NDC:25000-117-69	300 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	
8	NDC:25000-117-02	1 in 1 CARTON	12/20/2019	
8		24 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091239	07/25/2011	

**Labeler -** MARKSANS PHARMA LIMITED (925822975)

### Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

MARKSANS PHARMA LIMITED		925822975	LABEL(25000-117) , MANUFACTURE(25000-114, 25000-117)
-------------------------	--	-----------	--

Revised: 12/2021

MARKSANS PHARMA LIMITED