NAPROXEN SODIUM- naproxen sodium tablet Pharmacy Valuce Alliance, LLC

Naproxen Sodium Tablets USP 220mg (Round)

Active ingredient

(in each tablet)

Naproxen sodium 220 mg (NSAID)**

**nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

• take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing

- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

adults and children 12 years and older,

- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8- to 12- hour period
- do not exceed 3 tablets in a 24- hour period

children under 12 years,

■ ask a doctor

Other information


- each tablet contains: sodium 20 mg
- store at 20 25°C (68 to 77°F). Avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients

FD&C blue #2 aluminum lake, hypromellose 2910, maize starch, microcrystalline cellulose, polyethylene glycol, povidone k-30, sodium starch glycolate, stearic acid, titanium dioxide.

Questions or comments?

1-877-770-3183 Mon-Fri 9:00 AM to 4:00 PM EST.



DIE GIFCOOI

 $1750 \times 1750 \times 3.375$ Fig. Note: Ensure that the unvannished area is not less than the minimum area specified on this document.





DIE GIFCOOI

1.750 × 1.750 × 3.375 Flat Note: Ensure that the unvamished area is not less than the minimum area specified on this document.

GRANULES		JOB INFORMATION	HISTORY			COLORS	
		ITEM#::300000023_CDC_Nap_Tab_50ct_FC_India_R2 CAD#::J50009A	Æ	Version¥ 2 Printed 9-10-16	Rav-6: Rav-7:	4COLOR PRO	Dieline UPC Code Measurement info. Costing Free
			& Ree-1:	Rev-1: 10-15-15		PN6 295 Pn Blue PNS 072	
				Reportio			
				Rev-4s	Plan-Bi	PMS 012	
						Q.C	
			Rev-it:	Rate-10:	0.6.		
	GRAI	WLES ARTWORK APPROVAL (THIS PROOF	OTEL	ST BE SIGNED A	WO RETURN	(EO) Sobwill not be released for production unles	s signed off by the below.
ррго чег		Nam e (print)				Signature	Date
OT C Marketing							
ackaging Engineering							
Regulatory Mairs							
Duality Assurance							
		YORKSON TO	41.0	LEASE PROOFREAD	and the second con-		

NAPROXEN SODIUM

naproxen sodium tablet

Product Information					
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68016-661		
Route of Administration	ORAL				

Active Ingredient/Active Moiety						
Ingredient Name	Basis of Strength	Strength				
NAPRO XEN SODIUM (UNII: 9TN87S3A3C) (NAPRO XEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg				

Inactive Ingredients					
Ingredient Name	Strength				
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)					
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)					
STARCH, CORN (UNII: O8232NY3SJ)					
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)					
PO VIDONE K30 (UNII: U725QWY32X)					
STEARIC ACID (UNII: 4ELV7Z65AP)					
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)					
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)					
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)					

Product Characteristics					
Color	blue	Score	no score		
Shape	ROUND	Size	10 mm		
Flavor		Imprint Code	220		
Contains					

]	Packaging						
7	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:68016-661-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	0 2/25/20 16				
2	NDC:68016-661-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	02/25/2016				

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
ANDA	ANDA091353	02/25/2016			

Labeler - Pharmacy Valuce Alliance, LLC (101668460)

Revised: 1/2021 Pharmacy Valuce Alliance, LLC