COLD AND COUGH- brompheniramine maleate, dextromethorphan hbr, phenylephrine hcl solution

HEB

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

HEB Cold & Cough Drug Facts

Active ingredients (in each 10 mL)

Brompheniramine maleate, USP 2 mg Dextromethorphan HBr, USP 10 mg Phenylephrine HCl, USP 5 mg

Purposes

Antihistamine

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- heart disease
- high blood pressure

- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers

When using this product

- do not use more than directed
- may cause marked drowsiness
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

age	dose	
adults and children 12 years and over	20 mL every 4 hours	
children 6 to under 12 years	10 mL every 4 hours	
children under 6 years	do not use	

Other information

- each 10 mL contains: sodium 4 mg
- store at 20°-25°C (68°-77°F)

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Children's Dimetapp® Cold & Cough active ingredients

children's Cold & Cough

Brompheniramine Maleate/Antihistamine

Dextromethorphan HBr/Cough Suppressant

Phenylephrine HCl/Nasal Decongestant

Multi – Symptom

For Ages 6 Yrs. & Over

Relief of:

Nasal Congestion

Runny Nose

Sneezing

Itchy, Watery Eyes

Coughing

Grape Flavor

Alcohol Free

4 FL OZ (118 mL)



Dextromethorphan HBr/

Cough Suppressant

Phenylephrine HCI/

Nasal Decongestant

Multi-Symptom

For Ages 6 Yrs. & Over

Nasal Congestion

Relief of:

Runny Nose

Grape Flavor

Alcohol Free

Sneezing

Compare to Children's Dimetapp® Cold & Cough active ingredients*

NDC 37808-987-26

(H-E-B)

Children's

Brompheniramine Maleate/ Antihistamine Dextromethorphan HBr/ Cough Suppressant Phenylephrine HCI/ Nasal Decongestant

Multi-Symptom

For Ages 6 Yrs. & Over Relief of:

- Nasal Congestion
- Runny Nose Sneezing
- Itchy, Watery Eyes
- Itchy, Watery EyesCoughing Coughing

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING



www.StopMedicineAbuse.org

Drug Facts

Active Ingredients (In each 10 mL)

Brompheniramine maleate, USP 2 mg.....Antihistamine Dextromethorphan HBr, USP 10 mg...Cough suppressant Phenylephrine HCl, USP 5 mg......Nasal decongestant

Phenylephrine HCI/ Nasal Decongestant an ppressant

An tih is tamine Brompheniramine Maleate/

8-3-H

Purposes

Dextromethorphan HBr/Cough

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever
- or other upper respiratory allergies

 temporarily relieves these symptoms due to hay
 fever (allergic rhinitis):

- sneezing
 itchy, watery eyes
 itching of the nose or throat
 temporarily restores freer breathing through the nose

Warnings

- Do not use

 if you are now taking a prescription monoamine
 oxidase inhibitor (MAOI) (certain drugs for
 depression, psychiatric, or emotional conditions, or
 Parkinson's disease), of for 2 weeks after stopping
 the MAOI drug, If you do not know if your
 prescription drug contains an MAOI, ask a doctor or
 pharmacist before taking this product.

 to make a child sleepy

Ask a doctor before use if you have

- heart disease high blood pressure
 thyroid disease diabetes
 trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
 cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- do not use more than directed may cause marked drowsiness avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase
- be careful when driving a motor vehicle or operating
- machinery

 excitability may occur, especially in children

Drug Facts (continued)

Stop use and ask a doctor it

- you get nervous, dizzy, or sleepless
 symptoms do not get better within 7 days or are
- accompanied by fever cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

serrous commons

If pregnant or breast-feeding, ask a health
professional before use.

Keep out of reach of children. In case of overdose,
get medical help or contact a Poison Control Center
right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
 measure only with dosage cup provided
 keep dosage cup with product
 mL = milliliter

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
shildren under Cueere	d 4

Other Information

■ each 10 mL contains: sodium 4 mg ■ store at 20°-25°C (68°-77°F)

Inactive Ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solutior

Questions or comments?

*This product is not manufactured or distributed by Pfizer. distributor of Children's Dimetapp® Cold & Cough

MADE WITH PRIDE & CARE FOR H-E-B[®] SAN ANTONIO, TX 78204





EXP.

98726 1J C6

COLD AND COUGH

brompheniramine maleate, dextromethorphan hbr, phenylephrine hcl solution

Product Information HUMAN OTC DRUG NDC:37808-987 Product Type Item Code (Source) ORAL Route of Administration

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BROMPHENIRAMINE MALEATE (UNII: IXA7C9ZN03) (BROMPHENIRAMINE - UNII:H57G17P2FN)	BROMPHENIRAMINE MALEATE	2 mg in 10 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 10 mL
PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 10 mL

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)		
EDETATE DISO DIUM (UNII: 7FLD9 1C8 6 K)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
GLYCERIN (UNII: PDC6 A3C0 O X)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SACCHARIN SO DIUM (UNII: SB8 ZUX40 TY)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SORBITOL (UNII: 506T60A25R)		

Product Characteristics			
Color	PURPLE (Clear bluish-red)	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

	Packaging				
1	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
	NDC:37808-987-26	1 in 1 CARTON	09/21/2006		
		118 mL in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/21/2006	

Labeler - HEB (007924756)

Revised: 4/2018 HE B