HIGHLIGHTS OF PRESCRIBING INFORMATION These highlights do not include all the information needed to use LOVAZA safely and effectively. See full prescribing information for LOVAZA.
LOVAZA® (omega-3-acid ethyl esters) Capsules Initial U.S. Approval: 2004
RECENT MAJOR CHANGES ·
Warnings and Precautions, Recurrent Atrial Fibrillation or Flutter 08/2012 (5.3)
INDICATIONS AND USAGE
LOVAZA is a combination of ethyl esters of omega 3 fatty acids, principally EPA and DHA, indicated as an adjunct to diet to reduce triglyceride (TG) levels in adult patients with severe (\geq 500 mg/dL) hypertriglyceridemia. (1) Limitations of Use: The effect of LOVAZA on cardiovascular mortality and morbidity in patients with elevated triglycerides has not been determined. (1)
• The daily dose of LOVAZA is 4 grams per day taken as a single 4-gram dose (4 capsules) or as two 2-gram doses (2
capsules given twice daily). (2)
 Patients should be advised to swallow LOVAZA capsules whole. Do not break open, crush, dissolve or chew LOVAZA. (2)
DOSAGE FORMS AND STRENGTHS
1-gram transparent soft-gelatin capsules. (3)
CONTRAINDICATIONS
LOVAZA is contraindicated in patients with known hypersensitivity (e.g., anaphylactic reaction) to LOVAZA or any of its components. (4)
WARNINGS AND PRECAUTIONS
 • In patients with hepatic impairment, monitor ALT and AST levels periodically during therapy. (5.1) • LOVAZA may increase levels of LDL. Monitor LDL levels periodically during therapy. (5.1) • Use with caution in patients with known hypersensitivity to fish and/or shellfish. (5.2)
• There is a possible association between LOVAZA and more frequent recurrences of symptomatic atrial fibrillation or flutter in patients with paroxysmal or persistent atrial fibrillation, particularly within the first months of initiating therapy. (5.3)
ADVERSE REACTIONS
The most common adverse reactions (incidence >3% and greater than placebo) were eructation, dyspepsia, and taste perversion. (6) (6)
To report SUSPECTED ADVERSE REACTIONS, contact Glaxo SmithKline at 1-888-825-5249 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch (6)
Omega-3-acids may prolong bleeding time. Patients taking LOVAZA and an anticoagulant or other drug affecting coagulation should be monitored periodically. (7.1)
• Pregnancy: Use during pregnancy only if the potential benefit justifies the potential risk to the fetus. (8.1)
• Pediatric Use: The safety and effectiveness in pediatric patients have not been established. (8.4)
See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 4/2013

LOVAZA- omega-3-acid ethyl esters capsule, liquid filled Cardinal Health

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1 INDICATIONS AND USAGE
- 2 DOSAGE AND ADMINISTRATION
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Monitoring: Laboratory Tests
- 5.2 Fish Allergy
- 5.3 Recurrent Atrial Fibrillation (AF) or Flutter

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

7.1 Anticoagulants or Other Drugs Affecting Coagulation

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

9 DRUG ABUSE AND DEPENDENCE

- 10 OVERDOSAGE
- 11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Severe Hypertriglyceridemia
- 14.2 Other Clinical Experience

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

- 17.1 Information for Patients
- 17.2 FDA-Approved Patient Labeling
- * Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

LOVAZA[®] (omega-3-acid ethyl esters) is indicated as an adjunct to diet to reduce triglyceride (TG) levels in adult patients with severe (\geq 500 mg/dL) hypertriglyceridemia.

Usage Considerations: Patients should be placed on an appropriate lipid-lowering diet before receiving LOVAZA and should continue this diet during treatment with LOVAZA.

Laboratory studies should be done to ascertain that the lipid levels are consistently abnormal before instituting LOVAZA therapy. Every attempt should be made to control serum lipids with appropriate diet, exercise, weight loss in obese patients, and control of any medical problems such as diabetes mellitus and hypothyroidism that are contributing to the lipid abnormalities. Medications known to exacerbate hypertriglyceridemia (such as beta blockers, thiazides, estrogens) should be discontinued or

changed if possible prior to consideration of triglyceride-lowering drug therapy.

Limitations of Use: The effect of LOVAZA on cardiovascular mortality and morbidity in patients with elevated triglycerides has not been determined.

2 DOSAGE AND ADMINISTRATION

- Assess triglyceride levels carefully before initiating therapy. Identify other causes (e.g., diabetes mellitus, hypothyroidism, or medications) of high triglyceride levels and manage as appropriate. [see Indications and Usage (1)].
- Patients should be placed on an appropriate lipid-lowering diet before receiving LOVAZA, and should continue this diet during treatment with LOVAZA. In clinical studies, LOVAZA was administered with meals.

The daily dose of LOVAZA is 4 grams per day. The daily dose may be taken as a single 4-gram dose (4 capsules) or as two 2-gram doses (2 capsules given twice daily).

Patients should be advised to swallow LOVAZA capsules whole. Do not break open, crush, dissolve or chew LOVAZA.

3 DOSAGE FORMS AND STRENGTHS

LOVAZA (omega-3-acid ethyl esters) capsules are supplied as 1-gram transparent soft-gelatin capsules filled with light-yellow oil and bearing the designation LOVAZA.

4 CONTRAINDICATIONS

LOVAZA is contraindicated in patients with known hypersensitivity (e.g., anaphylactic reaction) to LOVAZA or any of its components.

5 WARNINGS AND PRECAUTIONS

5.1 Monitoring: Laboratory Tests

In patients with hepatic impairment, alanine aminotransferase (ALT) and aspartate aminotransferase (AST) levels should be monitored periodically during therapy with LOVAZA. In some patients, increases in ALT levels without a concurrent increase in AST levels were observed.

In some patients, LOVAZA increases LDL-C levels. LDL-C levels should be monitored periodically during therapy with LOVAZA.

Laboratory studies should be performed periodically to measure the patient's TG levels during therapy with LOVAZA.

5.2 Fish Allergy

LOVAZA contains ethyl esters of omega-3 fatty acids (EPA and DHA) obtained from the oil of several fish sources. It is not known whether patients with allergies to fish and/or shellfish, are at increased risk of an allergic reaction to LOVAZA. LOVAZA should be used with caution in patients with known hypersensitivity to fish and/or shellfish.

5.3 Recurrent Atrial Fibrillation (AF) or Flutter

In a double-blind, placebo-controlled trial of 663 patients with symptomatic paroxysmal AF (n=542) or persistent AF (n=121), recurrent AF or flutter was observed in patients randomized to LOVAZA who received 8 grams/day for 7 days and 4 grams/day thereafter for 23 weeks at a higher rate relative to placebo. Patients in this trial had median baseline triglycerides of 127 mg/dL, had no substantial

structural heart disease, were taking no anti-arrhythmic therapy (rate control permitted), and were in normal sinus rhythm at baseline.

At 24 weeks, in the paroxysmal AF stratum, there were 129 (47%) first recurrent symptomatic AF or flutter events on placebo and 141 (53%) on LOVAZA [primary endpoint, HR 1.19; 95% CI 0.93, 1.35]. In the persistent AF stratum, there were 19 (35%) events on placebo and 34 (52%) events on LOVAZA [HR 1.63; 95% CI 0.91, 2.18]. For both strata combined, the HR was 1.25; 95% CI 1.00, 1.40. Although the clinical significance of these results is uncertain, there is a possible association between LOVAZA and more frequent recurrences of symptomatic atrial fibrillation or flutter in patients with paroxysmal or persistent atrial fibrillation, particularly within the first 2 to 3 months of initiating therapy.

LOVAZA is not indicated for the treatment of AF or flutter.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse reactions reported in at least 3% and at a greater rate than placebo for patients treated with LOVAZA based on pooled data across 23 clinical studies are listed in Table 1.

Table 1. Adverse Reactions Occurring at Incidence ≥3% and Greater than Placebo in Clinical Studies of LOVAZA

BODY	LOV	AZA	Placebo	
SYSTEM	(N =	655)	(N:	= 370)
Adverse	n	%	n	%
Reaction*				
Eructation	29	4	5	1
Dyspepsia	22	3	6	2
Taste	27	4	1	<1
perversion				

^{*} Studies included subjects with HTG and severe HTG.

Additional adverse reactions from clinical studies are listed below:

Digestive System: Constipation, gastrointestinal disorder and vomiting.

Metabolic and Nutritional Disorders: Increased ALT and increased AST.

Skin: Pruritus and rash.

6.2 Postmarketing Experience

In addition to adverse reactions reported from clinical trials, the events described below have been identified during post-approval use of LOVAZA. Because these events are reported voluntarily from a population of unknown size, it is not possible to reliably estimate their frequency or to always establish a causal relationship to drug exposure.

The following events have been reported: anaphylactic reaction, hemorrhagic diathesis.

7 DRUG INTERACTIONS

7.1 Anticoagulants or Other Drugs Affecting Coagulation

Some studies with omega-3-acids demonstrated prolongation of bleeding time. The prolongation of bleeding time reported in these studies has not exceeded normal limits and did not produce clinically significant bleeding episodes. Clinical studies have not been done to thoroughly examine the effect of LOVAZA and concomitant anticoagulants. Patients receiving treatment with LOVAZA and an anticoagulant or other drug affecting coagulation should be monitored periodically (e.g., aspirin, NSAIDS, warfarin, coumarin).

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C: There are no adequate and well-controlled studies in pregnant women. It is unknown whether LOVAZA can cause fetal harm when administered to a pregnant woman or can affect reproductive capacity. LOVAZA should be used during pregnancy only if the potential benefit to the patient justifies the potential risk to the fetus.

Animal Data

Omega-3-acid ethyl esters have been shown to have an embryocidal effect in pregnant rats when given in doses resulting in exposures 7 times the recommended human dose of 4 grams/day based on a body surface area comparison.

In female rats given oral gavage doses of 100, 600, and 2,000 mg/kg/day beginning 2 weeks prior to mating and continuing through gestation and lactation, no adverse effects were observed in the high dose group (5 times human systemic exposure following an oral dose of 4 grams/day based on body surface area comparison).

In pregnant rats given oral gavage doses of 1,000, 3,000, and 6,000 mg/kg/day from gestation day 6 through 15, no adverse effects were observed (14 times human systemic exposure following an oral dose of 4 grams/day based on a body surface area comparison).

In pregnant rats given oral gavage doses of 100, 600, and 2,000 mg/kg/day from gestation day 14 through lactation day 21, no adverse effects were seen at 2,000 mg/kg/day (5 times the human systemic exposure following an oral dose of 4 grams/day based on a body surface area comparison). However, decreased live births (20% reduction) and decreased survival to postnatal day 4 (40% reduction) were observed in a dose-ranging study using higher doses of 3,000 mg/kg/day (7 times the human systemic exposure following an oral dose of 4 grams/day based on a body surface area comparison).

In pregnant rabbits given oral gavage doses of 375, 750, and 1,500 mg/kg/day from gestation day 7 through 19, no findings were observed in the fetuses in groups given 375 mg/kg/day (2 times human systemic exposure following an oral dose of 4 grams/day based on a body surface area comparison). However, at higher doses, evidence of maternal toxicity was observed (4 times human systemic exposure following an oral dose of 4 grams/day based on a body surface area comparison).

8.3 Nursing Mothers

It is not known whether omega-3-acid ethyl esters are excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when LOVAZA is administered to a nursing woman.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

8.5 Geriatric Use

A limited number of patients older than 65 years were enrolled in the clinical studies of LOVAZA. Safety and efficacy findings in subjects older than 60 years did not appear to differ from those of subjects younger than 60 years.

9 DRUG ABUSE AND DEPENDENCE

LOVAZA does not have any known drug abuse or withdrawal effects.

10 OVERDOSAGE

In the event of an overdose, the patient should be treated symptomatically, and general supportive care measures instituted, as required.

11 DESCRIPTION

LOVAZA, a lipid-regulating agent, is supplied as a liquid-filled gel capsule for oral administration. Each 1-gram capsule of LOVAZA contains at least 900 mg of the ethyl esters of omega-3 fatty acids sourced from fish oils. These are predominantly a combination of ethyl esters of eicosapentaenoic acid (EPA - approximately 465 mg) and docosahexaenoic acid (DHA - approximately 375 mg).

The empirical formula of EPA ethyl ester is $C_{22}H_{34}O_2$, and the molecular weight of EPA ethyl ester is 330.51. The structural formula of EPA ethyl ester is:

The empirical formula of DHA ethyl ester is $C_{24}H_{36}O_2$, and the molecular weight of DHA ethyl ester is 356.55. The structural formula of DHA ethyl ester is:

LOVAZA capsules also contain the following inactive ingredients: 4 mg α -tocopherol (in a carrier of soybean oil), and gelatin, glycerol, and purified water (components of the capsule shell).

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism of action of LOVAZA is not completely understood. Potential mechanisms of action include inhibition of acyl-CoA:1,2-diacylglycerol acyltransferase, increased mitochondrial and peroxisomal β -oxidation in the liver, decreased lipogenesis in the liver, and increased plasma lipoprotein lipase activity. LOVAZA may reduce the synthesis of triglycerides in the liver because EPA and DHA are poor substrates for the enzymes responsible for TG synthesis, and EPA and DHA

inhibit esterification of other fatty acids.

12.3 Pharmacokinetics

In healthy volunteers and in patients with hypertriglyceridemia, EPA and DHA were absorbed when administered as ethyl esters orally. Omega-3-acids administered as ethyl esters (LOVAZA) induced significant, dose-dependent increases in serum phospholipid EPA content, though increases in DHA content were less marked and not dose-dependent when administered as ethyl esters.

Specific Populations

Age

Uptake of EPA and DHA into serum phospholipids in subjects treated with LOVAZA was independent of age (<49 years versus ≥49 years).

Gender

Females tended to have more uptake of EPA into serum phospholipids than males. The clinical significance of this is unknown.

Pediatric

Pharmacokinetics of LOVAZA in pediatric patients have not been established [see *Use in Specific Populations (8.4)*].

Renal or Hepatic Impairment

LOVAZA has not been studied in patients with renal or hepatic impairment.

Drug-Drug Interactions

Simvastatin

In a 14-day study of 24 healthy adult subjects, daily co-administration of simvastatin 80 mg with LOVAZA 4 grams did not affect the extent (AUC) or rate (C_{max}) of exposure to simvastatin or the major active metabolite, beta-hydroxy simvastatin at steady state.

Atorvastatin

In a 14-day study of 50 healthy adult subjects, daily co-administration of atorvastatin 80 mg with LOVAZA 4 grams did not affect AUC or C_{max} of exposure to atorvastatin, 2-hydroxyatorvastatin, or 4-hydroxyatorvastatin at steady state.

Rosuvastatin

In a 14-day study of 48 healthy adult subjects, daily co-administration of rosuvastatin 40 mg with LOVAZA 4 grams did not affect AUC or C_{max} of exposure to rosuvastatin at steady state.

In vitro studies using human liver microsomes indicated that clinically significant cytochrome P450 mediated inhibition by EPA/DHA combinations are not expected in humans.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In a rat carcinogenicity study with oral gavage doses of 100, 600, and 2,000 mg/kg/day, males were treated with omega-3-acid ethyl esters for 101 weeks and females for 89 weeks without an increased incidence of tumors (up to 5 times human systemic exposures following an oral dose of 4 grams/day based on a body surface area comparison). Standard lifetime carcinogenicity bioassays were not

conducted in mice.

Omega-3-acid ethyl esters were not mutagenic or clastogenic with or without metabolic activation in the bacterial mutagenesis (Ames) test with *Salmonella typhimurium* and *Escherichia coli* or in the chromosomal aberration assay in Chinese hamster V79 lung cells or human lymphocytes. Omega-3-acid ethyl esters were negative in the in vivo mouse micronucleus assay.

In a rat fertility study with oral gavage doses of 100, 600, and 2,000 mg/kg/day, males were treated for 10 weeks prior to mating and females were treated for 2 weeks prior to and throughout mating, gestation, and lactation. No adverse effect on fertility was observed at 2,000 mg/kg/day (5 times human systemic exposure following an oral dose of 4 grams/day based on a body surface area comparison).

14 CLINICAL STUDIES

14.1 Severe Hypertriglyceridemia

The effects of LOVAZA 4 grams per day were assessed in 2 randomized, placebo-controlled, double-blind, parallel-group studies of 84 adult patients (42 on LOVAZA, 42 on placebo) with very high triglyceride levels. Patients whose baseline triglyceride levels were between 500 and 2,000 mg/dL were enrolled in these 2 studies of 6 and 16 weeks duration. The median triglyceride and LDL-C levels in these patients were 792 mg/dL and 100 mg/dL, respectively. Median HDL-C level was 23.0 mg/dL.

The changes in the major lipoprotein lipid parameters for the groups receiving LOVAZA or placebo are shown in Table 2.

Table 2. Median Baseline and Percent Change From Baseline in Lipid Parameters in Patients with Very High TG Levels (≥500 mg/dL)

Parameter	LOVAZA N = 42		Placebo N = 42		Difference
	BL			%	
		Change		Change	
TG	816	-44.9	788	+6.7	-51.6
Non-HDL-	271	-13.8	292	-3.6	-10.2
C					
TC	296	-9.7	314	-1.7	-8.0
VLDL-C	175	-41.7	175	-0.9	-40.8
HDL-C	22	+9.1	24	0.0	+9.1
LDL-C	89	+44.5	108	-4.8	+49.3

BL = Baseline (mg/dL); % Change = Median Percent Change from Baseline; Difference = LOVAZA Median % Change – Placebo Median % Change

LOVAZA 4 grams per day reduced median TG, VLDL-C, and non-HDL-C levels and increased median HDL-C from baseline relative to placebo. Treatment with LOVAZA to reduce very high TG levels may result in elevations in LDL-C and non-HDL-C in some individuals. Patients should be monitored to ensure that the LDL-C level does not increase excessively.

The effect of LOVAZA on the risk of pancreatitis in patients with very high TG levels has not been evaluated.

The effect of LOVAZA on cardiovascular mortality and morbidity in patients with elevated TG levels has not been determined.

14.2 Other Clinical Experience

The effects of LOVAZA 4 grams per day as add-on therapy to treatment with simvastatin were evaluated in a randomized, placebo-controlled, double-blind, parallel-group study of 254 adult patients (122 on LOVAZA and 132 on placebo) with persistent high triglycerides (200 to 499 mg/dL) despite simvastatin therapy. Patients were treated with open-label simvastatin 40 mg per day for 8 weeks prior to randomization to control their LDL-C to no greater than 10% above NCEP ATP III goal and remained on this dose throughout the study. Following 8 weeks of open-label treatment with simvastatin, patients were randomized to either LOVAZA 4 grams per day or placebo for an additional 8 weeks with simvastatin co-therapy. The median baseline triglyceride and LDL-C levels in these patients were 268 mg/dL and 89 mg/dL, respectively. Median baseline non-HDL-C and HDL-C levels were 138 mg/dL and 45 mg/dL, respectively.

The changes in the major lipoprotein lipid parameters for the groups receiving LOVAZA plus simvastatin or placebo plus simvastatin are shown in Table 3.

Table 3. Response to the Addition of LOVAZA 4 grams per day to Ongoing Simvastatin 40 mg per day Therapy in Patients with High Triglycerides (200 to 499 mg/dL)

Parameter		_	ZA + statin 122	Placebo + Simvastatin N = 132		Difference	P-Value	
	BL	EOT	Median %	BL	EOT	Median %		
			Change			Change		
Non-HDL-	137	123	-9.0	141	134	-2.2	-6.8	< 0.0001
C								
TG	268	182	-29.5	271	260	-6.3	-23.2	< 0.0001
TC	184	172	-4.8	184	178	-1.7	-3.1	< 0.05
VLDL-C	52	37	-27.5	52	49	-7.2	-20.3	< 0.05
Apo-B	86	80	-4.2	87	85	-1.9	-2.3	< 0.05
HDL-C	46	48	+3.4	43	44	-1.2	+4.6	< 0.05
LDL-C	91	88	+0.7	88	85	-2.8	+3.5	=0.05

BL = Baseline (mg/dL); EOT = End of Treatment (mg/dL); Median % Change = Median Percent Change from Baseline; Difference = LOVAZA Median % Change – Placebo Median % Change

LOVAZA 4 grams per day significantly reduced non-HDL-C, TG, TC, VLDL-C, and Apo-B levels and increased HDL-C and LDL-C from baseline relative to placebo.

16 HOW SUPPLIED/STORAGE AND HANDLING

LOVAZA (omega-3-acid ethyl esters) capsules are supplied as 1-gram transparent soft-gelatin capsules filled with light-yellow oil and bearing the designation LOVAZA.

Bottles of 120: NDC 0173-0783-02

Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Do not freeze. Keep out of reach of children.

17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (17.2).

17.1 Information for Patients

- LOVAZA should be used with caution in patients with known sensitivity or allergy to fish and/or shellfish [see Warnings and Precautions (5.2)].
- Patients should be advised that use of lipid-regulating agents does not reduce the importance of adhering to diet [see Dosage and Administration (2)].
- Patients should be advised not to alter LOVAZA capsules in any way and to ingest intact capsules only [see Dosage and Administration (2)].

17.2 FDA-Approved Patient Labeling

Patient labeling is provided as a tear-off leaflet at the end of this full prescribing information.

Manufactured for GlaxoSmithKline by:

Catalent Pharma Solutions

St. Petersburg, FL 33716

Distributed by:

GlaxoSmithKline

Research Triangle Park, NC 27709

LOVAZA is a registered trademark of the GlaxoSmithKline group of companies.

©2012, GlaxoSmithKline. All rights reserved.

LVZ:8PI

PHARMACIST-DETACH HERE AND GIVE INSTRUCTIONS TO PATIENT

PATIENT INFORMATION

LOVAZA® (lō-vā-ză)

(omega-3-acid ethyl esters)

Capsules

Read this Patient Information before you start taking LOVAZA, and each time you get a refill. There may be new information. This information does not take the place of talking with your doctor about your medical condition or your treatment.

What is LOVAZA?

LOVAZA is a prescription medicine used along with a low fat and low cholesterol diet to lower very high triglyceride (fat) levels in adults.

It is not known if LOVAZA prevents you from having a heart attack or stroke.

It is not known if LOVAZA is safe and effective in children.

Who should not take LOVAZA?

Do not take LOVAZA if you are allergic to omega-3-acid ethyl esters or any of the ingredients in LOVAZA. See the end of this leaflet for a complete list of ingredients in LOVAZA.

What should I tell my doctor before taking LOVAZA?

Before you take LOVAZA, tell your doctor if you:

- •have diabetes.
- •have a low thyroid problem (hypothyroidism).
- •have a liver problem.
- •have a pancreas problem.
- •have a certain heart rhythm problem called atrial fibrillation or flutter.
- •are allergic to fish or shellfish. It is not known if people who are allergic to fish or shellfish are also allergic to LOVAZA.
- • are pregnant or plan to become pregnant. It is not known if LOVAZA will harm your unborn baby.
- •are breastfeeding or plan to breastfeed. It is not known if LOVAZA passes into your breast milk. You and your doctor should decide if you will take LOVAZA or breastfeed.

Tell your doctor about all the medicines you take, including prescription and non-prescription medicine, vitamins, and herbal supplements.

LOVAZA can interact with certain other medicines that you are taking. Using LOVAZA with medicines that affect blood clotting (anticoagulants or blood thinners) may cause serious side effects.

Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

How should I take LOVAZA?

- • Take LOVAZA exactly as your doctor tells you to take it.
- You should not take more than 4 capsules of LOVAZA each day. Either take all 4 capsules at one time, or 2 capsules two times a day.
- •Do not change your dose or stop LOVAZA without talking to your doctor.
- Take LOVAZA with or without food.
- Take LOVAZA capsules whole. Do not break, crush, dissolve, or chew LOVAZA capsules before swallowing. If you cannot swallow LOVAZA capsules whole, tell your doctor. You may need a different medicine.
- Your doctor should start you on a low fat and low cholesterol diet before giving you LOVAZA. Stay on a low fat and low cholesterol diet while you take LOVAZA.
- Your doctor should do blood tests to check your triglyceride, bad cholesterol and liver function levels while you take LOVAZA.

What are the possible side effects of LOVAZA?

LOVAZA may cause serious side effects, including:

- •increases in the results of blood tests used to check your liver function (ALT and AST) and your bad cholesterol levels (LDL-C) cholesterol.
- •increases in the frequency of a heart rhythm problem (atrial fibrillation or flutter) may especially happen in the first few months of taking LOVAZA if you already have that problem.

The most common side effects of LOVAZA include:

- burping
- •upset stomach
- a change in your sense of taste.

Talk to your doctor if you have a side effect that bothers you or does not go away.

These are not all the possible side effects of LOVAZA. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store LOVAZA?

- •Store LOVAZA at room temperature between 68°F to 77°F (20°C to 25°C).
- •Do not freeze LOVAZA.
- •Safely throw away medicine that is out of date or no longer needed.

Keep LOVAZA and all medicines out of the reach of children.

General information about the safe and effective use of LOVAZA

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use LOVAZA for a condition for which it was not prescribed. Do not give LOVAZA to other people, even if they have the same symptoms you have. It may harm them.

This Patient Information Leaflet summarizes the most important information about LOVAZA. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about LOVAZA that is written for health professionals.

For more information go to www.LOVAZA.com or call 1-888-825-5249.

What are the ingredients in LOVAZA?

Active Ingredient: omega-3-acid ethyl esters, mostly EPA and DHA

Inactive Ingredients: alpha-tocopherol (in soybean oil), gelatin, glycerol, purified water

This patient labeling has been approved by the U.S. Food and Drug Administration.

Manufactured for GlaxoSmithKline by:

Catalent Pharma Solutions

St. Petersburg, FL 33716

Distributed by:

GlaxoSmithKline

Research Triangle Park, NC 27709

©2012, GlaxoSmithKline. All rights reserved.

August 2012

LVZ:6PIL

LVZ:4PIL

LOVAZA (omega-3-acid ethyl esters) is available from Cardinal Health in unit dose packages of 120 capsules.

LOVAZA, unit dose package of 120 capsules, NDC 55154-1134-4

Cardinal Health

IU42458660312

PRINCIPAL DISPLAY PANEL - Carton

NDC 55154-1134-4

Lovaza®

(omega-3-acid ethyl esters)

Capsules

1 Gram

120 Capsules

Each capsule contains 1 gram omega-3-acid ethyl ester liquid concentrate consisting of at least 900 mg omega-3-acid ethyl esters. SWALLOW CAPSULES WHOLE.

Each capsule provides: Eicosapentaenoic acid (EPA) ethyl ester: 465 mg Docosahexaenoic acid (DHA) ethyl ester: 375 mg

Usual Dosage: See product insert for prescribing information, precautions and warnings.

STORAGE: Store at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F) [see USP Controlled Room Temperature].

Do not freeze.

RX ONLY

WARNING: This package is intended for in-patient institutional use only. This package is not child resistant. Keep this and all drugs out of the reach of children.

See window for lot number and expiration date.

Mfd for: GlaxoSmithKline, RTP, NC 27709

Mfd by: Catalent Pharma Solutions, St. Petersburg, FL 33716 or

Made in Norway

Repackaged by: Cardinal Health Zanesville, OH 43701

LUC424586600712

NDC 55154-1134-4

Lovaza®

(omega-3-acid ethyl esters) Capsules

1 Gram

120 Capsules



(01)00355154113442

NDC 55154-1134-4 Lovaza® (omega-3-acid ethyl esters) Capsules 1 Gram 120 Capsules

Each capsule contains 1 gram omega-3-acid ethyl ester liquid concentrate consisting of at least 900 mg omega-3-acid ethyl esters. SWALLOW CAPSULES WHOLE.

Each capsule provides: Eicosapentaenoic acid (EPA) ethyl ester: 465 mg Docosahexaenoic acid (DHA) ethyl ester: 375 mg

Usual Dosage: See product insert for prescribing information, precautions and warnings.

STORAGE: Store at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F) [see USP Controlled Room Temperature]. Do not freeze.

RX ONLY

WARNING: This package is intended for in-patient institutional use only. This package is not child resistant. Keep this and all drugs out of the reach of children.

See window for lot number and expiration date.

Mfd for: GlaxoSmithKline, RTP, NC 27709

Mfd by: Catalent Pharma Solutions, St. Petersburg, FL 33716

Made in Norway

CardinalHealth

Repackaged by: Cardinal Health

Zanesville, OH 43701

LUC42458660712

Principal Display Panel - Pouch

Lovaza®

(omega-3-acid ethyl esters)

Capsule

1 Gram



PRINCIPAL DISPLAY PANEL - Bag

Lovaza

Omega-3-Acid Ethyl Esters Capsules

10 Capsules



168

LOVAZA

Omega-3-Acid Ethyl Esters Capsules

10 CAPSULES

Swallow capsules whole.

Each capsule contains 1 gram omega-3-acid ethyl ester liquid concentrate consisting of at least 900 mg omega-3-acid ethyl esters.

Each capsule provides: Eicosapentaenoic acid (EPA) ethyl ester: 465 mg Docosahexaenoic acid (DHA) ethyl ester: 375 mg

Usual Dosage: See product insert for prescribing information, precautions and warnings.

STORAGE: Store at 25 C (77 F); excursions permitted to 15 - 30 C (59 - 86 F) [see USP Controlled Room Temperature]. Do not freeze.

RX ONLY

WARNING: This package is intended for in-patient institutional use only. This package is not child resistant. Keep this and all drugs out of the reach of children.

Mfd for:

GlaxoSmithKline, RTP, NC 27709

Mfd by:

Catalent Pharma Solutions St. Petersburg, FL 33716

Made in Norway

LOVAZA is a trademark owned by its respective owner.

Repackaged by Cardinal Health Zanesville, OH 43701 L42458660712

LOT#: XXXX

EXP. DATE: 12/34



LOVAZA

omega-3-acid ethyl esters capsule, liquid filled

Product Information

Active Ingredient/Active Moiety

WATER

Ingredient Name	Basis of Strength	Strength
OMEGA-3-ACID ETHYL ESTERS (OMEGA-3 FATTY ACIDS)	OMEGA-3-ACID ETHYL ESTERS	1 g

Inactive Ingredients				
Ingredient Name	Strength			
.ALPHATO CO PHERO L				
GELATIN				
GLYCERIN				

Product Characteristics					
Color	YELLOW (light yellow)	Score	no score		
Shape	CAPSULE	Size	24mm		
Flavor		Imprint Code	LOVAZA		
Contains					

P	Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:55154-1134-4	120 in 1 CARTON					
1	NDC:55154-1134-6	1 in 1 POUCH					
2	NDC:55154-1134-0	10 in 1 BAG					
2		1 in 1 POUCH					

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA	NDA021654	10/21/2008		

Labeler - Cardinal Health (188557102)

Establishment			
Name	Address	ID/FEI	Business Operations
Cardinal Health		188557102	REPACK(55154-1134)

Revised: 4/2013 Cardinal Health