

GOOD NEIGHBOR PAIN RELIEF- benzocaine liquid
AmerisourceBergen Drug Corporation (Good Neighbor Pharmacy)

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient

Benzocaine, 20%

□ Purpose

Oral anesthetic

Uses

Provides temporary relief of pain associated with the following mouth and gum irritations:

- toothache
- sore gums
- canker sores
- braces
- minor dental procedures
- dentures

□ Warnings

□ Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

□ When using this product

- avoid contact with eyes
- do not exceed recommended dosage
- do not use for more than 7 days unless directed by a doctor or dentist

□ Stop use and ask a doctor or dentist if

- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling, rash, or fever develops

□ Keep out of reach of children

- If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away

□ Directions

- ***□ Adults and children 2 years of age and older:***
- wipe liquid on with cotton, or cotton swab, or finger tip.
- apply to the affected area up to 4 times daily of as directed by a doctor or dentist.

- **Children under 12 years of age:** adult supervision should be given in the use of this product.
- **Children under 2 years of age:** consult a doctor or dentist.

Inactive ingredients

Benzyl alcohol, FD&C Blue #1, FD&C Red #40, D&C yellow #10, Flavor, Methylparaben, Polyethylene Glycol, Propylene Glycol, Saccharin

7.6X3.8CM

NDC 46122-030-05



Active ingredient
Benzocaine 20%

Purpose
Oral anesthetic

Uses Provides temporary relief of pain associated with the following mouth and gum irritations: toothache, sore gums, canker sores, braces, minor dental procedures, dentures

Warnings Allergy alert: Do not use this product if you have a history of allergy of local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics

When using this product : avoid contact with the eyes; do not exceed recommended dosage; do not use for more than 7 days unless directed by a doctor/dentist

Stop use and ask a doctor if: sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; swelling, rash, or fever develops

Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

Directions Adults and children 2 years of age and older: wipe liquid on with cotton, cotton swab, or fingertip. Apply to the affected area up to 4 times daily or as directed by a doctor/dentist. **Children under 12 years of age:** adult supervision should be given in the use of this product. **Children under 2 years of age:** consult a doctor/dentist

Other information • store at room temperature

SEE BLISTER CARD FOR COMPLETE LABELING INFO.

Distributed By: AmerisourceBergen
1300 Morris Drive Chesterbrook, PA 19087
Visit us at www.goodneighborpharmacy.com
160386A, Rev. 3/12

LOT: _____

EXP: _____



Manufacturer: Lee Pharmaceuticals



7.6x14.9cm

* Compare to Anbesol®
active ingredient
NDC 46122-030-05



Oral Pain Relief

Liquid Oral Anesthetic

benzocaine 20%
Instant Pain Relief for:

- Toothache
- Canker Sores
- Denture Pain
- Gum Pain

0.5 FL OZ (15 mL)



Oral Pain Relief

Liquid Oral Anesthetic

Do not use if plastic blister or backing material is broken or separated

Drug Facts	
Active ingredient Benzocaine, 20%	Purpose Oral anesthetic
Uses Provides temporary relief of pain associated with the following mouth and gum irritations: • toothache • sore gums • canker sores • braces • minor dental procedures • dentures	
Warnings Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other 'caine' anesthetics.	
When using this product • avoid contact with eyes • do not exceed recommended dosage • do not use for more than 7 days unless directed by a doctor or dentist	
Stop use and ask a doctor or dentist if • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops	
Keep out of the reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.	
Directions • Adult and children 2 years of age and older: • wipe liquid on with cotton, or cotton swab, or fingertip • apply to the affected area up to 4 times daily or as directed by a doctor or dentist • Children under 12 years of age: adult supervision should be given in the use of this product • Children under 2 years of age: consult a doctor or dentist	
Other information • Store at room temperature	
Inactive ingredients benzyl alcohol, FD&C Blue #1, FD&C Red #40, FD&C Yellow #10, flavor, methylparaben, polyethylene glycol, propylene glycol, saccharin	

*This product is not manufactured or distributed by Wyeth Consumer Healthcare, owner of the registered trademark Anbesol®

Distributed By
AmerisourceBergen
1300 Morris Drive
Chesterbrook, PA 19087
Visit us at
www.goodneighborpharmacy.com
Questions or comments:
800-950-5337
Cat. No. 0386-000
180386A, Rev. 3/12



ABC# 070-932



0 187701 40627 0

9 font HELVETICA Condensed	8 font HELVETICA Condensed	6 font HELVETICA Condensed
9 font HELVETICA Condensed	8 font HELVETICA Condensed	6 font HELVETICA Condensed

Manufacturer: Lee Pharmaceuticals



GOOD NEIGHBOR PAIN RELIEF

benzocaine liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-030
---------------------	----------------	---------------------------	---------------

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN (UNII: FST467XS7D)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-030-05	1 in 1 BLISTER PACK		
1	NDC:46122-030-06	15 mL in 1 BOTTLE, GLASS		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	08/01/2008	

Labeler - AmerisourceBergen Drug Corporation (Good Neighbor Pharmacy) (007914906)

Registrant - Lee Pharmaceuticals (056425432)

Establishment

Name	Address	ID/FEI	Business Operations
AmerisourceBergen Drug Corporation		007914906	label(46122-030)

Establishment

Name	Address	ID/FEI	Business Operations
Lee Pharmaceuticals		056425432	pack(46122-030)

Establishment

Name	Address	ID/FEI	Business Operations
Unipack		009248480	manufacture(46122-030)

Establishment

Name	Address	ID/FEI	Business Operations
Samson Pharmaceuticals		088169581	manufacture(46122-030)

Revised: 6/2013

AmerisourceBergen Drug Corporation (Good Neighbor Pharmacy)