

PAIN RELIEVER REGULAR STRENGTH- acetaminophen tablet
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/ fever reducer

Uses

- temporarily relieves minor aches and pains due to
 - headache
 - the common cold
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has

liver disease.

Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 tablets every 4 to 6 hours while symptoms last• do not take more than 10 tablets in 24 hours• do not use for more than 10 days unless directed by a doctor
children 6-11 years	<ul style="list-style-type: none">• take 1 tablet every 4 to 6 hours while symptoms last• do not take more than 5 tablets in 24 hours• do not use for more than 5 days unless directed by a doctor

children under 6 years	• ask a doctor
------------------------	----------------

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients

povidone, pregelatinized starch, sodium starch glycolate*, stearic acid

*may contain this ingredient

Questions or comments?

Call **1-888-309-9030**

Principal Display Panel

Compare to the active ingredient of Tylenol® Regular Strength**

Regular Strength

Pain Reliever

Acetaminophen 325 mg

Pain Reliever/ Fever Reducer

For ages 6 years and over

Tablets

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Regular Strtength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY DOLGENCORP, LLC

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

Product Label

Exp. Date:
Lot No.:
PLD-B205B
FC006153



A0907

100%
Satisfaction
Guaranteed!
(888) 309-9030

DISTRIBUTED BY DOLGENCORP, LLC
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072

Drug Facts

Active ingredient
(in each tablet)

Acetaminophen 325 mg.....Pain reliever/
fever reducer

Uses

- temporarily relieves minor aches and pains due to
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include

- skin redness
 - skin redness
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Drug Facts (continued)

Ask a doctor before use if the user has liver disease.

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
 - pain gets worse or lasts more than 5 days in children under 12 years
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Drug Facts (continued)

Directions

do not take more than directed (see Overdose warning)

adults and children 12 years and over

- take 2 tablets every 4 to 6 hours while symptoms last
- do not take more than 10 tablets in 24 hours
- do not use for more than 10 days unless directed by a doctor

children 6-11 years

- take 1 tablet every 4 to 6 hours while symptoms last
- do not take more than 5 tablets in 24 hours
- do not use for more than 5 days unless directed by a doctor

children under 6 years

- ask a doctor

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients

povidone, pregelatinized starch, sodium starch glycolate*, stearic acid

*may contain this ingredient

Questions or comments?

Call 1-888-309-9030

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Regular Strength.

DG™ health

Regular Strength Pain Relief

Acetaminophen 325 mg

Pain Reliever/Fever Reducer

325
mg

For ages 6 years and over

Compare to the active ingredient of Tylenol® Regular Strength**

100 Tablets

Actual Tablet Size

DOLLAR GENERAL Regular Strength Pain Relief

PAIN RELIEVER REGULAR STRENGTH

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-691
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	TCL340
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-691-10	1 in 1 BOX	04/30/2018	04/30/2025
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/30/2018	04/30/2025

Labeler - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

