ACID CONTROLLER- famotidine tablet, film coated
Walgreen Company

Walgreen Co. Acid Controller Drug Facts

Active ingredient (in each tablet)

Famotidine 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Walgreens

WALGREENS PHARMACIST RECOMMNEDED

Compare to the active ingredient in Maximum Strength Pepcid® AC

Acid Controller

FAMOTIDINE TABLETS, 20 mg / ACID REDUCER

Maximum Strength

• Just one tablet prevents & relieves heartburn due to acid indigestion

85 TABLETS

ACTUAL SIZE

SEE NEW WARNINGS

ACTUAL SIZE

W3ORG1021-F2

NDC 0363-0701-01

Walgreens

Compare to the active ingredient in Maximum Strength Pepcid® AC*

WALGREENS PHARMACIST RECOMMENDED

Acid Controller

FAMOTIDINE TABLETS, 20 mg / ACID REDUCER

Maximum Strength

- Just one tablet prevents & relieves heartburn due to acid indigestion

85 TABLETS

ACTUAL SIZE

SEE NEW WARNINGS

1946 94 CS

CODE AREA

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.

ITEM 551851 WH0106-02224-L

3 11917 08767 2

Gluten Free

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet) — **Purpose**

Famotidine 20 mg ..... Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition. ▶

Drug Facts (continued)

- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) ▼

Drug Facts (continued)

Directions

- adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
  - to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes

Drug Facts (continued)

- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol

†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

††This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Maximum Strength Pepcid® AC.

PAPER BOX | PLASTIC BOTTLE



ACID CONTROLLER

famotidine tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:0363-0701
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients	
Ingredient Name	**Strength**
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	WHITE	**Score**	no score
Shape	ROUND	**Size**	8mm
Flavor		**Imprint Code**	L194
Contains			

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:0363-0701-71	1 in 1 CARTON	09/26/2006	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		

2	NDC:0363-0701-72	1 in 1 CARTON	06/02/2011	09/30/2020
2		60 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0363-0701-01	1 in 1 CARTON	10/29/2007	
3		85 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0363-0701-02	25 in 1 CARTON	09/26/2006	
4		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:0363-0701-39	30 in 1 CARTON	04/10/2015	04/10/2015
5		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
6	NDC:0363-0701-82	1 in 1 CARTON	05/08/2019	
6		200 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077351	09/26/2006	

Labeler - Walgreen Company (008965063)

Revised: 5/2022 Walgreen Company