

DG HEALTH NAPROXEN SODIUM- naproxen sodium tablet, film coated
Dolgencorp, LLC

Dolgencorp, LLC Naproxen Sodium Tablets, 220 mg Drug Facts

Active ingredient (in each tablet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none"> • take 1 tablet every 8 to 12 hours while symptoms last • for the first dose you make take 2 tablets within the first hour • do not exceed 2 tablets in any 8- to 12-hour period • do not exceed 3 tablets in a 24-hour period
Children under 12 years	<ul style="list-style-type: none"> • ask a doctor

Other information

- **each tablet contains:** sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?

1-888-309-9030

Package/Label Principal Display Panel

Compare to the active ingredient of Aleve® Headache Pain

Naproxen Sodium Tablets, 220 mg

Pain Reliever/Fever Reducer (NSAID)

For temporary relief of headache pain

STRENGTH TO LAST 12 HOURS

HEADACHE PAIN

24 Tablets

220 mg

Actual Tablet Size

Actual Size



DO NOT USE IF PRINTED FOIL UNDER CAP
IS BROKEN OR MISSING

KEEP CARTON FOR REFERENCE

**This product is not manufactured or distributed by Bayer
HealthCare LLC, distributor of Aleve® Headache Pain.

DG™ | health

Compare
to the
active
ingredient
of Aleve®
Headache
Pain**

Naproxen Sodium

Tablets, 220 mg

**Pain Reliever/
Fever Reducer (NSAID)**

For temporary relief of
headache pain



HEADACHE PAIN




24 Tablets

**220
mg**

Actual Tablet Size





48C62 VT C1



Drug Facts

Active ingredient (in each tablet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/

fever reducer

Uses ■ temporarily relieves minor aches and pains due to:

■ headache ■ minor pain of arthritis ■ muscular aches

■ backache ■ menstrual cramps ■ toothache

■ the common cold ■ temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin ■ taking any other drug

When using this product

■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ you have difficulty swallowing ■ it feels like the pill is stuck in your throat ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

■ do not take more than directed

■ the smallest effective dose should be used

■ drink a full glass of water with each dose

Adults and children

12 years and older

■ take 1 tablet every 8 to 12 hours

while symptoms last

■ for the first dose you may take

2 tablets within the first hour

■ do not exceed 2 tablets in any

8- to 12-hour period

■ do not exceed 3 tablets in a

24-hour period

Children under 12 years


■ ask a doctor

DISTRIBUTE
OLDEST M
100 MISSION
GOODLETT
TN37072

Questions or

Inactive ingre
hy promellose, m
cellulose & polyeth

Drug Facts (co
Other in form:
sodium 21 mg
humidity and exc



DG HEALTH NAPROXEN SODIUM

naproxen sodium tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-481
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name		Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)		NAPROXEN SODIUM 220 mg

Inactive Ingredients	
Ingredient Name	Strength
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	BLUE	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	L490
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date

1	NDC:55910-481-62	1 in 1 CARTON	05/26/2021	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55910-481-71	1 in 1 CARTON	05/26/2021	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	05/26/2021	

Labeler - Dolgencorp, LLC (068331990)

Revised: 5/2021

Dolgencorp, LLC