TENSION HEADACHE ASPIRIN FREE- acetaminophen, caffeine tablet, film coated CVS Pharmacy

CVS 44-428

Active ingredients (in each caplet)

Acetaminophen 500 mg (formulated with 65 mg caffeine)

Purpose

Pain reliever

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen

Ask a doctor before use if you have

liver disease.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- new symptoms occur
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- adults and children 12 years and over: take 2 caplets every 6 hours. Do not take more than 6 caplets in 24 hours.
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, crospovidone, D&C red #27 aluminum lake, FD&C blue #2 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

♥CVS Health™

Compare to the active ingredients in Excedrin® Tension Headache*

Coated Caplets

Aspirin Free Tension Headache

ACETAMINOPHEN & CAFFEINE Pain reliever

100 CAPLETS

Actual Size


TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*This product is not manufactured or distributed by GSK Consumer Healthcare S.A., owner of the registered trademark Excedrin® Tension Headache.

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895 © 2020 CVS/pharmacy CVS.com® 1-800-SHOP CVS V-11112

50844 REV1119A42812



Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe If a skin reaction occurs, stop use and seek ■ blisters ■ rash ■ skin reddening skin reactions. Symptoms may include medical help right away.

Caffeine warning: The recommended dose of this medications, foods, or beverages while taking this product contains about as much caffeine as a cup product because too much caffeine may cause of coffee. Limit the use of caffeine-containing nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat

Do not use

(prescription or nonprescription). If you are not with any other drug containing acetaminophen sure whether a drug contains acetaminophen ask a doctor or pharmacist

Ask a doctor before use if you have liver disease ■ if you are allergic to acetaminophen

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

■ painful area is red or swollen Stop use and ask a doctor if ■ new symptoms occur

■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days These could be signs of a serious condition. If pregnant or breast-feeding, ask a health

accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical children even if you do not notice any signs or attention is critical for adults as well as for professional before use.

Keep out of reach of children. In case of symptoms

CVS 44-428

FD&C blue #2 aluminum lake, FD&C yellow #6 microcrystalline cellulose, polyethylene glycol

aluminum lake, magnesium stearate,

alycolate, stearic acid, talc, titanium dioxide polyvinyl alcohol, povidone, sodium starch

Questions or comments?

1-800-426-939

crospovidone, D&C red #27 aluminum lake,

Inactive ingredients corn starch

store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

use by expiration date on package

TENSION HEADACHE ASPIRIN FREE

adults and children 12 years and over: take

do not take more than directed

Directions

STOP PEELING

Drug Facts (continued)

2 caplets every 6 hours. Do not take more

children under 12 years: ask a doctor

Other information

than 6 caplets in 24 hours.

acetaminophen, caffeine tablet, film coated

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59779-824
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	500 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients	
Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPOVIDONE (UNII: 2S7830E561)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ 989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	red	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	44;428
Contains			

P	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779- 824-21	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/17/2007	
2	NDC:59779- 824-06	200 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/17/2007	
3	NDC:59779- 824-08	1 in 1 CARTON	01/17/2007	12/01/2024
3		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:59779- 824-12	1 in 1 CARTON	01/17/2007	02/11/2018
4		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M013	01/17/2007			

Labeler - CVS Pharmacy (062312574)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(59779-824)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(59779-824)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	pack(59779-824)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(59779-824)

Revised: 7/2023 CVS Pharmacy