HEARTBURN RELIEF- cimetidine tablet, film coated CVS Pharmacy

CVS Pharmacy, Inc. Heartburn Relief Drug Facts

Active ingredient (in each tablet)

Cimetidine 200 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- **prevents** heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

Ask a doctor or pharmacist before use if you are taking

a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- you need to take this product for more than 14 days
- stomach pain continues
- your heartburn continues or worsens

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water
- to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30minutes before eating food or drinking beverages that cause heartburn
- do not take more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

store at 20°-25°C (68°-77°F)

Inactive ingredients

hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

CVS Health®

Compare to the active ingredient in Tagamet $^{\circledR}$ HB 200

Heartburn Relief

CIMETIDINE TABLETS USP, 200 mg

Acid reducer

Just one tablet:

Relieves and prevents heartburn and acid indigestion

Take ANY TIME you need it:


- Before meal
- During meal
- After meal

Actual Size

60 TABLETS (60 Doses)

Actual Bottle Size on Side Panel





Drug Facts

Active ingredient (in each tablet)

drinking certain foods and beverages

Purpose .. Acid reducer

Cimetidine 200 mg......

USES ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents hear burn associated with acid indiges fron and sour stomach brought on by eating or

Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers

- if you have trouble or pain swall owing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Aska doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

Ask a doctor or pharmacist before use if you are taking

- a prescription drug. Acid reducers may interact with certain prescription drugs.
- Stop use and ask a doctor if
- you need to take this product for more than 14 days
- your heartburn continues or worsens ■ stomach pain continues

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults and children 12 years and over:
- to relieve symptoms, swallow 1 tablet with a glass of water
- to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn
- do not take more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Drug Facts (continued)

Other information

■ store at 20°-25°C (68°-77°F)

Inactive ingredients hypromellose, magnesium stearate, microcrystalline cellulose, poly ethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions or comments? 1-800-719-9260


- Tips for Managing Heartburn
 Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- · Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
 If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Do not use if printed foil under cap is broken or missing.

Distributed by: CVS Pharmacy, Inc. One CVS Drive, Woonsocket, RI 02895 @ 2021 CVS/pharmacy CVS.come 1-800-SHOPCVS

V-16467







Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:59779-022

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CIMETIDINE (UNII: 80061L1WGD) (CIMETIDINE - UNII:80061L1WGD)	CIMETIDINE	200 mg

Inactive Ingredients	
Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 60ZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ 989GH94E)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics					
Color	WHITE	Score	no score		
Shape	OVAL	Size	13mm		
Flavor		Imprint Code	L022		
Contains					

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:59779-022- 72	1 in 1 CARTON	08/07/2009			
1		60 in 1 BOTTLE; Type 0: Not a Combination Product				
2	NDC:59779-022- 76	1 in 1 CARTON	01/05/2022			
2		120 in 1 BOTTLE; Type 0: Not a Combination Product				

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075285	08/07/2009	

Labeler - CVS Pharmacy (062312574)

Revised: 1/2022 CVS Pharmacy