

**DG HEALTH ANTI DIARRHEAL- loperamide hcl solution**  
**Dolgencorp Inc**

-----  
**Dolgencorp, LLC Anti-Diarrheal Drug Facts**

**Active ingredient (in each 7.5 mL)**

Loperamide HCl 1 mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
children 9-11 years (60-95 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

- **each 30 mL contains:** sodium 15 mg
- store between 20-25°C (68-77°F)

**Inactive ingredients**

anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

**Questions or comments?**

**1-888-309-9030**

**Package/Label Principal Display Panel**

SEE NEW WARNING AND DIRECTIONS

Compare to the active ingredient of Imodium® A-D

Loperamide Hydrochloride Oral Solution 1 mg per 7.5 mL

Anti-Diarrheal

Controls the Symptoms of Diarrhea

Anti-Diarrheal Oral Solution

Mint Flavor

4 FL OZ (118mL)

## Drug Facts

**Active ingredient**      **Purpose**  
(in each 7.5 mL)  
Loperamide HCl 1 mg....Anti-diarrheal

**Use** controls symptoms of diarrhea,  
including Travelers' Diarrhea

### Warnings

**Allergy alert:** Do not use if you  
have ever had a rash or other  
allergic reaction to loperamide HCl  
**Heart alert:** Taking more than  
directed can cause serious heart  
problems or death

Peel open to read complete directions  
and warnings before purchase

**DG™** | health

**SEE NEW WARNING  
AND DIRECTIONS**

Compare to the  
active ingredient  
of Imodium® A-D\*

Loperamide Hydrochloride  
Oral Solution 1 mg per 7.5 mL

# Anti-Diarrheal

Controls the Symptoms of Diarrhea  
Anti-Diarrheal Oral Solution

Mint Flavor

4 FL OZ (118mL)

Do not use if  
printed plastic  
neckband is  
broken or missing.

DISTRIBUTED BY  
OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE,  
TN 37072

**100%**  
**Satisfaction  
Guaranteed!**  
(888) 309-9030

\*This product is not  
manufactured or  
distributed by  
Johnson & Johnson  
Consumer Inc.,  
distributor of  
Imodium® A-D.

A1502



**CODE  
AREA**  
• no varnish • no color

: 64526 VT F5

### Drug Facts (continued)

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- fever    ■ mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use  
if you are taking a prescription drug.  
Loperamide may interact with certain  
prescription drugs.

When using this product tiredness,  
drowsiness or dizziness may occur. Be careful  
when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
  - diarrhea lasts for more than 2 days
  - you get abdominal swelling or bulging.
- These may be signs of a serious condition.

If pregnant or breast-feeding, ask a  
health professional before use.

Keep out of reach of children. In case of  
overdose, get medical help or contact a  
Poison Control Center right away.  
(1-800-222-1222)

### Directions

- drink plenty of clear fluids to help  
prevent dehydration caused by  
diarrhea    ■ find right dose on chart.
- If possible, use weight to dose; otherwise  
use age.    ■ shake well before using ▼

**ADHESIVE  
AREA  
NO COPY**

### Drug Facts (continued)

- use only enclosed dosing cup specifically designed for  
use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
children 9-11 years (60-95 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

### Other information

- each 30 mL contains: sodium 15 mg
- store between 20-25°C (68-77°F)

**Inactive ingredients** anhydrous citric acid,  
carboxymethylcellulose sodium, D&C yellow no. 10, FD&C  
blue no. 1, glycerin, microcrystalline cellulose, natural and  
artificial mint flavor, propylene glycol, purified water,  
simethicone, sodium benzoate, sucralose, titanium  
dioxide, xanthan gum

Questions or comments? 1-888-309-9030

**ADHESIVE  
AREA  
NO COPY**

loperamide hcl solution					
Product Information					
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-867		
Route of Administration	ORAL				
Active Ingredient/Active Moiety					
Ingredient Name		Basis of Strength	Strength		
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)		LOPERAMIDE HYDROCHLORIDE	1 mg in 7.5 mL		
Inactive Ingredients					
Ingredient Name			Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)					
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)					
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)					
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)					
GLYCERIN (UNII: PDC6A3C0OX)					
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)					
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)					
WATER (UNII: 059QF0KO0R)					
SODIUM BENZOATE (UNII: OJ245FE5EU)					
SUCRALOSE (UNII: 96K6UQ3ZD4)					
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)					
XANTHAN GUM (UNII: TTV12P4NEE)					
DIMETHICONE (UNII: 92RU3N3Y1O)					
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)					
Product Characteristics					
Color	GREEN (opaque, viscous)	Score			
Shape		Size			
Flavor	MINT	Imprint Code			
Contains					
Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:55910-867-26	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/27/2022		
Marketing Information					
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	

ANDA	ANDA091292	07/27/2022	
------	------------	------------	--

**Labeler -** Dolgencorp Inc (068331990)

Revised: 7/2022

Dolgencorp Inc