

TUSSIN CF- dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Publix Super Markets Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Publix Super Markets, Inc. Tussin CF Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 200 mg

Phenylephrine HCl, USP 10 mg

Purposes

Cough suppressant

Expectorant

Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

Other information

- **each 10 mL contains:** sodium 5 mg
- store at 20-25°C (68-77°F).
Do not refrigerate.

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Principal Display Panel

ADULT tussin CF

COUGH SUPPRESSANT (DEXTROMETHORPHAN HBr)

EXPECTORANT (GUAIFENESIN)

NASAL DECONGESTANT (PHENYLEPHRINE HCl)

Relieves:

Cough

Nasal congestion

Mucus

Compare to the active ingredients in Robitussin® Multi-Symptom Cold

FOR AGES 12 & OVER NON-DROWSY

4 FL OZ (118 mL)

ADULT
NDC 56062-516-26

tussinCF
COUGH SUPPRESSANT
(DEXTROMETHORPHAN HBr)
EXPECTORANT (GUAIFENESIN)
NASAL DECONGESTANT
(PHENYLEPHRINE HCl)

Relieves:

- Cough
- Nasal congestion
- Mucus

FOR AGES 12 & OVER
NON-DROWSY

4 FL OZ (118 mL)

**Compare to the active ingredients in Robitussin® Multi-Symptom Cold*

ADULT
NDC 56062-516-26

tussinCF
COUGH SUPPRESSANT
(DEXTROMETHORPHAN HBr)
EXPECTORANT (GUAIFENESIN)
NASAL DECONGESTANT
(PHENYLEPHRINE HCl)

Relieves:

- Cough
- Nasal congestion
- Mucus

FOR AGES 12 & OVER
NON-DROWSY

4 FL OZ (118 mL)

**Compare to the active ingredients in Robitussin® Multi-Symptom Cold*

ADULT
NDC 56062-516-26

tussinCF
COUGH SUPPRESSANT
(DEXTROMETHORPHAN HBr)
EXPECTORANT (GUAIFENESIN)
NASAL DECONGESTANT
(PHENYLEPHRINE HCl)

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 20 mg.....Cough suppressant
Guaifenesin, USP 200 mg.....Expectorant
Phenylephrine HCl, USP 10 mg.....Nasal decongestant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

Other information

- each 10 mL contains: sodium 5 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

*This product is not manufactured or distributed by the owner of the registered trademark Robitussin®.

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

DISTRIBUTED BY PUBLIX SUPER MARKETS, INC., 3300 PUBLIX CORPORATE PARKWAY, LAKELAND, FL 33811 publix.com 1-888-267-3037

PUBLIX
PUBLIX GUARANTEE: COMPLETE SATISFACTION OR YOUR MONEY BACK

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

LOT NO.
EXP.
: 51424 43 C9

TUSSIN CF

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:56062-516
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	RED	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56062-516-26	1 in 1 CARTON	05/26/2006	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:56062-516-34	1 in 1 CARTON	05/10/2006	
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
---------------------------	---	-----------------------------	---------------------------

OTC monograph final

part341

03/19/1993

Labeler - Publix Super Markets Inc (006922009)

Revised: 10/2018

Publix Super Markets Inc