

ALKA-SELTZER EXTRA STRENGTH- anhydrous citric acid, sodium bicarbonate, and aspirin granule, effervescent
Bayer HealthCare LLC.

Alka-Seltzer Extra Strength

Drug Facts

<i>Active ingredients (in each tablet)</i>	<i>Purpose</i>
Anhydrous citric acid 1000 mg	Antacid
Aspirin 500 mg	Analgesic
Sodium bicarbonate (heat-treated) 1985 mg	Antacid

Uses

for the temporary relief of:

- heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- taking a prescription drug for anticoagulation (blood thinning), diabetes, gout, or arthritis

When using this product do not exceed recommended dosage**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
- feel faint • vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- symptoms get worse or last more than 10 days
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- fully dissolve 2 tablets in 4 ounces of water before taking

adults and children 12 years and over	2 tablets every 6 hours, or as directed by a doctor	do not exceed 7 tablets in 24 hours
	2 tablets every 6	

adults 60 years and over	2 tablets every 6 hours, or as directed by a doctor	do not exceed 3 tablets in 24 hours
children under 12 years	consult a doctor	

Other information

- **each tablet contains:** sodium 588 mg
- store at room temperature. Avoid excessive heat.
- Alka-Seltzer Extra Strength in water contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate

Inactive ingredients

flavor

Questions or comments?

1-800-986-0369 (Mon - Fri 9AM - 5PM EST)

PRINCIPAL DISPLAY PANEL - 24 Tablets Carton

Alka-

Seltzer ®

ANTACID & PAIN RELIEF MEDICINE

FAST

RELIEF

OF:

Acid Indigestion

Upset Stomach

Heartburn with

Headache

Body Aches

XTRA

STRENGTH

24 EFFERVESCENT TABLETS



LIFT HERE

<p>Drug Facts</p> <p>Warnings</p> <p>• heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains</p> <p>• upset stomach with headache from overindulgence in food or drink</p> <p>• headache, body aches, and pain alone</p> <p>Uses</p> <p>• for the temporary relief of heartburn, acid indigestion, and sour stomach</p> <p>• upset stomach with headache or body aches and pains</p> <p>• overindulgence in food or drink</p> <p>• headache, body aches, and pain alone</p> <p>Active Ingredients</p> <p>(in each tablet)</p> <p>Anhydrous citric acid 1000 mg</p> <p>Aspirin 500 mg (NSAID)*</p> <p>Sodium bicarbonate (heat-treated) 1985 mg</p> <p>Antacid</p> <p>Nonsteroidal anti-inflammatory drug</p> <p>Purpose</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older</p> <p>• have had stomach ulcers or bleeding problems</p> <p>• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</p> <p>• use this product</p> <p>• take more or for a longer time than directed</p> <p>Do not use</p> <p>• if you are allergic to aspirin or any other pain reliever/fever reducer</p> <p>• if you have ever had an allergic reaction to this product or any of its ingredients</p> <p>Ask a doctor before use if</p> <p>• you have a history of stomach problems, such as heartburn</p> <p>• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease</p> <p>• you are taking a diuretic</p> <p>• you have asthma</p> <p>• you have a sodium-restricted diet</p> <p>Drug Facts (continued)</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older</p> <p>• have had stomach ulcers or bleeding problems</p> <p>• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</p> <p>• use this product</p> <p>• take more or for a longer time than directed</p> <p>Do not use</p> <p>• if you are allergic to aspirin or any other pain reliever/fever reducer</p> <p>• if you have ever had an allergic reaction to this product or any of its ingredients</p> <p>Ask a doctor before use if</p> <p>• you have a history of stomach problems, such as heartburn</p> <p>• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease</p> <p>• you are taking a diuretic</p> <p>• you have asthma</p> <p>• you have a sodium-restricted diet</p>	<p>Drug Facts</p> <p>Warnings</p> <p>• heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains</p> <p>• upset stomach with headache from overindulgence in food or drink</p> <p>• headache, body aches, and pain alone</p> <p>Uses</p> <p>• for the temporary relief of heartburn, acid indigestion, and sour stomach</p> <p>• upset stomach with headache or body aches and pains</p> <p>• overindulgence in food or drink</p> <p>• headache, body aches, and pain alone</p> <p>Active Ingredients</p> <p>(in each tablet)</p> <p>Anhydrous citric acid 1000 mg</p> <p>Aspirin 500 mg (NSAID)*</p> <p>Sodium bicarbonate (heat-treated) 1985 mg</p> <p>Antacid</p> <p>Nonsteroidal anti-inflammatory drug</p> <p>Purpose</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older</p> <p>• have had stomach ulcers or bleeding problems</p> <p>• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</p> <p>• use this product</p> <p>• take more or for a longer time than directed</p> <p>Do not use</p> <p>• if you are allergic to aspirin or any other pain reliever/fever reducer</p> <p>• if you have ever had an allergic reaction to this product or any of its ingredients</p> <p>Ask a doctor before use if</p> <p>• you have a history of stomach problems, such as heartburn</p> <p>• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease</p> <p>• you are taking a diuretic</p> <p>• you have asthma</p> <p>• you have a sodium-restricted diet</p>
--	--



24 EFFERVESCENT TABLETS
Do not use if pouch is torn or open.



14


PS

<p>Alka-Seltzer</p> <p>EXTRA STRENGTH</p> <p>Fast Relief of: Heartburn Upset Stomach Acid Indigestion with Headache Body Aches</p> <p>LOT & EXP (Bright Background for Black Printing)</p>	<p>Alka-Seltzer</p> <p>EXTRA STRENGTH</p> <p>Fast Relief of: Heartburn Upset Stomach Acid Indigestion with Headache Body Aches</p> <p>Anhydrous citric acid / Antacid Aspirin (NSAID) / Analgesic Sodium bicarbonate / Antacid</p> <p>BAYER</p> <p>24 EFFERVESCENT TABLETS</p>	<p>Alka-Seltzer</p> <p>EXTRA STRENGTH</p> <p>Fast Relief of: Heartburn Upset Stomach Acid Indigestion with Headache Body Aches</p> <p>Place two (2) tablets in 4 oz. of water. Dissolve tablets completely. You do not have to drink any of the residue that may be on the bottom of the glass. The medicines of Alka-Seltzer® are in the water.</p> <p>Take Alka-Seltzer® any time — morning, noon, or night — when you need relief from heartburn, upset stomach, acid indigestion with headache or body aches.</p>
--	--	--

84200913

Bayer
Bayer, the Bayer Cross and Alka-Seltzer are registered trademarks of Bayer.
Whippany, NJ 07981
Bayer Healthcare LLC





ALKA-SELTZER EXTRA STRENGTH

anhydrous citric acid, sodium bicarbonate, and aspirin granule, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0280-0211
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)	ANHYDROUS CITRIC ACID	1000 mg
SODIUM BICARBONATE (UNII: 8MDF5V39QO) (SODIUM CATION - UNII:LYR4M0NH37)	SODIUM BICARBONATE	1985 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg

Inactive Ingredients

Ingredient Name	Strength
LEMON (UNII: 24RS0A988O)	

Product Characteristics

Color	white (White tablet)	Score	no score
Shape	ROUND	Size	25mm
Flavor		Imprint Code	ALKA;SELTZER;EX;S

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-0211-24	12 in 1 CARTON	04/18/2011	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	04/11/2011	

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 12/2023

Bayer HealthCare LLC.