

UNIT DOSE BISACODYL- bisacodyl tablet, delayed release
Cardinal Health 107, LLC

Major 44-327

Active ingredient (in each tablet)

Bisacodyl 5 mg

Enteric Coated Tablets

Purpose

Stimulant laxative

Uses

- relieves occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

Warnings

This Unit Dose package is not child resistant and is Intended for Institutional Use Only. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- you may have stomach discomfort, faintness or cramps

Stop use and ask a doctor if

- have rectal bleeding or fail to have a bowel movement after use of a laxative. These may be signs of a serious condition.
- you need to use a laxative for more than 1 week

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take with a glass of water

adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- avoid excessive humidity

Inactive ingredients

acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, iron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac glaze, simethicone, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

(800) 616-2471

Distributed By:

MAJOR® PHARMACEUTICALS

17177 N Laurel Park Drive, Suite 233

Livonia, MI 48152 USA

Distributed By:

Cardinal Health

Dublin, OH 43017

L5071915-11223


L5071915-21223

Principal Display Panel

Bisacodyl 5 mg

Enteric Coated Tablets

10 Tablets



S105

NDC 55154-6897-0

BISACODYL USP 5 mg
Enteric Coated Tablets

10 TABLETS

Stimulant laxative

Drug Facts
Active ingredient (in each tablet) Purpose
Bisacodyl USP, 5 mg Stimulant laxative

Uses - for relief of occasional constipation and irregularity
 - this product generally produces bowel movement in
 6 to 12 hours

Directions - take with a glass of water
adults and children 12 take 1 to 3 tablets in a
years and over single daily dose
children 6 to under 12 take 1 tablet in a single
years daily dose
children under 6 years ask a doctor

STORAGE: Store at 25° C (77° F); excursions permitted
between 15° - 30° C (59° - 86° F)
- avoid excessive humidity

WARNING: This Unit Dose package is not child resistant
and is Intended for Institutional Use Only. If pregnant or
breast-feeding, ask a health professional before use. Keep out
of reach of children. In case of overdose, get medical help or
contact a Poison Control Center right away.

Questions or comments? (800) 616-2471

Distributed by MAJOR® PHARMACEUTICALS
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152
MFGD By LNK, Hauppauge, NY 11788

Distributed by Cardinal Health
Dublin, OH 43017
L5071915-11223

Lot: Exp:

Distributed by Cardinal Health
Dublin, OH 43017
L5071915-21223

Warnings
Do not use if you cannot swallow without chewing.
Ask a doctor before use if you have
- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks
When using this product
- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- you may have stomach discomfort, faintness or cramps
Stop use and ask a doctor if
- you have rectal bleeding or fail to have a bowel movement after
use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN,
BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

Inactive ingredients: acacia, ammonium hydroxide, calcium
carbonate, carnauba wax, colloidal anhydrous silica, corn starch,
D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake,
hypromellose, iron oxide black, lactose anhydrous, magnesium
stearate, methylparaben, polydextrose, polyethylene glycol,
polyvinyl acetate phthalate, povidone, propylene glycol,
propylparaben, shellac glaze, simethicone, sodium alginate,
sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc,
titanium dioxide, triacetin, triethyl citrate

UNIT DOSE BISACODYL

bisacodyl tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55154-6897(NDC:0904-6407)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	

CALCIUM CARBONATE (UNII: H0G9379FGK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
Polyvinyl Acetate Phthalate (UNII: 58QVG85GW3)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
SHELLAC (UNII: 46N107B71O)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	5
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154-6897-0	10 in 1 BAG	03/25/2002	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	03/25/2002	

Revised: 1/2024

Cardinal Health 107, LLC