

IBUPROFEN- ibuprofen tablet, film coated
COSTCO WHOLESALE CORPORATION

Kirkland 44-393

Active ingredient (in each orange caplet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - menstrual cramps
 - backache
 - the common cold
 - toothache
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- blisters
- skin reddening
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- taking any other drug
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - leg swelling
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 caplet, 2 caplets may be used
 - do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- use by expiration date on package
- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

KIRKLAND

Signature

NDC 63981-393-14

ITM. / ART. 808083

COMPARE TO

MOTRIN® IB

*active ingredient***

IBUPROFEN

IB TABLETS

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

500

Caplets[†]

[†]capsule-shaped tablets

Actual Size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**This product is not manufactured or distributed by
Johnson & Johnson Corporation, owner of the registered
trademark Motrin® IB.

Made in the USA

Manufactured by: LNK INTERNATIONAL, INC.

60 Arkay Drive, Hauppauge, NY 11788 USA

For: Costco Wholesale Corporation,

P.O. Box 34535, Seattle, WA 98124-1535 USA

1-800-774-2678 www.costco.com 20V09016a

50844 REV1221G39314

No print/No varnish area
Lot no/Exp date

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Motrin® IB.

Made in the USA

PEEL HERE FOR MORE DRUG FACTS

KIRKLAND
Signature

COMPARE TO
MOTRIN® IB
active ingredient**

NDC 63981-393-14
ITM. / ART. 808083

**IBUPROFEN
IB TABLETS**

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

500
Caplets[†]

Actual Size

[†]capsule-shaped tablets

Drug Facts

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

Active ingredient (in each orange caplet) **Purpose**
Ibuprofen USP, 200 mg (NSAID)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ headache ■ menstrual cramps
■ backache ■ the common cold
■ toothache ■ muscular aches ■ minor pain of arthritis
■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ blisters ■ skin reddening ■ rash
If an allergic reaction occurs, stop use and seek medical help right away.

Manufactured by: LNK INTERNATIONAL, INC.
60 Arkay Drive, Hauppauge, NY 11788 USA
For: Costco Wholesale Corporation,
P.O. Box 34535, Seattle, WA 98124-1535 USA
1-800-774-2678 www.costco.com
20V09016a

50844 REV1221G39314

PLASTIC BOTTLE

how2recycle.info

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- taking any other drug
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: ■ feel faint
- have bloody or black stools ■ vomit blood
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke: ■ chest pain ■ slurred speech ■ leg swelling ■ trouble breathing
- weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than directed
■ the smallest effective dose should be used
■ adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 caplet, 2 caplets may be used
■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
■ children under 12 years: ask a doctor

Other information

- use by expiration date on package
- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)

Inactive ingredients carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

STOP PEELING

Kirkland 44-393

IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63981-393
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	14mm
Flavor		Imprint Code	44;393
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63981-393-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/1999	
2	NDC:63981-393-58	750 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/1999	05/20/2016
3	NDC:63981-393-03	1 in 1 PACKAGE	03/01/1999	07/27/2021
3		10 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075139	03/01/1999	

Labeler - COSTCO WHOLESALE CORPORATION (103391843)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(63981-393)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(63981-393)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(63981-393)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(63981-393)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(63981-393)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(63981-393)

Revised: 8/2023

COSTCO WHOLESALE CORPORATION