

**NINJACOF-D- chlophedianol hydrochloride, pyrilamine maleate,  
pseudoephedrine hydrochloride liquid  
Centurion Labs, LLC**

-----  
**Ninjacof D**

**Drug Facts**

<b>Active Ingredients (in each 5 mL (1 tsp))</b>	<b>Purpose</b>
Chlophedianol HCl 12.5 mg.....	Cough Suppressant
Pyrilamine Maleate 12.5 mg.....	Antihistamine
Pseudoephedrine HCl 30 mg.....	Nasal Decongestant

**Purpose**

- Antitussive
- Antihistamine
- Nasal Decongestant

**Uses:**

Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies.

- Cough due to minor throat and bronchial irritation
- Runny Nose
- Sneezing
- Itching of the nose or throat
- Itchy, watery eyes
- Nasal Congestion
- Reduces swelling of nasal passages

**WARNINGS**

**Do not exceed recommended dosage**

**Do not use this product if**

- you are now taking prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema
- a cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland
- heart disease
- high blood pressure
- thyroid disease
- diabetes

**Ask doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occurs
- symptoms do not improve within 7 days or are accompanied by a fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition
- new symptoms occur

### **If Pregnant or breast feeding**

Ask a health professional before use.

### **Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

Do not exceed recommended dosage.

---

Adults and Children 12 Years or Older.	5-10ml (1-2 tsp) every 6 to 8 hours not to exceed 40ml in 24hr
Children 6-12	5ml (1 tsp) every 6 to 8 hours not to exceed 20ml in 24hr
Children under 6 years of age	Consult a doctor.

---

### **Other Information**

Store at 59° - 85° F (15° - 30° C)

### **Inactive Ingredients**

Citric Acid, Glycerin, Natural & Artificial Flavors, Propylene Glycol, Sodium Benzoate, Sodium Citrate, Sucralose, Water

### **Questions? Comments?**

To report a serious adverse event or obtain product information, contact 1-866-959-4880

### **PRINCIPAL DISPLAY PANEL**

23359-034-16

**Ninjacof-D™**

Antitussive • Antihistamine • Nasal Decongestant

Each teaspoonful (5 mL) Contains:

Chlophedianol HCl ..... 12.5 mg  
 Pyrilamine Maleate ..... 12.5 mg  
 Pseudoephedrine HCl ..... 30.0 mg

Alcohol Free • Dye Free • Sugar Free

**Cotton Candy Flavor**

**CENTURION**  
 16 fl oz (473 mL)

Manufactured for: Centurion Labs, LLC Birmingham, AL 35243  
 Rev 07/2017

3 2335-9034-16 9

Drug Facts	
<b>Active Ingredients</b> (in each 5 mL (1 tsp))	<b>Purpose</b>
Chlophedianol HCl 12.5 mg	Cough Suppressant
Pyrilamine Maleate 12.5 mg	Antihistamine
Pseudoephedrine HCl 30mg	Nasal Decongestant
<b>Uses:</b> Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies.	
<ul style="list-style-type: none"> <li>Cough due to minor throat and bronchial irritation</li> <li>Runny Nose • Sneezing</li> <li>Itching of the nose or throat</li> <li>Itchy, watery eyes</li> <li>Nasal Congestion</li> <li>Reduces swelling of nasal passages</li> </ul>	
<b>WARNINGS</b> Do not exceed recommended dosage.	
<ul style="list-style-type: none"> <li>Do not use this product if               <ul style="list-style-type: none"> <li>you are now taking prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul> </li> </ul>	
<b>Ask a doctor before use if you have</b> <ul style="list-style-type: none"> <li>a cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema</li> <li>a cough that occurs with too much phlegm (mucus)</li> <li>a breathing problem such as emphysema or chronic bronchitis</li> <li>glaucoma</li> <li>trouble urinating due to an enlarged prostate gland</li> <li>heart disease</li> <li>high blood pressure</li> <li>thyroid disease</li> <li>diabetes</li> </ul>	
<b>Ask doctor or pharmacist before use if you are</b> <ul style="list-style-type: none"> <li>taking sedatives or tranquilizers</li> </ul>	
<b>When using this product</b> <ul style="list-style-type: none"> <li>excitability may occur, especially in children</li> <li>marked drowsiness may occur</li> <li>avoid alcoholic beverages</li> <li>alcohol, sedatives, and tranquilizers may increase drowsiness</li> <li>be careful when driving a motor vehicle or operating machinery</li> </ul>	

Drug Facts (continued)	
<b>Stop use and ask a doctor if</b>	
<ul style="list-style-type: none"> <li>irritation, dizziness, or sleepiness occurs</li> <li>symptoms do not improve within 7 days or are accompanied by a fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition</li> <li>new symptoms occur</li> </ul>	
<b>If pregnant or breast-feeding</b> ask a health professional before use.	
<b>Keep out of reach of children</b> In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> Do not exceed recommended dosage.	
Adults and Children 12 Years or Older:	5-10ml (1-2 tsp) every 6 to 8 hours not to exceed 40ml in 24hr
Children 6-12:	5ml (1 tsp) every 6 to 8 hours not to exceed 20ml in 24hr
Children under 6 years of age:	Consult a doctor.
<b>Other information</b> Store at 59° - 82° F (15° - 30°C)	
<b>Inactive Ingredients</b> Citric Acid, Citrus, Natural & Artificial Flavors, Propylene Glycol, Sodium Benzoate, Sodium Citrate, Sucralose, Water	
<b>Questions? Comments?</b> To report a serious adverse event or obtain product information, contact 1-866-959-4880	

23359-034-16

Ninjacof™ -D

Antitussive, Antihistamine, Nasal Decongestant

Each teaspoonful (5mL) Contains:

Chlophedianol HCl.....12.5 mg

Pyrilamine Maleate.....12.5 mg

Pseudoephedrine HCl.....30.0 mg

Alcohol Free, Dye Free, Sugar Free

Cotton Candy Flavor

Centurion

16 fl oz (473 mL)

Manufactured for: Centurion Labs, LLC Birmingham, AL 35243

Rev 07/2017

**PRINCIPAL DISPLAY PANEL**

23359-034-04

**Ninjacof-D™**

Antitussive • Antihistamine • Nasal Decongestant

Each teaspoonful (5 mL) Contains:

Chlophedianol HCl ..... 12.5 mg  
 Pyrilamine Maleate ..... 12.5 mg  
 Pseudoephedrine HCl ..... 30.0 mg

Alcohol Free • Dye Free • Sugar Free

**Cotton Candy Flavor**

**CENTURION**  
 4 fl oz (120 mL)

Manufactured for: Centurion Labs, LLC Birmingham, AL 35243  
 Rev 10/2018

3 2335-9034-04 6

Drug Facts	
<b>Active Ingredients</b> (in each 5 mL (1 tsp))	<b>Purpose</b>
Chlophedianol HCl 12.5 mg	Cough Suppressant
Pyrilamine Maleate 12.5 mg	Antihistamine
Pseudoephedrine HCl 30mg	Nasal Decongestant
<b>Uses:</b> Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies.	
<ul style="list-style-type: none"> <li>Cough due to minor throat and bronchial irritation</li> <li>Runny Nose • Sneezing</li> <li>Itching of the nose or throat</li> <li>Itchy, watery eyes</li> <li>Nasal Congestion</li> <li>Reduces swelling of nasal passages</li> </ul>	
<b>WARNINGS</b> Do not exceed recommended dosage.	
<ul style="list-style-type: none"> <li>Do not use this product if               <ul style="list-style-type: none"> <li>you are now taking prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul> </li> </ul>	
<b>Ask a doctor before use if you have</b> <ul style="list-style-type: none"> <li>a cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema</li> <li>a cough that occurs with too much phlegm (mucus)</li> <li>a breathing problem such as emphysema or chronic bronchitis</li> <li>glaucoma</li> <li>trouble urinating due to an enlarged prostate gland</li> <li>heart disease</li> <li>high blood pressure</li> <li>thyroid disease</li> <li>diabetes</li> </ul>	
<b>Ask doctor or pharmacist before use if you are</b> <ul style="list-style-type: none"> <li>taking sedatives or tranquilizers</li> </ul>	
<b>When using this product</b> <ul style="list-style-type: none"> <li>excitability may occur, especially in children</li> <li>marked drowsiness may occur</li> <li>avoid alcoholic beverages</li> <li>alcohol, sedatives, and tranquilizers may increase drowsiness</li> <li>be careful when driving a motor vehicle or operating machinery</li> </ul>	

Drug Facts (continued)	
<b>Stop use and ask a doctor if</b>	
<ul style="list-style-type: none"> <li>irritation, dizziness, or sleepiness occurs</li> <li>symptoms do not improve within 7 days or are accompanied by a fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition</li> <li>new symptoms occur</li> </ul>	
<b>If pregnant or breast-feeding</b> ask a health professional before use.	
<b>Keep out of reach of children</b> In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> Do not exceed recommended dosage.	
Adults and Children 12 Years or Older:	5-10ml (1-2 tsp) every 6 to 8 hours not to exceed 40ml in 24hr
Children 6-12:	5ml (1 tsp) every 6 to 8 hours not to exceed 20ml in 24hr
Children under 6 years of age:	Consult a doctor.
<b>Other information</b> Store at 59° - 82° F (15° - 30°C)	
<b>Inactive Ingredients</b> Citric Acid, Citrus, Natural & Artificial Flavors, Propylene Glycol, Sodium Benzoate, Sodium Citrate, Sucralose, Water	
<b>Questions? Comments?</b> To report a serious adverse event or obtain product information, contact 1-866-959-4880	

23359-034-04

Ninjacof™ -D

Antitussive, Antihistamine, Nasal Decongestant

Each Teaspoonful (5 mL) Contains:

Chlohedianol HCl.....12.5 mg

Pyrilamine Maleate.....12.5 mg

Pseudoephedrine HCl.....30.0 mg

Alcohol Free, Dye Free, Sugar Free

Cotton Candy Flavor

Centurion

4 fl oz (120 mL)

Manufactured for: Centurion Labs, LLC Birmingham, AL 35243

Rev. 10/2018

### NINJACOF-D

chlophedianol hydrochloride, pyrilamine maleate, pseudoephedrine hydrochloride liquid

Product Information				
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:23359-034	
<b>Route of Administration</b>	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
PYRILAMINE MALEATE (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)		PYRILAMINE MALEATE	12.5 mg in 5 mL	
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)		PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg in 5 mL	
CHLOPHEDIANOL HYDROCHLORIDE (UNII: 69QQ58998Y) (CHLOPHEDIANOL - UNII:42C50P12AP)		CHLOPHEDIANOL HYDROCHLORIDE	12.5 mg in 5 mL	
Inactive Ingredients				
Ingredient Name			Strength	
CITRIC ACID ACETATE (UNII: DSO12WL7AU)				
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SODIUM CITRATE (UNII: 1Q73Q2JULR)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
WATER (UNII: 059QF0K0OR)				
Product Characteristics				
<b>Color</b>		<b>Score</b>		
<b>Shape</b>		<b>Size</b>		
<b>Flavor</b>	COTTON CANDY	<b>Imprint Code</b>		
<b>Contains</b>				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23359-034-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/16/2017	
2	NDC:23359-034-	120 mL in 1 BOTTLE; Type 0: Not a Combination	11/12/2018	

04

Product

11/12/2010

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/16/2017	

**Labeler** - Centurion Labs, LLC (016481957)

Revised: 1/2024

Centurion Labs, LLC