

**UP AND UP JUNIORS ACETAMINOPHEN- acetaminophen tablet, chewable**  
**Target Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**up and up juniors' acetaminophen melty tabs**

***Active ingredient (in each tablet)***

Acetaminophen 160 mg

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - the common cold
  - flu
  - headache
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has**

liver disease.

**Ask a doctor or pharmacist before use if your child is**

taking the blood thinning drug warfarin.

**When using this product**

**do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or complete warnings for adult use.**
- **do not give more than directed (see overdose warning)**
- find the right dose on chart below. If possible, use weight to dose; otherwise use age.
- dissolve in mouth or chew before swallowing
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

<b>Weight (lb)</b>	under 48	48-59	60-71	72-95	96 and over
<b>Age (yr)</b>	under 6 years	6-8 years	9-10 years	11 years	12 years
<b>Tablets</b>	ask a doctor	2	2 1/2	3	4

**Other information**

- store at between 20°-25°C (68°-77°F). Avoid high humidity.
- see end flap for expiration date and lot number

**Inactive ingredients**

citric acid, corn starch, crospovidone, red no. 27 aluminum lake, dextrose, ethylcellulose, flavors, gelatin, maltodextrin, magnesium stearate, sucralose

**Questions or comments ?**

**1-800-910-6874**

**Principal Display Panel**

NDC 11673-160-31

**Compare to active ingredient in Junior Strength Tylenol® Meltaways®\***

juniors'

acetaminophen

melty tabs,160 mg

pain reliever/fever reducer

aspirin-free

ibuprofen-free

AGES 6-11 YEARS

BUBBLEGUM FLAVOR

24 CHEWABLE TABLETS

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN,  
BROKEN OR SHOWS ANY SIGNS OF TAMPERING

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the  
registered trademark Junior Strength Tylenol® Meltaways®.

50844 REV1013G45008

094 01 0307 R01 ID225450

Distributed by Target Corporation

Minneapolis, MN 55403

© 2014 Target Brands, Inc.

Shop Target.com

# juniors' acetaminophen

mety tabs, 160 mg



Compare to active ingredient in  
Junior Strength Tylenol® Meltaways®

NDC 11673-160-31

# juniors' acetaminophen

mety tabs, 160 mg



pain reliever/fever reducer  
aspirin free  
ibuprofen free

AGES  
**6-11**  
YEARS



**24**  
CHEWABLE  
TABLETS

24 CHEWABLE TABLETS

EXP:

LOT:



1608521011C

### Drug Facts

**Active ingredient (in each tablet)** Acetaminophen 160 mg **Purpose** Pain reliever/fever reducer

**Uses** temporarily relieves minor aches and pains due to:  
 ■ the common cold ■ flu ■ headache  
 ■ temporarily reduces fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount  
 ■ with other drugs containing acetaminophen.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription).  
 If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

### Drug Facts (continued)

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.  
 When using this product do not exceed recommended dose (see overdose warning).

Stop use and ask a doctor if ■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present  
 These could be signs of a serious condition.

Keep out of reach of children. **Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions** ■ this product does not contain directions or complete warnings for adult use.  
 ■ do not give more than directed (see overdose warning)  
 ■ find the right dose on chart below. If possible, use weight to dose; otherwise, use age.  
 ■ dissolve in mouth or chew before swallowing ■ if needed, repeat dose every 4 hours while symptoms last  
 ■ do not give more than 5 times in 24 hours ■ do not give for more than 5 days unless directed by a doctor

Weight (lb)	under 48	48-59	60-71	72-95	96 and over
Age (yr)	under 6 years	6-8 years	9-10 years	11 years	12 years
Tablets	ask a doctor	2	2 1/2	3	4

**Other information** ■ store at between 20°-25°C (68°-77°F). Avoid high humidity.  
 ■ see end flap for expiration date and lot number

**Inactive ingredients** citric acid, corn starch, crospovidone, red no. 27 aluminum lake, dextrose, ethylcellulose, flavors, gelatin, maltodextrin, magnesium stearate, sucralose

**Questions or comments?** 1-800-910-6874

# UP AND UP JUNIORS ACETAMINOPHEN

acetaminophen tablet, chewable

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11673-160
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE (UNII: 68401960MK)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	
DEXTROSE (UNII: IY9XDZ35W2)	
ETHYLCELLULOSES (UNII: 7Z8S9VYZ4B)	
GELATIN (UNII: 2G86QN327L)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

## Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	16mm
<b>Flavor</b>	BUBBLE GUM	<b>Imprint Code</b>	RP160
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-160-31	4 in 1 CARTON		
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph not final	part343	10/19/2015	

