

**EQUATE DIARRHEA CONTROL- loperamide hcl suspension**  
**Wal-Mart Stores Inc**

-----  
**Wal-Mart Diarrhea Control Drug Facts**

**Active ingredient (in each 7.5 mL)**

Loperamide HCl 1 mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
children 9-11 years (60-95 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

### Other information

- **each 30 mL contains:** sodium 15 mg
- store between 20-25°C (68-77°F)

### Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

### Questions or comments?

**1-888-287-1915**

### Package/Label Principal Display Panel

SEE NEW WARNING AND DIRECTIONS

Compare to Imodium® A-D active ingredient

DIARRHEA CONTROL

Loperamide Hydrochloride Oral Solution, 1 mg per 7.5 mL

Anti-Diarrheal

FOR AGES 6 YEARS AND UP

Controls the Symptoms of Diarrhea

Anti-Diarrheal Oral Solution

Mint Flavor

4 FL OZ (118mL)

## Drug Facts

**Active ingredient Purpose**  
(in each 7.5 mL)

Loperamide HCl 1 mg....Anti-diarrheal

**Use** controls symptoms of diarrhea, including Travelers' Diarrhea

### Warnings

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

Peel open to read complete directions and warnings before purchase

SEE NEW WARNING AND DIRECTIONS

**equate™**

NDC 49035-695-26



## DIARRHEA CONTROL

Loperamide Hydrochloride Oral Solution, 1 mg per 7.5 mL

Anti-Diarrheal

FOR AGES 6 YEARS AND UP

Controls the Symptoms of Diarrhea  
Anti-Diarrheal Oral Solution

4 FL OZ (118mL)

Mint Flavor



Do not use if printed plastic neckband is broken or missing

Discard Seals, Empty & Replace Cap



PLASTIC BOTTLE

\*Not recycled in all communities

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716

\*\*This product is not manufactured or distributed by Johnson & Johnson, owner of the registered trademark Imodium®.

164152



: 75826 2E F1

## Drug Facts (continued)

**Do not use** if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use** if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using ▼

## Drug Facts (continued)

- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
children 9-11 years (60-95 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

### Other information

- each 30 mL contains: sodium 15 mg
- store between 20-25°C (68-77°F)

**Inactive ingredients** anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

**Questions or comments?** 1-888-287-1915

## EQUATE DIARRHEA CONTROL

loperamide hcl suspension

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49035-695	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)		LOPERAMIDE HYDROCHLORIDE	1 mg in 7.5 mL	
Inactive Ingredients				
Ingredient Name			Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
GLYCERIN (UNII: PDC6A3C0OX)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
DIMETHICONE (UNII: 92RU3N3Y1O)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
XANTHAN GUM (UNII: TTV12P4NEE)				
Product Characteristics				
Color	GREEN (opaque, viscous))	Score		
Shape		Size		
Flavor	MINT	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-695-26	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/03/2020	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA091292		09/03/2020	

