


OXYBUTYNIN CHLORIDE- oxybutynin chloride syrup
Atlantic Biologicals Corps

OXYBUTYNIN CHLORIDE
SYRUP, USP
5 mg/5 mL

DESCRIPTION

Chemically, oxybutynin chloride is 4-(Diethylamino)-2-butynyl (±)-α-phenylcyclohexaneglycolate hydrochloride. The molecular formula of oxybutynin chloride is $C_{22}H_{31}NO_3 \cdot HCl$. The structural formula appears below:



Oxybutynin chloride is a white crystalline solid with a molecular weight of 393.96. It is readily soluble in water and acids, but relatively insoluble in alkalis.

Each 5 mL, for oral administration, contains 5 mg of oxybutynin chloride. In addition, the following inactive ingredients are present: Artificial raspberry flavor; citric acid, USP; D&C Yellow No. 10; FD&C Blue No. 1; glycerin, USP; liquid sugar; methylparaben, NF; propylene glycol, USP; sodium citrate, USP and sorbitol solution, USP.

Therapeutic Category: Antispasmodic, anticholinergic.

CLINICAL PHARMACOLOGY

Oxybutynin chloride exerts direct antispasmodic effect on smooth muscle and inhibits the muscarinic action of acetylcholine on smooth muscle. Oxybutynin chloride exhibits only one fifth of the anticholinergic activity of atropine on the rabbit detrusor muscle, but four to ten times the antispasmodic activity. No blocking effects occur at skeletal neuromuscular junctions or autonomic ganglia (antinicotinic effects).

Oxybutynin chloride relaxes bladder smooth muscle. In patients with conditions characterized by involuntary bladder contractions, cystometric studies have demonstrated that oxybutynin chloride increases bladder (vesical) capacity, diminishes the frequency of uninhibited contractions of the detrusor muscle, and delays the initial desire to void. Oxybutynin chloride thus decreases urgency and the frequency of both incontinent episodes and voluntary urination.

Oxybutynin chloride was well tolerated in patients administered the drug in controlled studies of 30 days duration and in uncontrolled studies in which some patients received the drug for 2 years. Pharmacokinetic information is not currently available.

INDICATIONS AND USAGE

Oxybutynin chloride syrup is indicated for the relief of symptoms of bladder instability associated with voiding in patients with uninhibited neurogenic or reflex neurogenic bladder (ie, urgency, frequency, urinary leakage, urge incontinence, dysuria).

CONTRAINDICATIONS

Oxybutynin chloride is contraindicated in patients with untreated angle closure glaucoma and in patients with untreated narrow anterior chamber angles since anticholinergic drugs may aggravate these conditions.

It is also contraindicated in partial or complete obstruction of the gastrointestinal tract, paralytic ileus, intestinal atony of the elderly or debilitated patient, megacolon, toxic megacolon complicating ulcerative colitis, severe colitis, and myasthenia gravis. It is contraindicated in patients with obstructive uropathy and in patients with unstable cardiovascular status in acute hemorrhage.

Oxybutynin chloride is contraindicated in patients who have demonstrated hypersensitivity to the product.

WARNINGS

Oxybutynin chloride, when administered in the presence of a high environmental temperature, can cause heat prostration (fever and heat stroke due to decreased sweating). Diarrhea may be an early symptom of incomplete intestinal obstruction, especially in patients with ileostomy, or colostomy. In this instance treatment with oxybutynin chloride would be inappropriate and possibly harmful.

Oxybutynin chloride may produce drowsiness or blurred vision. The patient should be cautioned regarding activities requiring mental alertness such as operating a motor vehicle or other machinery or performing hazardous work while taking this drug.

Alcohol or other sedative drugs may enhance the drowsiness caused by oxybutynin chloride.

PRECAUTIONS

Oxybutynin chloride should be used with caution in the elderly and in all patients with autonomic neuropathy, hepatic or renal disease. Oxybutynin chloride may aggravate the symptoms of hyperthyroidism, coronary heart disease, congestive heart failure, cardiac arrhythmias, hiatal hernia, tachycardia, hypertension, and prostatic hypertrophy.

Administration of oxybutynin chloride to patients with ulcerative colitis may suppress intestinal motility to the point of producing a paralytic ileus and precipitate or aggravate toxic megacolon, a serious complication of the disease.

Carcinogenesis, Mutagenesis, Impairment of Fertility

A 24-month study in rats at dosages up to approximately 400 times the recommended human dosage showed no evidence of carcinogenicity.

Oxybutynin chloride showed no increase of mutagenic activity when tested in *Schizosaccharomyces pompholiciformis*, *Saccharomyces cerevisiae* and *Salmonella typhimurium* test systems. Reproduction studies in the hamster, rabbit, rat, and mouse have shown no definite evidence of impaired fertility.

Pregnancy

Teratogenic Effects-Pregnancy Category B

Reproduction studies in the hamster, rabbit, rat, and mouse have shown no definite evidence of impaired fertility or harm to the animal fetus. The safety of oxybutynin chloride administered to women who are or who may become pregnant has not been established. Therefore, oxybutynin chloride should not be

given to pregnant women unless, in the judgment of the physician, the probable clinical benefits outweigh the possible hazards.

Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when oxybutynin chloride is administered to a nursing woman.

Pediatric Use

The safety and efficacy of oxybutynin chloride administration have been demonstrated for pediatric patients 5 years of age and older (see **DOSAGE AND ADMINISTRATION**). However, as there is insufficient clinical data for pediatric patients under age 5, oxybutynin chloride is not recommended for this age group.

ADVERSE REACTIONS

Following administration of oxybutynin chloride, the symptoms that can be associated with the use of other anticholinergic drugs may occur:

Cardiovascular: Palpitations, tachycardia, vasodilatation

Dermatologic: Decreased sweating, rash

Gastrointestinal/Genitourinary: Constipation, decreased gastrointestinal motility, dry mouth, nausea, urinary hesitance and retention

Nervous System: Asthenia, dizziness, drowsiness, hallucinations, insomnia, restlessness

Ophthalmic: Amblyopia, cycloplegia, decreased lacrimation, mydriasis

Other: Impotence, suppression of lactation

OVERDOSAGE

The symptoms of overdosage with oxybutynin chloride may be any of those seen with other anticholinergic agents. Symptoms may include signs of central nervous system excitation (eg, restlessness, tremor, irritability, convulsions, delirium, hallucinations), flushing, fever, nausea, vomiting, tachycardia, hypotension or hypertension, respiratory failure, paralysis, and coma. In the event of an overdose or exaggerated response, treatment should be symptomatic and supportive. Maintain respiration and induce emesis or perform gastric lavage (emesis is contraindicated in precomatose patients, convulsive or psychotic state). Activated charcoal may be administered as well as a cathartic. Physostigmine may be considered to reverse symptoms of anticholinergic intoxication. Hyperpyrexia may be treated symptomatically with ice bags or other cold applications and sponges.

DOSAGE AND ADMINISTRATION

Adults

The usual dose is one teaspoonful (5 mg/5 mL) syrup two to three times a day. The maximum recommended dose is one teaspoonful (5 mg/5 mL) syrup four times a day.

Children over 5 years of age

The usual dose is one teaspoonful (5 mg/5 mL) two times a day. The maximum recommended dose is one teaspoonful (5 mg/5 mL) three times a day.

HOW SUPPLIED

Product: 17856-0092

NDC: 17856-0092-1 5 mL in a CUP

Rx Only

Product No.: 8092

Manufactured By:
Morton Grove Pharmaceuticals, Inc.
Morton Grove, IL 60053

A50-8092-16

REV. 11-04


OXYBUTYNIN CHLORIDE SYRUP

17856-0092-01
OXYBUTYNIN CHLORIDE
SYRUP, USP



See package insert for indications and dosage schedule

STORE AT CONTROLLED ROOM
TEMPERATURE 15°-30°C (59°-86°F). *****
Keep this and all Medication out of the reach of
childrer.



17856-0092-01

Dosage: 5 mg/ 5 ML

OXYBUTYNIN CHLORIDE

Qty: 72 CUPS



GTIN: 00117856009214

S/N: 01280901

Exp: 07/26/21

Lot: 012809

RX

Packaged by: Unit Dose Solutions
Morrisville, NC 27560

Distributed by: Atlantic Biologicals Corp.
Miami FL 33176

Rev. 09/13

Call to Reorder: 800.509.7592

OXYBUTYNIN CHLORIDE

oxybutynin chloride syrup

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:17856-0092(NDC:60432-092)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYBUTYNIN CHLORIDE (UNII: L9F3D9RENQ) (OXYBUTYNIN - UNII:K9P6MC7092)	OXYBUTYNIN CHLORIDE	5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
Water (UNII: 059QF0KO0R)	
Trisodium Citrate Dihydrate (UNII: B22547B95K)	
Anhydrous Citric Acid (UNII: XF417D3PSL)	
Sucrose (UNII: C151H8M554)	
Sorbitol (UNII: 506T60A25R)	
Glycerin (UNII: PDC6A3C0OX)	
Propylene Glycol (UNII: 6DC9Q167V3)	
Methylparaben (UNII: A218C7HI9T)	
FD&C Blue No. 1 (UNII: H3R47K3TBD)	
D&C Yellow No. 10 (UNII: 35SW5USQ3G)	

Product Characteristics

Color	BLUE, GREEN	Score	
Shape		Size	
Flavor	RASPBERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17856-0092-1	72 in 1 CASE	01/27/2021	
1		5 mL in 1 CUP; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074868	02/12/1997	

Labeler - Atlantic Biologicals Corps (047437707)

Establishment

Name	Address	ID/FEI	Business Operations
Atlantic Biologicals Corps		047437707	RELABEL(17856-0092) , REPACK(17856-0092)

Revised: 1/2021

Atlantic Biologicals Corps