

**GOOD NEIGHBOR PHARMACY LOPERAMIDE HYDROCHLORIDE- loperamide  
hcl solution**

**Amerisource Bergen**

-----

**Amerisource Bergen Loperamide Hydrochloride Oral Solution, 1 mg per 7.5 mL Drug Facts**

**Active ingredient (in each 7.5 mL)**

Loperamide HCl 1 mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
children 9-11 years (60-95 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

## Other information

- **each 30 mL contains:** sodium 15 mg
- store between 20-25°C (68-77°F)

## Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

## Questions or comments?

**1-800-719-9260**

## Package/Label Principal Display Panel

Compare to Imodium® A-D active ingredient

See New Warning and Directions

Loperamide Hydrochloride Oral Solution, 1 mg per 7.5 mL

anti-diarrheal

Controls the Symptoms of Diarrhea

mint flavor

anti-diarrheal oral solution

4 fl oz (120 mL)

## Drug Facts

**Active ingredient Purpose**  
(in each 7.5 mL)  
Loperamide HCl 1 mg....Anti-diarrheal

**Use** controls symptoms of diarrhea, including Travelers' Diarrhea

### Warnings

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl  
**Heart alert:** Taking more than directed can cause serious heart problems or death

Peel open to read complete directions and warnings before purchase



Compare to  
Imodium® A-D  
active ingredient\*

NDC 46122-544-26

See New Warning and Directions

# Loperamide Hydrochloride

Oral Solution, 1 mg per 7.5 mL anti-diarrheal

Controls the Symptoms of Diarrhea

mint flavor  
anti-diarrheal oral solution

4 fl oz (120 mL)

Do not use if printed plastic neckband is broken or missing.

Distributed By  
AmerisourceBergen  
1300 Morris Drive  
Chesterbrook,  
PA 19087  
Questions or  
Concerns?  
[www.mygnp.com](http://www.mygnp.com)

**GLUTEN FREE**

\*Good Neighbor Pharmacy®  
Loperamide Hydrochloride  
Oral Solution, 1 mg  
per 7.5 mL is not  
manufactured or  
distributed by Johnson &  
Johnson Consumer Inc.,  
distributor of  
Imodium® A-D.

ABC#: 10188131



: 64526 29 F4

## Drug Facts (continued)

**Do not use** if you have bloody or black stool

**Ask a doctor before use if you have**

- fever ■ mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

■ **drink plenty of clear fluids to help prevent dehydration caused by diarrhea** ■ find right dose on chart. If possible, use weight to dose; otherwise use age. ■ shake well before using ▼

## Drug Facts (continued)

■ use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.  
■ mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
children 9-11 years (60-95 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

### Other information

- each 30 mL contains: sodium 15 mg
- store between 20-25°C (68-77°F)

**Inactive ingredients** anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

Questions or comments? 1-800-719-9260

# GOOD NEIGHBOR PHARMACY LOPERAMIDE HYDROCHLORIDE

loperamide hcl solution

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:46 122-544
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	1 mg in 7.5 mL

## Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XANTHAN GUM (UNII: TTV12P4NEE)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

## Product Characteristics

<b>Color</b>	GREEN (opaque, viscous)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MINT	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46 122-544-26	120 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/13/2018	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091292	08/13/2018	

**Labeler** - Amerisource Bergen (007914906)

Revised: 1/2020

Amerisource Bergen