

CHILDRENS PLUS MULTI SYMPTOM COLD- acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hydrochloride suspension
Best Choice (Valu Merchandisers Company)

Drug Facts

Active ingredients (in each 5 mL)

Acetaminophen 160 mg

Chlorpheniramine maleate 1 mg

Dextromethorphan HBr 5 mg

Phenylephrine HCl 2.5 mg

Purposes

Pain reliever/fever reducer

Antihistamine

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves the following cold/flu symptoms:
 - minor aches and pains
 - sore throat
 - cough
 - sneezing and runny nose
 - headache
 - nasal congestion
 - stuffy nose
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses (10 mL) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- to make a child sleepy
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the child has

- liver disease
- glaucoma
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

Ask a doctor or pharmacist before use if your child is taking

- sedatives or tranquilizers
- the blood thinning drug warfarin

When using this product

- **do not exceed recommended dose (see overdose warning)**
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or last more than 3 days
- nervousness, dizziness or sleeplessness occurs
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adult as well as children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- mL=milliliter
- shake well before using
- do not give more than 5 doses in any 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose, otherwise, use age.

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

Other Information

- store between 20-25°C (68-77°F). Do not refrigerate

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C red #33, FD&C blue #1, FD&C red #40, flavor, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

For Ages 6-11 Years

COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S TYLENOL® PLUS MULTI-SYMPTOM COLD*

Children's Plus Multi-Symptom Cold

Acetaminophen

- Pain Reliever / Fever Reducer

Chlorpheniramine Maleate

- Antihistamine

Dextromethorphan HBr

- Cough Suppressant

Phenylephrine HCl

- Nasal Decongestant

Grape Flavor

Oral Suspension

FL OZ (mL)

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Plus Multi-Symptom Cold.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

PROUDLY DISTRIBUTED BY:

VALU MERCHANDISERS, CO.

5000 KANSAS AVE

KANSAS CITY, KS 66106

www.bestchoicebrand.com

Package Label

Drug Facts (continued)

- nervousness, dizziness or sleeplessness occurs
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- mL = milliliter
- shake well before using
- do not give more than 5 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

Other information

- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients: carboxymethylcellulose sodium, citric acid, D&C red #33, FD&C blue #1, FD&C red #40, flavor, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Plus Multi-Symptom Cold.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

PARENTS:
Learn about your medicine at
www.StopMedicineAbuse.org

Best Choice,
100% Guaranteed
www.bestchoicebrand.com

PROUDLY DISTRIBUTED BY:
VALU MERCHANDISERS, CO.
5000 KANSAS AVE
KANSAS CITY, KS 66106

For Ages 6 to 11 Years

Best Choice®

COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S TYLENOL® PLUS MULTI-SYMPTOM COLD*

Children's PLUS MULTI-SYMPTOM COLD

Acetaminophen

- Pain Reliever / Fever Reducer

Chlorpheniramine Maleate

- Antihistamine

Dextromethorphan HBr

- Cough Suppressant

Phenylephrine HCl

- Nasal Decongestant

GRAPE FLAVOR

Oral Suspension
4 FL OZ (118 mL)

For Ages 6 to 11 Years

Best Choice®

COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S TYLENOL® PLUS MULTI-SYMPTOM COLD*

Children's PLUS MULTI-SYMPTOM COLD

Acetaminophen

- Pain Reliever / Fever Reducer

Chlorpheniramine Maleate

- Antihistamine

Dextromethorphan HBr

- Cough Suppressant

Phenylephrine HCl

- Nasal Decongestant

GRAPE FLAVOR

Oral Suspension
4 FL OZ (118 mL)

Drug Facts

Active ingredients (in each 5 mL)

Acetaminophen 160 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 1 mg	Antihistamine
Dextromethorphan HBr 5 mg	Cough suppressant
Phenylephrine HCl 2.5 mg	Nasal decongestant

Uses

- temporarily relieves the following cold/flu symptoms
- minor aches and pains
- sore throat
- cough
- sneezing and runny nose
- headache
- nasal congestion
- stuffy nose
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses (10 mL) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- to make a child sleepy
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has

- liver disease
- glaucoma
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

Ask a doctor or pharmacist before use if your child is taking

- the blood thinning drug warfarin
- sedatives or tranquilizers

When using this product

- do not exceed recommended dose (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

BEST CHOICE

0 70038 61689 4

PLD-H253C FC004114

Lot No.:
Exp. Date:

BEST CHOICE Children's Plus Multi-Symptom Cold

CHILDRENS PLUS MULTI SYMPTOM COLD

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hydrochloride suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-304
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	1 mg in 5 mL

DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII: 7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-304-04	1 in 1 BOX	05/31/2015	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/31/2015	

Labeler - Best Choice (Valu Merchandisers Company) (868703513)