

**IBUPROFEN- ibuprofen capsule, liquid filled**  
**The Kroger Co.**

-----

**DRUG FACTS**

**Active ingredient (in each capsule)**

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*

(present as the free acid and potassium salt)

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/ fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- headache
- backache
- the common cold
- minor pain of arthritis
- temporarily reduces fever
- toothache
- menstrual cramps
- muscular aches

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery

**Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- **each capsule contains:** potassium 20 mg

- read all warnings and directions before use.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F).

**Inactive ingredients**

FD&C blue #1, gelatin, polyethylene glycol, potassium hydroxide, purified water, sorbitol

**Questions or Comments ?**

Call toll free **1-800-632-6900**

**Principal Display Panel**

Compare to the active ingredient in Advil® Liqui- Gels® †See bottom panel

**Important: Read all product information before using**

EASY OPEN CAP

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

Ibuprofen Capsules 200 mg

Pain reliever/ fever reducer (NSAID)

160 CAPSULES\*\*

soft gelatin capsules\*\*

\*\* (Liquid filled capsules)

† Advil® is a registered trademark of pfizer consumer healthcare, Madison, NJ 07940. Pfizer consumer healthcare is not affiliated with the Kroger co. or this product

**DISTRIBUTED BY THE KROGER CO.**

**CINCINNATI, OHIO 45202**

**Do not use if seal under cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.**

**Product Label**



**Compare to the Active Ingredient in ADVIL® Liquid-Gels®**  
\*See Bottom of Label

NDC 30142-123-16

**Important: Read all product information before using**

**THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN**

**Easy Open Cap**

**Ibuprofen**  
**Capsules 200 mg**

**Pain Reliever/Fever Reducer (NSAID)**

**160 Capsules**  
**SOFT GELATIN CAPSULES\*\***  
**\*\*(LIQUID FILLED CAPSULES)**

**Do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**

**Drug Facts**

**Active ingredient (in each capsule)**  
Solubilized ibuprofen equal to 200 mg (ibuprofen (NSAID))  
(present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**Purpose**  
Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Drug Facts (continued)**

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product:**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - stomach pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Other information**

- each capsule contains: potassium 20 mg
- read all warnings and directions before use.
- store at 20° to 25° C (68° to 77° F)
- avoid excessive heat above 40° C (104° F)

**Inactive ingredients:** FD&C Blue #1, gelatin, polyethylene glycol, potassium hydroxide, purified water, sorbitol

**Questions or comments?** Call toll free 1-800-632-6900

**Drug Facts (continued)**

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product:**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - stomach pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Lot No.:** 041260135227

**Exp. Date:**

**PLD-A** 1B000679

**Rev. 03-2012**

**QUALITY GUARANTEED**  
If you are not completely satisfied with this product, return it for a refund or replacement.

**Comments or questions?** [www.Kroger.com](http://www.Kroger.com)

**Peel Here**

## Ibuprofen capsules 200 mg

### IBUPROFEN

ibuprofen capsule, liquid filled

#### Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:30 142-123
Route of Administration	ORAL	DEA Schedule	

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (IBUPROFEN)	IBUPROFEN	200 mg

#### Inactive Ingredients

Ingredient Name		Strength		
FD&C BLUE NO. 1				
GELATIN				
POLYETHYLENE GLYCOLS				
POTASSIUM HYDROXIDE				
WATER				
SORBITOL				
Product Characteristics				
Color	BLUE	Score	no score	
Shape	CAPSULE	Size	19mm	
Flavor		Imprint Code	A356	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30 142-123-16	1 in 1 BOX		
1		160 in 1 BOTTLE, PLASTIC		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA202300	07/12/2012		

**Labeler** - The Kroger Co. (006999528)