

HARRIS TEETER TUSSIN- dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Harris Teeter, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Harris Teeter, LLC Tussin Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 200 mg

Phenylephrine HCl, USP 10 mg

Purposes

Cough suppressant

Expectorant

Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

Other information

- each 10 mL contains: sodium 5 mg
- store at 20-25°C (68-77°F).

Do not refrigerate.

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredients in Robitussin® Multi-Symptom Cold

Adult

Tussin

Cough Suppressant (Dextromethorphan HBr)

Expectorant (Guaifenesin)

Nasal Decongestant (Phenylephrine HCl)

MULTI-SYMPTOM COLD

Relieves:

Cough

Nasal Congestion


Mucus

Non-Drowsy

CF

For Ages 12 & Over

8 FL OZ (237 mL)



Compare to the active ingredients in Robitussin® Multi-Symptom Cold*



Compare to the active ingredients in Robitussin® Multi-Symptom Cold*



Adult
Tussin

**Cough Suppressant (Dextromethorphan HBr)
Expectorant (Guaifenesin)
Nasal Decongestant (Phenylephrine HCl)**

**MULTI-SYMP TOM
COLD**

Relieves:

- Cough
- Nasal Congestion
- Mucus

Non-Drowsy

CF

For Ages 12 & Over



8 FL OZ (237 ml)

Adult
Tussin

**Cough Suppressant (Dextromethorphan HBr)
Expectorant (Guaifenesin)
Nasal Decongestant (Phenylephrine HCl)**

**MULTI-SYMP TOM
COLD**

Relieves:

- Cough
- Nasal Congestion
- Mucus

Non-Drowsy

CF


For Ages 12 & Over



8 FL OZ (237 ml)

ST L 02 (207 HIL)

ST L 02 (207 HIL)



Cough Suppressant (Dextromethorphan HBr)
Expectorant (Guaifenesin)
Nasal Decongestant (Phenylephrine HCl)

Tussin

Adult



Drug Facts

Active ingredients Purposes (in each 10 mL)

Dextromethorphan HBr,
USP 20 mg.....Cough suppressant
Guaifenesin, USP 200 mg.....Expectorant
Phenylephrine HCl,
USP 10 mg.....Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition. ▶

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

Other information

- each 10 mL contains: sodium 5 mg
- store at 20-25°C (68-77°F).
Do not refrigerate.

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

1-800-719-9260

*This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Multi-Symptom Cold.

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

PROUDLY DISTRIBUTED BY:
HARRIS TEETER, LLC, MATTHEWS, NC 28105
 **yourplus** guarantee.

1-800-432-6111 or harristeeter.com

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



LOT NO.

EXP.

151634 HT C2

HARRIS TEETER TUSSIN

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69256-516
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	RED (clear)	Score	
Shape		Size	
Flavor	FRUIT (Menthol)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69256-516-34	1 in 1 CARTON	11/21/2015	
1		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	11/21/2015	

Labeler - Harris Teeter, LLC (048463103)

Revised: 12/2019

Harris Teeter, LLC