

LEADER TUSSIN DM- dextromethorphan hydrobromide and guaifenesin liquid

Cardinal Health

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Cardinal Health Tussin DM Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 200 mg

Purposes

Cough suppressant

Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, FD&C red no. 40, flavor, glycerin, high fructose corn syrup, menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Robitussin® Cough + Chest Congestion DM active ingredients

For Ages 12 & Over

Adult

Tussin DM

Cough Suppressant (Dextromethorphan HBr)

Expectorant (Guaifenesin)

Cough + Chest Congestion DM

Relieves:

Cough

Mucus

Non-Drowsy

Alcohol Free

4 FL OZ (118 mL)



LEADER®

NDC 37205-970-26

**Compare to Robitussin®
Cough + Chest
Congestion DM**
active ingredients*

For Ages 12 & Over

Adult Tussin DM

*Cough Suppressant (Dextromethorphan HBr)
Expectorant (Guaifenesin)*

*Cough + Chest
Congestion DM*

Relieves:

Cough

Mucus

Non-Drowsy



Alcohol Free



4 FL OZ (118 ml)

LEADER®

NDC 37205-970-26

**Compare to Robitussin®
Cough + Chest
Congestion DM**
active ingredients*

For Ages 12 & Over

Adult Tussin DM

*Cough Suppressant (Dextromethorphan HBr)
Expectorant (Guaifenesin)*

*Cough + Chest
Congestion DM*

Relieves:

Cough

Mucus


Non-Drowsy



Alcohol Free



4 FL OZ (118 ml)



DO NOT USE IF PRINTED NECKBAND
IS BROKEN OR MISSING

Cough Suppressant (Dextromethorphan HBr)
Expectorant (Guaifenesin)

**Adult
Tussin
DM**



Drug Facts

Active ingredients **Purposes**
(in each 10 mL)
Dextromethorphan HBr,
USP 20 mg.....Cough suppressant
Guaifenesin, USP 200 mg.....Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

Drug Facts (continued)

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, FD&C red no. 40, flavor, glycerin, high fructose corn syrup, menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose

Questions or comments?

1-800-719-9260

Gluten Free

*This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Cough + Chest Congestion DM.

All Leader® Brand products are 100% satisfaction guaranteed or return to place of purchase for a full refund.

DISTRIBUTED BY CARDINAL HEALTH
DUBLIN, OHIO 43017
CIN 1388081
www.myleader.com
1-800-200-6313

Use Dosing Cup Included

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



LEADER TUSSIN DM

dextromethorphan hydrobromide, guaifenesin liquid

Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:37205-970
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (DEXTROMETHORPHAN)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
GUAIFENESIN (GUAIFENESIN)	GUAIFENESIN	200 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID	
FD&C RED NO. 40	
GLYCERIN	
HIGH FRUCTOSE CORN SYRUP	
MENTHOL	
PROPYLENE GLYCOL	
WATER	
SODIUM BENZOATE	
SODIUM CITRATE	
SUCRALOSE	

Product Characteristics

Color	RED (Orange-Red)	Score	
--------------	------------------	--------------	--

Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37205-970-26	1 in 1 CARTON		
1		118 mL in 1 BOTTLE		
2	NDC:37205-970-34	1 in 1 CARTON		
2		237 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/21/1991	

Labeler - Cardinal Health (097537435)