

ASPIRIN CHEWABLE- aspirin 81 mg tablet
McKesson Corporation dba SKY Packaging

Drug Facts

Active ingredient

Aspirin 81 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include hives, facial swelling, asthma (wheezing), shock.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use

if you are allergic to aspirin or any other pain reliever/fever reducer this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away
- you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep Out of Reach of Children

In case of accidental overdose, get medical help or contact a Poison Control Center(1-800-222-1222) immediately.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Inactive Ingredients

corn starch, dextrans, FD&C yellow #6 aluminum lake, orange flavor, saccharin sodium

HOW SUPPLIED

Product: 63739-434

NDC: 63739-434-01 30 TABLET in a BLISTER PACK / 25 in a BOX, UNIT-DOSE

Drug Facts

Drug Facts

Active ingredient (in each tablet) Purpose

Aspirin 81 mg (NSAID*) Pain Reliever

*nonsteroidal anti-inflammatory drug

Uses: temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness

Allergy alert: Aspirin may cause a severe allergic reaction which may include hives, facial swelling, asthma (wheezing), shock.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer • this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for • gout • diabetes • arthritis

Stop use and ask a doctor if • an allergic reaction occurs. Seek medical help right away • you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better • pain gets worse or lasts more than 10 days • redness or swelling is present • new symptoms occur • ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep Out of Reach of Children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) immediately.

Directions • drink a full glass of water with each dose • adults and children 12 years and over: take 4 to 6 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor • children under 12 years: consult a doctor

Other information • save carton for full directions and warnings • store at room temperature • do not use if blister card is damaged or appears to have been tampered

Inactive ingredients: corn starch, dextrates, FD&C yellow #6, aluminum lake, orange flavor, saccharin sodium

Mfg. By: Allegiant Health, Deer Park, NY 11729

Dist. By: McKesson Corporation, Memphis, TN 38141

†This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Bayer®



ASPIRIN CHEWABLE

NDC 63739-434-02



Tablets, 81 mg (NSAID*)
UD 300 Tablets (30x10)

LOT 1234567890

EXP YYYY-MM-DD



(01)10363739434025

91055
R-00

00000

ASPIRIN CHEWABLE

aspirin 81 mg tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:63739-434(NDC:69168-288)

Route of Administration	ORAL
-------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
DEXTRATES (UNII: G263MI44RU)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	7mm
Flavor	ORANGE	Imprint Code	AZ013
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63739-434-01	25 in 1 BOX, UNIT-DOSE	08/02/2016	05/31/2021
1		30 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:63739-434-02	30 in 1 BOX	03/18/2021	
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/18/2014	

Labeler - McKesson Corporation dba SKY Packaging (140529962)

Revised: 2/2024

McKesson Corporation dba SKY Packaging