

**CHLORASEPTIC SORE THROAT MAX LIQUID CENTER- menthol and benzocaine lozenge  
Prestige Brands Holdings, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Chloraseptic Max Berry Lozenges**

**Drug Facts**

**Active Ingredients**

**(in each lozenge)**

Benzocaine 15 mg

**Purpose**

Oral Anesthetic/Analgesic

**Active Ingredients**

**(in each lozenge)**

Menthol 10 mg

**Purpose**

Oral Anesthetic/Analgesic

**Uses**

Temporarily relieves:

- occasional minor irritation, pain, sore throat and sore mouth

**Warnings**

**Allergy Alert:** Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other “caine” anesthetics.

**Sore Throat Warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea and vomiting may be serious. Consult a doctor promptly. Do not use more than 2 days or administer to children under 6 years of age unless directed by a doctor.

**When using this product:**

do not exceed recommended dosage.

**Stop use and ask a dentist or doctor if:**

- sore mouth symptoms do not improve in 7 days
- irritation, pain or redness persists or worsens
- swelling, rash or fever develops

**If pregnant or breast-feeding,**

ask a health care professional before use.

**Keep out of reach of children.**

In case of overdose or accidental poisoning, get medical help or contact a Poison Control Center right away.

**Directions**

- **Adults and children 6 years of age and over:** Take 1 lozenge every 2 hours as needed. Allow lozenges to dissolve slowly in the mouth.
- **Children under 6 years of age:** Consult a doctor or dentist.

**Other Information**

- Store between 59° F and 86° F (15° C - 30° C).
- Protect from moisture.
- Check expiration date before using.
- **Tamper Evident:** Do not use if blister package with “Chloraseptic®” name has been disturbed or opened.

**Inactive Ingredients**

Corn Syrup, FD&C Blue #1, FD&C Red #40, Flavor, Glycerine, Propylene Glycol, Soy Lecithin, Sucrose, Water

**Questions?**

**1-800-552-7932 or [www.chloraseptic.com](http://www.chloraseptic.com)**

**PRINCIPAL DISPLAY PANEL**

Chloraseptic® Max  
SORE THROAT FAST ACTING RELIEF  
15 lozenges | Wild Berries

SOOTHING  
LIQUID CENTER

Pharmacist #1 Recommended  
**Chloraseptic**<sup>®</sup>

Menthol | Benzocaine Oral Anesthetic | Analgesic Lozenges

**Max**

**SORE THROAT**  
FAST ACTING RELIEF



- **MAX Strength** pain relief
- Cools nasal passages

15 lozenges | Wild Berries

### Drug Facts

#### Active Ingredients (in each lozenge) Purpose

Benzocaine 15 mg ..... Oral Anesthetic/Analgesic  
Menthol 10 mg ..... Oral Anesthetic/Analgesic

**Uses** Temporarily relieves: ■ occasional minor irritation, pain, sore throat and sore mouth

#### Warnings

**Allergy Alert:** Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

**Sore Throat Warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult a doctor promptly. Do not use more than 2 days or administer to children under 6 years of age unless directed by a doctor.

**When using this product:** do not exceed recommended dosage.

#### Stop use and ask a dentist or doctor if:

- sore mouth symptoms do not improve in 7 days
- irritation, pain or redness persists or worsens
- swelling, rash or fever develops

### Drug Facts (continued)

If pregnant or breast-feeding, ask a health care professional before use.

#### Keep out of reach of children.

In case of overdose or accidental poisoning, get medical help or contact a Poison Control Center right away.

#### Directions

- **Adults and children 6 years of age and over:** Take 1 lozenge every 2 hours as needed. Allow lozenges to dissolve slowly in the mouth.
- **Children under 6 years of age:** Consult a doctor or dentist.

#### Other Information

- Store between 59°F and 86°F (15°C -30°C). ■ Protect from moisture. ■ Check expiration date before using.
- **Tamper Evident:** Do not use if blister package with "Chloraseptic<sup>®</sup>" name has been disturbed or opened.

**Inactive Ingredients** Corn Syrup, FD&C Blue #1, FD&C Red #40, Flavor, Glycerine, Propylene Glycol, Soy Lecithin, Sucrose, Water

#### Questions?

1-800-552-7932 or [www.chloraseptic.com](http://www.chloraseptic.com)

CHLORASEPTIC SORE THROAT MAX LIQUID CENTER

menthol and benzocaine lozenge

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67172-174
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	10 mg
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	15 mg

### Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
CORN SYRUP (UNII: 9G5L16BK6N)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
SUCROSE (UNII: C151H8M554)	
WATER (UNII: 059QF0KO0R)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

### Product Characteristics

<b>Color</b>	RED	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	4mm
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	C
<b>Contains</b>			

### Packaging

#	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:67172-174-61	1 in 1 CARTON		
1		15 in 1 BLISTER PACK		

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC MONOGRAPH FINAL	part341	08/01/2010	

**Labeler** - Prestige Brands Holdings, Inc. (159655021)

**Establishment**

Name	Address	ID/FEI	Business Operations
BestSweet, Inc.		002149 136	MANUFACTURE

Revised: 11/2010

Prestige Brands Holdings, Inc.